MASTER

SYDNEY GRAMMAR SCHOOL



2016 Half-Yearly Examination

FORM V

MATHEMATICS 2 UNIT

Tuesday 17th May 2016

General Instructions

- Writing time 1 hour 30 minutes
- Write using black pen.
- Board-approved calculators and templates may be used.

Total – 80 Marks

• All questions may be attempted.

Section I – 8 Marks

- Questions 1-8 are of equal value.
- Record your answers to the multiple choice on the sheet provided.

Section II – 72 Marks

- Questions 9–14 are of equal value.
- All necessary working should be shown.
- Start each question in a new booklet.

5A:	DNW	5B:	$\mathbf{P}\mathbf{K}\mathbf{H}$
5E:	WJM	$5\mathrm{F}$:	GMC
5P:	TCW	5Q:	SDP

Checklist

- SGS booklets 6 per boy
- Multiple choice answer sheet
- Candidature 178 boys

Collection

- Write your name, class and Master on each answer booklet and on your multiple choice answer sheet.
- Hand in the booklets in a single wellordered pile.
- Hand in a booklet for each question in Section II, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Write your name, class and Master on this question paper and hand it in with your answers.
- Place everything inside the answer booklet for Question Nine.

5C: LRP	5D: FMW
5G: NL	5H: SO
5R: RCF	

Examiner NL

SECTION I - Multiple Choice

Answers for this section should be recorded on the separate answer sheet handed out with this examination paper.

QUESTION ONE

Which of the following bearings is due west?


- (A) $000^{\circ} T$
- (B) 090° T
- (C) 180° T
- (D) 270° T

QUESTION TWO

What is the correct expansion of (x-1)(x+4)?

(A) $x^{2} - 3x + 4$ (B) $x^{2} + 4x - 4$ (C) $x^{2} + 3x - 4$ (D) $x^{2} + 4x - 3$

QUESTION THREE




Which expression is the exact area of the triangle above?

(A) $5\sqrt{2}$ (B) $\frac{20\sqrt{2}}{3}$ (C) $\frac{15\sqrt{3}}{2}$ (D) $10\sqrt{2}$

Examination continues next page ...

QUESTION FOUR



The equation of the circle drawn above is:

- (A) $(x+2)^2 + (y+3)^2 = 6$
- (B) $(x-3)^2 + (y-2)^2 = 6$
- (C) $(x+3)^2 + (y+2)^2 = 36$
- (D) $(x-2)^2 + (y-3)^2 = 36$

QUESTION FIVE


Which expression is equivalent to $x^2 + 4x + 9$?

(A)
$$(x + 2)^2 + 13$$

(B) $(x - 2)^2 + 5$
(C) $(x + 2)^2 + 9$
(D) $(x + 2)^2 + 5$

Examination continues overleaf

QUESTION SIX



Using the graph above, the solution to the inequation $x^2 - x - 2 > 0$ is:

(A) $-1 \le x \le 2$ (B) x < -1 or x > 2(C) -1 > x > 2(D) x < -1 or x < 2

QUESTION SEVEN

Which of these functions is even?

(A)
$$f(x) = x^2 - 8x + 15$$

(B) $f(x) = (x+2)^2 - 4$
(C) $f(x) = \sqrt{9 - x^2}$
(D) $f(x) = x^3$

QUESTION EIGHT

Which of the following identities is <u>not</u> true?

(A)
$$\sec^2 \theta + 1 = \cot^2 \theta$$

(B)
$$\sin^2 \theta + \cos^2 \theta = 1$$

(C)
$$1 + \cot^2 \theta = \csc^2 \theta$$

(D)
$$\tan^2 \theta + 1 = \sec^2 \theta$$

End of Section I

Examination continues next page ...

SECTION II - Written Response

Answers for this section should be recorded in the booklets provided.

Show all necessary working.

Start a new booklet for each question.

QUESTION NINE (12 marks) Use a separate writing booklet.

- (a) If f(x) = 4x − 3, find f(2).
 (b) Evaluate |-5| |-7|.
- (c) Write $\frac{2}{b} + \frac{1}{3}$ as a single fraction.
- (d) Express $\frac{3}{2-\sqrt{2}}$ as a simplified fraction with a rational denominator.
- (e) What is the exact value of $\sqrt{27} + \sqrt{12}$?
- (f) (i) Factorise $x^2 x 6$.
 - (ii) Hence solve $x^2 x 6 = 0$.
- (g) Consider the points A(6,9) and B(2,1).
 - (i) Find the coordinates of the midpoint of the interval AB.
 - (ii) Find the gradient of AB.
 - (iii) Find the equation of the line which passes through points A and B.

Marks

1

1

1

 $\mathbf{2}$

1

1

1

1

1


 $\mathbf{2}$

SGS Half-Yearly 2016 Form V Mathematics 2 Unit Page 6

QUESTION TEN (12 marks) Use a separate writing booklet.	Marks
(a) Show that the lines $y = 3x - 4$ and $x + 3y = 12$ are perpendicular.	2
(b) Solve $2x - 4y = -12$ and $3x + 2y - 2 = 0$ simultaneously.	2
(c) Consider the parabola with equation $y = x^2 - 6x + 7$.	
(i) Show that the parabola has x-intercepts $x = 3 + \sqrt{2}$ and $x = 3 - \sqrt{2}$.	1
(ii) Write down the y -intercept.	1
(iii) Write down the equation of the axis of symmetry.	1
(iv) Find the coordinates of the vertex.	1

(v) Sketch the graph of $y = x^2 - 6x + 7$, clearly marking all of the above features.

(d) Given that $\cos \theta = \frac{3}{7}$ and $\tan \theta > 0$, find $\sin \theta$. Leave your answer in exact form.






SGS Half-Yearly 2016 Form V Mathematics 2 Unit Page 7

QUESTION ELEVEN (12 marks) Use a separate writing booklet.

- (a) (i) Solve |x 1| = 5.
 - (ii) Solve |3x + 2| > 1.
- (b) Solve $\cos x = 0.6$, for $0^{\circ} \le x \le 360^{\circ}$. Give your answer to the nearest degree.
- (c) Find the perpendicular distance from the point P(2,3) to the line x + 2y + 3 = 0.
- (d)



The graph of y = g(x) is shown above. It has horizontal asymptote y = 1. State the coordinates of the point A and the equation of the asymptote after each of the following transformations.

(i)
$$y = g(x) + 1$$

(ii)
$$y = -g(x)$$

	2	
l	2	

2 2 2

 $\mathbf{2}$

Marks


Examination continues overleaf ...

QUESTION TWELVE (12 marks) Use a separate writing booklet.

(a) Sketch each function below on separate axes, including labelled asymptotes and intercepts with axes if they exist.

(i) $y = \sqrt{x+4}$	1
(ii) $y = x+5 $	2
(iii) $y = \frac{1}{x-1} + 3$	3

(b)



The diagram above shows the path of a hiker. The hiker starts from point A and walks on a bearing of 035° T for 2 km to point B. She leaves point B on a bearing of 320° T and walks for 3.5 km until she reaches point C.

- (i) Show that $\angle ABC$ is 105°. Justify your answer with geometrical reasoning.
- (ii) Find the distance AC. Give your answer correct to the nearest metre.
- (iii) Find the bearing of C from A. Give your answer correct to the nearest degree.

2	
2	
2	

Marks

SGS Half-Yearly 2016 Form V Mathematics 2 Unit Page 9

QUESTION THIRTEEN (12 marks) Use a separate writing booklet.

- (a) Consider the function $f(x) = 2^x + 1$.
 - (i) Sketch the graph of y = f(x).
 - (ii) State the range of the function.
- (b) Find the angle of inclination of the line y = -4x + 2. Give your answer correct to the nearest degree.

(c) Solve
$$\cos 2\alpha = \frac{\sqrt{3}}{2}$$
, for $0^{\circ} \le \alpha \le 360^{\circ}$.

- (d) Consider the points A(3,9) and B(1,1) and the circle with diameter AB.
 - (i) Show that the equation of the circle is $(x-2)^2 + (y-5)^2 = 17$.
 - (ii) The line y = x + 6 cuts the circle at point A and again at a second point C. Find the coordinates of the point C.

2	
1	

 $\mathbf{2}$

Marks

3	


2	
2	

Examination continues overleaf

QUESTION FOURTEEN (12 marks) Use a separate writing booklet.

- (a) (i) Express $A^3 + B^3$ as a product of two factors.
 - (ii) Hence prove the identity $\frac{\sin^3 x + \cos^3 x}{\sin x + \cos x} = 1 \sin x \cos x.$

(b)



Consider the functions f(x) = |x+3| - 2 and $g(x) = -(x+2)^2 + 5$ in the diagram above.

- (i) Show that y = f(x) and y = g(x) intersect at (0, 1).
- (ii) Find the exact values of the coordinates of the second point of intersection of y = f(x) and y = g(x).
- (iii) Hence solve $|x+3| 2 \le -(x+2)^2 + 5$.
- (c) Two points A and B lie on a line l. The midpoint of AB is M(6,7). The line 2x + 5y 15 = 0 passes through A and the line 3x 2y + 5 = 0 passes through B. Let A have coordinates A(h, k).
 - (i) Show that the coordinates of B are B(12 h, 14 k).
 - (ii) Find the equation of l.

End of Section II

END OF EXAMINATION

	1	
1	3	

T

3

Marks

1

 $\mathbf{2}$

NAME:

SYDNEY GRAMMAR SCHOOL



2016 Half-Yearly Examination FORM V MATHEMATICS 2 UNIT Tuesday 17th May 2016

- Record your multiple choice answers by filling in the circle corresponding to your choice for each question.
- Fill in the circle completely.
- Each question has only one correct answer.

Question One								
A 🔿	В ()	С ()	D ()					
Question Two								
A ()	В ()	С ()	D ()					
Question Three								
A 🔿	В ()	С ()	D ()					
Question Four								
A 🔿	В ()	С ()	D ()					
Question Five								
A 🔿	В ()	С ()	D ()					
Question Six								
A 🔿	В ()	С ()	D ()					
Question Seven								
A 🔿	В ()	С ()	D ()					
Question Eight								
А ()	В ()	$C \bigcirc$	D ()					

(,		2016	Mathema	tics Soluti	ons.
	1), D				
	2) C				
	3,)C				
	4)D				
	5)D				
	6)B				
	7)0				
	8)A				
	0/1				

 Q^{q} a) $f(2) = 4 \times 2 - 3$ =5 6) 1-51-1-71 = -2 c) 2+1 $= \frac{6}{3b} + \frac{b}{3b}$ $= \frac{6+b}{3b}$ d) $\frac{3}{2-52}$ $=\frac{3(2+\sqrt{2})}{(2-\sqrt{2})(2+\sqrt{2})}$ = $\frac{6+3\sqrt{2}}{2}$ V (or factorised) e) 3/3+2/3=5/3 f(x) = (x+a)(x-3)ii) x = -2 or x = 3g) i) $M = \begin{pmatrix} 6+2 & 9+1 \\ -2 & -2 \end{pmatrix}$ $=\left(\frac{8}{2},\frac{10}{2}\right)$ $-(2'\bar{2})$ = (4,5) V $m = \frac{9-1}{6-2}$ m=2 V iii) y-1 = 2(x-2) y = 2x - 4 + 1 y = 2x - 3

Q10. a) y = 3pc - 4 $m_1 = 3$ x+3y = 12 3y = -x+12 y = -1x+4 3either $\sqrt{-2} m_2 = -\frac{1}{3}$ $3x - \frac{1}{2} = -1$ or m, xm2 = -1 (if explicitly stated m, and m2 above) 5) (1) 2x - 4y = -122 3x + 2y = 26x + 4y = 4 $\begin{array}{rcl}
(2)+1 & 6x+4y = 4 \\
+ (2x-4y = -12) \\
8z &= -8 \\
\hline
x &= -1
\end{array}$ V for correctly formines an equation with one variable; 3(2y-6)+2y-2=0 Sub: -2 - 4y = -12y = 5/2

c) $y = x^2 - 6x + 7$ i) a=1 b=-6 c=7 $\int_{x=3\pm\sqrt{2}}^{2} (x-3) = 2$ $\frac{6 + \sqrt{(-6)^2 - 4 \times 1 \times 7}}{2 \times 1}$ χ= $x = 6 \pm \sqrt{8}$ x= 3± 12 (0,7)ā) ar 7 iii) x = -b= 6 $y = 3^2 - 6 \times 3 + 7$ iv) : (3,-2) V vfor missing -intercepts - poor shape - arxis labels 3-52 ¥ 3+√Z (3,-2) vertex label x=250 V (or 1st guad but d 76 $Sin O = \frac{250}{7} \sqrt{Accept \frac{540}{7}}$ 0

QII. a) i) x - 1 = 5 or x - 1 = -5nc = − 4 ∕ x=b / (or other valid method I mark por working) ;i) 30C+271 or 320+22-1 3x 4-3 3x>-1 >c>-1/3 x 2 - 1 xx-1 V ar x>-13V ÷. 6) cos x = 0.6. n = 53.13 x = 53 (to remest degree) 5 A 53 53 53 and x= 306.86° x= 307° (to nearest degree). c) P(2,3) = +2y+3=0 $x_1 = 2$ $y_1 = 3$ a = 1 b = 2 c = 3 $d = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$ = |l(2) + 2(3) + 3| $\sqrt{l^2 + 2^2}$ = |||| $\sqrt{5}$ = 115

10 CT - - d) i) (1, 4) y = 2ii) (1, -3) y = -1

a) i) >2 shape v intercepts ii20= 1 shape ~ iii) asymptotes / = 3 2 $35^{\circ} + 5c = 180^{\circ}$ (Cointerior angles between Illines $x = 145^{\circ}$ $360^{\circ} - 320^{\circ} = 40^{\circ}$ $145^{\circ} - 40^{\circ} = 105^{\circ}$ (Adjacent Angles) (Cointerior argles between Illines) b) i) ii) a2= 102+ c2- 2bclosA a2= 35002 + 20002 - 2x 3500x 2000 x Cos 105 a = 4458m (to rearest m) Suft = sin 105 ìï) 4458 3500 = 49.3197 A = 360-(49.31996-35) = 346 degrees to nearest degree.

Q13) a)i) y= 2+1 asymptote, labelled; V Shape + intercept; V - y= 1 >>c ii) / y> 6 m = -4 \propto tan B = 4 β=75.96375... 104.036 = 104° (to rearest degree) $\cos 2\alpha = \sqrt{3}$ for 0° 4 × 4 3 60° c)let u=2R for O'Eu E 720° $\cos u = \sqrt{3}$ related angle: U = 30° 33) $\mu = 30^{\circ}, 330^{\circ}, 390^{\circ}, 690^{\circ}$ $\chi = 15^{\circ}, 115^{\circ}, 195^{\circ}, 345^{\circ}$

(d) (i) A(3,9) B(1,1). $M = \left(\frac{3+1}{2}, \frac{9+1}{2}\right) \checkmark$ M= (2,5) certre of circle. $MB = \sqrt{(2-i)^{2} + (5-i)^{2}}$ = $\sqrt{17}$: Radius is JIT ii) y= >c+ 6 ~ $(x-2)^{2} + ((x+6) - 5)^{2} = 17$ $x^2 - 4x + 4 + x^2 + 2x + 1 = 17$ $2\pi c^2 - 2\pi c + 5 = 17$ $2x^2 - 2x - 12 = 0$ $x^2 - x - 6 = 0$ (x+2)(x-3) = 0y = -2, y = 4ar $\chi = 3$... (-2,4) is the other point of intersection

Q14) a) i) $A^{3} + B^{3} = (A + B)(A^{2} - AB + B^{2})$ ii) LHS = (Sinx + cosx) (Sin2x - Sinx cosx + cos2x) V Sinc + COSX = Sizic + cosix - Sixcosi = 1- six cosx = Rots as required $g(x) = -(x+2)^{2}+5$ $g(0) = -(0+2)^{2}+5$ $g(0) = -(0+2)^{2}+5$ (b) i) f(x) = |x+3| - 2f(0) = 10+31-2 =1 (both) ii) when x 2-3 f(x) = - (x+3)-2 $-(x+3) - 2 = -(x+2)^2 + 5$ ->c - 3 - 5 = - (x2 + 4x + 4) + 5 -x - 5 = -x2 - 4x +1 $x^2 + 3x - 6 = 0$ $x = -3 - \sqrt{9 - 4 \times 1 \times - 6}$ 2+1 $\frac{2}{2} = -3 \pm \sqrt{33}$ as x2-3, z=-3-53 $f\left(\frac{-3-\sqrt{33}}{2}\right) = \frac{-3-\sqrt{33}+3}{2} - 2$ $= \frac{3-\sqrt{33}}{2} - 2$ = 133-7 -3-53, 53-7 -3-133 = x 40 ini

 $M = \left(\frac{x_{1} + x_{2}}{2}, \frac{y_{1} + y_{2}}{2}\right) = \left(6, 7\right).$ 2) i) A(h,k). let B heme coordinates B(x, y,). $b = \alpha_1 + h$ $12 = x_1 + h$ (or equivalent) $2c_1 = 12 - h$ $7 = \frac{y_1 + k}{2}$ $14 = y_1 + k$ $y_1 = 14 - k$. : B(12-h, 14-k) " A (h, k) lies on lie 2x + 5y - 15 = 0. : 1 2h+5K-15=0 B(12-h, 14-k) lies on live 3x-2y+5= · (2) 3(12-n) -2(14-k)+5=0 36-3h - 28 + 2k + 5= 0 -3h + 2k + 18 = 0. 0×3+0×2: 6h + 15k - 45 = 0-6h + 4k + 2b = 019k - 19 = 0k = 1h = 5Sub into O . A(5,1) M(6,7) $y - 1 = \frac{7 - 1}{6 - 5} (x - 5)$ y=6x-29