



**GIRRAWEEN HIGH SCHOOL**

**YEAR 11 - TASK 1**

**2006**

**MATHEMATICS**

*Time allowed – 90 minutes*

**DIRECTIONS TO CANDIDATES**

- Attempt ALL questions.
- All necessary working should be shown in every question. Marks may be deducted for careless or badly arranged work.
- Board-approved calculators may be used.
- Start each question on a *new* sheet of paper.

**Question 1 (16 marks)**

(a) Find  $\frac{\sqrt{2.8+7.6}}{4.7-2.9}$  correct to 3 decimal places. 2

(b) Evaluate  $\sqrt{\frac{4.91 \times 10^5}{7.46 \times 10^9}}$  correct to 3 significant figures. 1

(c) Write 6 340 000 000 in scientific notation. 1

(d) Evaluate  $\frac{|10-12|}{|5| \times |-7|}$  2

(e) If  $S = \frac{a}{1-r}$  find the value of a when  $S = 30$  and  $r = \frac{3}{4}$  2

(f) After GST of 10% the new price of an item is \$16.50  
What was the price of the item before GST ? 2

(g) Express the following as a fraction in simplest form. 6

(i) 0.14

(ii) 0. $\dot{3}2\dot{4}$

**Question 2 (22 marks)**

(a) Simplify.

(i)  $4\sqrt{5} \times 3\sqrt{7}$

1

(ii)  $\sqrt{8} + \sqrt{32} - \sqrt{2}$

2

(iii)  $\frac{5\sqrt{3} \times 6\sqrt{15}}{2\sqrt{5}}$

3

(b) Expand and simplify :

(i)  $\sqrt{2}(\sqrt{6} - 1)$

2

(ii)  $(5\sqrt{3} - 2)^2$

3

(c) Express with a rational denominator :

(i)  $\frac{4}{3\sqrt{2}}$

2

(ii)  $\frac{2}{2 - \sqrt{3}}$

3

(iii)  $\frac{2\sqrt{3}}{\sqrt{2} + \sqrt{3}}$

3

(d) Show that  $\frac{1}{5 - \sqrt{7}} + \frac{1}{5 + \sqrt{7}}$  is a rational number.

3

**Question 3 (21 marks)**

(a) Expand and simplify :

(i)  $4(a - 2b) - 4(b - 4a)$

2

(ii)  $(2x^2 - 1)(2x^2 + 1)$

2

(iii)  $(x - y)(x^2 + xy + y^2)$

2

(iv)  $(x + 1)(x - 3)^2$

3

(b) Factorise :

(i)  $a^2 + 6a + 9$

2

(ii)  $3x^3 - 27$

2

(iii)  $2x^2 - 7x - 15$

3

(iv)  $ax + by - bx - by$

2

(v)  $8x^3 - 125$

3

**Question 4 (17 marks)**

(a) Simplify :

$$(i) \frac{x}{3} - \frac{x}{7}$$

2

$$(ii) \frac{1}{x+1} - \frac{1}{x-1}$$

3

$$(iii) \frac{3}{x+2} + \frac{x}{x-3}$$

3

(b) Simplify :

$$(i) \frac{1-t^2}{t-1}$$

3

$$(ii) \frac{m^2 + 2m - 8}{m^2 - 4}$$

3

$$(iii) \frac{x^3 + 8}{x^2 - 2x + 4}$$

3

**Question 5 (20 marks)**

(a) Solve the following equations :

(i)  $4(x + 1) - 3(x - 1) = 6$  3

(ii)  $(3x - 1)(x - 2) = 0$  2

(iii)  $\frac{2-x}{2x} = -5$  2

(iv)  $4x^2 + 7x - 2 = 0$  3

(b) Use the method of completing the square to solve the equation  $x^2 + 6x + 7 = 0$ . 3

[ Give answer in surd form]

(c) Use the quadratic formula to solve the equation  $3x^2 - 3x - 1 = 0$ . 3

[ Give answer in exact form]

(d) A rectangle has a perimeter of 30 cm and a length of  $x$  cm. 4(i) Show that its area,  $A$ , is given by  $A = 15x - x^2$ .(ii) Find the length of the rectangle when the area is  $56 \text{ cm}^2$

**Question 6** <sup>17</sup> (19 marks)

(a) Solve :

(i)  $-3x \geq 6$

1

(ii)  $\frac{3x}{2} < x + 3$

2

(b) Solve  $|2x - 3| < 5$  and graph the solution on the number line.

4

(c) Solve the following simultaneous equations :

(i)  $x - 5y = -11$   
 $x + 2y = -4$

3

(ii)  $3x - 7y = 1$   
 $4x - 5y = -3$

3

(iii)  $y = x^2 - 3x + 3$   
 $2x - y - 3 = 0$

4

End of Assessment Task 1



# Mathematics Yr 11 Task 1, 2006 (Solutions)

## QUESTION 1

$$(a) \frac{\sqrt{10.4}}{1.8} = 1.792 \# \quad (2)$$

$$(b) 8.11 \times 10^{-3} \# \quad (1)$$

$$(c) 6.34 \times 10^9 \# \quad (1)$$

$$(d) \frac{|10-12|}{(5)(1-7)} = \frac{2}{5 \times 7} = \frac{2}{35} \# \quad (2)$$

$$(e) 30 = \frac{a}{1-\frac{1}{4}}$$

$$30 = \frac{a}{\frac{3}{4}} \quad (2)$$

$$a = 7\frac{1}{2} \#$$

$$(f) 110\% = \$16.50$$

$$\therefore 1\% = \frac{\$16.50}{110}$$

$$100\% = \frac{\$16.50}{110} \times 100 = \$15 \# \quad (2)$$

$$(g) (i) \text{ Let } x = 0.14444\ldots \quad (1)$$

$$\therefore 110x = 1.44444\ldots \quad (2)$$

$$(2) - (1) \Rightarrow 9x = 1.3$$

$$\therefore x = \frac{13}{90} \# \quad (3)$$

$$(ii) \text{ Let } x = 0.324324324\ldots \quad (1)$$

$$\therefore 1000x = 324.324324\ldots \quad (2)$$

$$(2) - (1) \Rightarrow 999x = 324$$

$$x = \frac{324}{999}$$

$$x = \frac{12}{37} \# \quad (3)$$

QUESTION 2

$$(a) (i) 4\sqrt{5} \times 3\sqrt{7} = 12\sqrt{35} \quad (1)$$

$$\begin{aligned} (ii) \quad & \sqrt{8} + \sqrt{32} - \sqrt{2} \\ &= 2\sqrt{2} + 4\sqrt{2} - \sqrt{2} \\ &= 5\sqrt{2}. \end{aligned} \quad (2)$$

$$\begin{aligned} (iii) \quad & \frac{5\sqrt{3} \times 6\sqrt{15}}{2\sqrt{5}} \\ &= \frac{5\sqrt{3} \times 6\sqrt{5} \cdot \sqrt{3}}{2\sqrt{5}} = 45 \end{aligned} \quad (3)$$

$$\begin{aligned} (b) \quad (i) \quad & \sqrt{2}(\sqrt{8}-1) = \sqrt{12} - \sqrt{2} \\ &= 2\sqrt{3} - \sqrt{2}. \end{aligned} \quad (2)$$

$$\begin{aligned} (ii) \quad & (5\sqrt{3}-2)^2 \\ &= (5\sqrt{3})^2 + 2 \times 5\sqrt{3} \times -2 + (-2)^2 \\ &= 75 - 20\sqrt{3} + 4 \\ &= 79 - 20\sqrt{3} \end{aligned} \quad (3)$$

$$(c) \quad (i) \quad \frac{4}{3\sqrt{2}} = \frac{4}{3\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{4\sqrt{2}}{6} = \frac{2\sqrt{2}}{3} \quad (2)$$

$$\begin{aligned} (ii) \quad & \frac{2}{2-\sqrt{3}} = \frac{2}{2-\sqrt{3}} \times \frac{2+\sqrt{3}}{2+\sqrt{3}} \\ &= \frac{2(2+\sqrt{3})}{4-3} \\ &= 4+2\sqrt{3} \# \end{aligned} \quad (3)$$

$$\begin{aligned} (iii) \quad & \frac{2\sqrt{3}}{\sqrt{2}+\sqrt{3}} = \frac{2\sqrt{3}}{\sqrt{2}+\sqrt{3}} \times \frac{\sqrt{2}-\sqrt{3}}{\sqrt{2}-\sqrt{3}} \\ &= \frac{2\sqrt{6}-6}{2-3} \\ &= 6-2\sqrt{6} \# \end{aligned} \quad (3)$$

$$(d) \frac{1}{5-\sqrt{7}} + \frac{1}{5+\sqrt{7}}$$

$$= \frac{5+\sqrt{7} + 5-\sqrt{7}}{(5-\sqrt{7})(5+\sqrt{7})}$$

$$= \frac{10}{25-7} = \frac{10}{18} = \frac{5}{9} \text{ which is rational. } \# \quad (3)$$

### QUESTION 3

$$(a) (i) 4(a-2b) - 4(b-4a)$$

$$= 4a - 8b - 4b + 16a$$

$$= 20a - 12b. \# \quad (2)$$

$$(ii) (2x^2-1)(2x^2+1)$$

$$= 4x^4 - 1 \# \quad (2)$$

$$(iii) (x-y)(x^2+xy+y^2)$$

$$= x^3 + x^2y + xy^2 - x^2y - xy^2 - y^3$$

$$= x^3 - y^3 \# \quad (2)$$

$$(iv) (x+1)(x-3)^2$$

$$= (x+1)(x^2 - 6x + 9)$$

$$= x^3 - 6x^2 + 9x + x^2 - 6x + 9$$

$$= x^3 - 5x^2 + 3x + 9 \# \quad (3)$$

$$(b) (i) a^2 + 6a + 9 = (a+3)^2 \quad (2)$$

$$(ii) 3x^3 - 27$$

$$= 3(x^3 - 9) \# \quad (2)$$

$$(iii) 2x^2 - 7x - 15 \quad ab = -30 \quad a+b = -7 \quad \left\{ -10, 3 \right.$$

$$= 2x^2 - 10x + 3x - 15$$

$$= 2x(x-5) + 3(x-5) = (2x+3)(x-5) \# \quad (3)$$

$$\begin{aligned}
 (iv) \quad & ax + bx - bx - by \\
 &= ax - bx \\
 &= x(a - b) \quad \# \quad (2)
 \end{aligned}$$

$$\begin{aligned}
 (v) \quad & 8x^3 - 125 \\
 &= (2x)^3 - 5^3 \\
 &= (2x - 5)(4x^2 + 10x + 25) \quad \# \quad (3)
 \end{aligned}$$

QUESTION 4 (a) (i)  $\frac{x}{3} - \frac{x}{7} = \frac{7x - 3x}{21} = \frac{4x}{21} \# \quad (2)$

$$\begin{aligned}
 (ii) \quad & \frac{1}{x+1} - \frac{1}{x-1} \\
 &= \frac{x-1 - (x+1)}{(x+1)(x-1)} = \frac{-2}{x^2 - 1} \# \quad (3)
 \end{aligned}$$

$$\begin{aligned}
 (iii) \quad & \frac{3}{x+2} + \frac{x}{x-3} \\
 &= \frac{3(x-3) + x(x+2)}{(x+2)(x-3)} = \frac{3x-9+x^2+2x}{(x+2)(x-3)} \\
 &= \frac{x^2+5x-9}{(x+2)(x-3)} \# \quad (3)
 \end{aligned}$$

(b) (i)  $\frac{1-t^2}{t-1} = \frac{(1-t)(1+t)}{-(1-t)} = -1-t \# \quad (3)$

$$\begin{aligned}
 (ii) \quad & \frac{m^2+2m-8}{m^2-4} = \frac{(m+4)(m-2)}{(m+2)(m-2)} = \frac{m+4}{m+2} \# \quad (3)
 \end{aligned}$$

$$\begin{aligned}
 (iii) \quad & \frac{x^3+8}{x^2-2x+4} = \frac{(x+2)(x^2-2x+4)}{(x^2-2x+4)} = x+2 \# \quad (3)
 \end{aligned}$$

QUESTION 5

(a) (i)  $4(x+1) - 3(x-1) = 6$

$$4x + 4 - 3x + 3 = 6$$

$$x = 6 - 7$$

$$\therefore x = -1 \quad \text{#}$$

(3)

(ii)  $(3x-1)(x-2) = 0$

$$\therefore 3x-1 = 0 \text{ or } x-2 = 0$$

$$\therefore x = \frac{1}{3} \text{ or } x = 2.$$

(2)

(iii)  $\frac{2-x}{2x} = -5$

$$2-x = -10x$$

$$10x-x = -2$$

$$9x = -2$$

$$x = -\frac{2}{9} \quad \text{#}$$

(2)

(iv)  $4x^2 + 7x - 2 = 0$

$$(4x-1)(x+2) = 0$$

$$\therefore 4x-1 = 0 \text{ or } x+2 = 0$$

$$x = \frac{1}{4} \text{ or } x = -2 \quad \text{#}$$

(3)

(b)  $x^2 + 6x = -7$

$$x^2 + 6x + 3^2 = -7 + 3^2$$

$$(x+3)^2 = 2$$

$$\therefore x+3 = \pm\sqrt{2}$$

$$\therefore x = -3 \pm \sqrt{2} \quad \text{#}$$

(3)

(c)  $x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4 \times 3 \times -1}}{2 \times 3}$

$$= \frac{3 \pm \sqrt{9+12}}{6}$$

$$= \frac{3 \pm \sqrt{21}}{6} \quad \text{#}$$

(3)

(d)

$$\text{Area } A = xy$$

$$A = x(15-x) \text{ using } (1)$$

$$\therefore A = 15x - x^2$$

$$(ii) 15x - x^2 = 56$$

$$\therefore x^2 - 15x + 56 = 0$$

$$(x-8)(x-7) = 0$$

$$\therefore x=8 \text{ or } x=7$$

$$\therefore \text{length is } 8 \text{ cm } (x>y).$$
4

### QUESTION 6

(a) (i)  $-3x > 6, \therefore \frac{-3x}{-3} < \frac{6}{-3}, \therefore x < -2$  # (1)

$$(ii) \frac{3x}{2} < x+3$$

$$3x < 2x+6$$

$$3x-2x < 6$$

$$x < 6$$
 #
2

(b)  $|2x-3| < 5$

$$\therefore 2x-3 < 5 \text{ or } -2x+3 < 5$$

$$2x < 8 \quad -2x < 2$$

$$x < 4 \quad x > -1$$
4

(c) (i)  $x-5y = -11 \dots (1)$

$$x+2y = -4 \dots (2)$$

(2) - (1)  $\Rightarrow 7y = 7$

$$\therefore y = 1.$$

From (1),  $x = -11 + 5$

$$\therefore x = -6$$

$$\therefore x = -6, y = 1$$
 #

$$\begin{aligned}
 & 6(c) \quad (ii) \quad 3x - 7y = 1 \quad \dots \quad (1) \\
 & \quad 4x - 5y = -3 \quad \dots \quad (2) \\
 & 4 \times (1) \Rightarrow 12x - 28y = 4 \quad \dots \quad (3) \\
 & 3 \times (2) \Rightarrow 12x - 15y = -9 \quad \dots \quad (4) \\
 & (3) - (4) \Rightarrow -13y = 13 \\
 & \therefore y = -1 \\
 & \text{From (1)} \quad 3x + 7 = 1 \\
 & \quad 3x = -6 \\
 & \quad x = -2 \\
 & \therefore x = -2, y = -1 \quad \# \quad (3)
 \end{aligned}$$

$$\begin{aligned}
 & (iii) \quad y = x^2 - 3x + 3 \quad \dots \quad (1) \\
 & \quad 2x - y - 3 = 0 \quad \dots \quad (2) \\
 & \text{Sub. (1) into (2),} \\
 & \therefore 2x - (x^2 - 3x + 3) - 3 = 0 \\
 & 2x - x^2 + 3x - 3 - 3 = 0 \\
 & -x^2 + 5x - 6 = 0 \\
 & x^2 - 5x + 6 = 0 \\
 & (x-3)(x-2) = 0 \\
 & \therefore x = 3 \text{ or } x = 2
 \end{aligned}$$

$$\begin{aligned}
 & \text{When } x = 3, \quad y = 3^2 - 3 \times 3 + 3 = 3 \\
 & \quad x = 2, \quad y = 2^2 - 3 \times 2 + 3 = 1 \\
 & \therefore x = 3, y = 3 \quad \# \\
 & \quad x = 2, y = 1 \quad \# \quad (4)
 \end{aligned}$$

