

SYDNEY TECHNICAL HIGH SCHOOL
MATHEMATICS EXTENSION 1
YEAR 11 YEARLY EXAMINATION

2002

Time allowed: 90 minutes

Instructions:

- Show all necessary working
- Start each question on a new page
- All questions are of equal value
- Marks will not be awarded for careless or badly arranged work
- Non-programmable calculators may be used
- This paper must be handed in with your answer sheets
- Answers must be written in blue or black pen

Name: _____

Class: _____


Question 1	Question 2	Question 3	Question 4	Question 5	Question 6	TOTAL

Question 1**Marks**

a) If $(x + 1)$ is a factor of $P(x) = x^3 - ax + 3$. Find the value of a .

1

b)

2

Write down the equation of the polynomial function (in factored form)

c) A parabola is symmetrical about the line $y = 2$ it has a focal length 3 units and the equation of the directrix is $x = 1$.

3

i) How many parabolas satisfy these conditions?

ii) If the vertex is $(4, 2)$ find the equation of the parabola

d) Solve $\frac{x+1}{x-1} \leq 0$

2

e) The roots of the quadratic equation $(k + 2)x^2 - 4x + k^2 = 0$ are reciprocals. Find the value/s of k .

3

Question 2**Marks**

- a) A polynomial of degree 7 is divided by the polynomial $Q(x)$, the remainder is $x^2 + x + 2$. What is the least degree of $Q(x)$. 1
- b) For the quadratic equation $x^2 + (k - 3)x + 2 - k = 0$ 3
- i) Find the value of the discriminant in the terms of k
- ii) Explain why the roots of this quadratic equation are real for all values of k
- c) If $a + b = 1$ and $a^2 + b^2 = 2$ 3
- i) Find the value of ab
- ii) Hence find the value of $a^3 + b^3$
- d) i) Write $x^{-\frac{1}{2}}$ with a positive index 4
- ii) Solve $x^{\frac{1}{2}} + 10x^{-\frac{1}{2}} = 7$

Question 3

Marks

a) For the function $y = \sqrt{x^2 - 4}$

3

Write down


- i) the domain
- ii) the range

b) The points $P(12t, 6t^2)$ and $Q(36, 54)$ are points on a parabola

3

- i) Find the cartesian equation of the parabola
- ii) If PQ is a focal chord find the value of t

c)



A cyclist riding along a straight flat road passes by three stop signs A, B and C spaced 200m apart. From these three signs the respective angles of elevation to the top of a mobile phone tower are $45^\circ, 45^\circ$ and 30° . If 'h' is the height of the tower GH.

5

- i) Show that $CG = \sqrt{3} h$.
- ii) If $\angle GBA = \alpha$. Find two different expressions for $\cos \alpha$ in terms of h .
- iii) Hence find the height of the tower.

Question 4**Marks**

- a) i) Simplify
 $(\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$ 3
- ii) The roots of $x^3 - 4x^2 - 8 = 0$ are α , β and γ . Use the result in part (i) to find
The value of $\alpha^2 + \beta^2 + \gamma^2$.
- b) i) Show that $\frac{1 + \cos 2A}{\sin 2A} = \cot A$ 4
- ii) Hence find the exact value of $\cot 15^\circ$
- c) T $(2t, t^2)$ is a point on the parabola $x^2 = 4y$ with focus S. 4
P is the point which divides ST internally in the ratio 1 : 2.
- i) Write down the coordinates of P in terms of t .
- ii) Hence show that as T moves on the parabola $x^2 = 4y$ that
the locus of P is the parabola $9x^2 = 12y - 8$

Question 5

- a) The roots of the equation $x^3 - 6x^2 + 5x + 8 = 0$ are α, β, γ 3
- The roots of the equation $x^3 + ax^2 + bx + 512 = 0$ are $k\alpha, k\beta, k\gamma$
- i) Find the value of k
- ii) Hence find the value of b .
- b) Consider the points A(-2, 3) B(6, 5) the point P(x, y) moves 4
so that the angle APB = 90°
- i) Write down an expression for the gradient of AP
- ii) Show that the locus of P is a circle
- iii) Find its centre and radius.
- c) i) Expand $\tan(A + B)$ 4
- ii) The roots of $x^2 - 2x - 1 = 0$ are $\tan A$ and $\tan B$. If A and B are acute
find the size of $A + B$.


Question 6**Marks**

- a) i) Derive the equation of the tangent to the parabola $x^2 = 4ay$ at the point $P(2at, at^2)$ 7
- ii) The tangent cuts the y axis at R . Find the co-ordinates of R .
- iii) If S is the focus of the parabola. Find the length of PS .
- iv) Prove that the triangle PSR is isosceles
- v) If $\angle PSR = 120^\circ$. Find the numerical value of t .
-
- b) If $P(x) = 4x^3 + 9x - 4$ 4
- i) Find $P(\alpha + 1)$
- ii) If α is a root of $P(x)$ use part (i) to help show that $P(\alpha + 1) > 0$

Question 1

a) $P(-1) = 0 \therefore -1 + a + 3 = 0$
 $a = -2$


b) $P(x) = A(x+2)^2(x-1)$
 sub $(0, -12) \therefore A = 3$
 $P(x) = 3(x+2)^2(x-1)$



ii) $(y-2)^2 = 12(x-4)$

d) $\frac{x+1}{x-1} \leq 0$

$\frac{(x-1)^2(x+1)}{(x-1)^2} \leq 0 \cdot (x-1)^2$

$(x-1)(x+1) \leq 0$ 

$x: -1 \leq x < 1$

e) $\alpha, \frac{1}{\alpha}$ roots

$\therefore \text{prod} = 1$

$\frac{k^2}{k+2} = 1$

$k+2$

$k^2 = k+2$

$k^2 - k - 2 = 0$

$(k-2)(k+1) = 0$

$\therefore k = 2, k = -1$

$\therefore \sqrt{x} = 5 \quad \sqrt{x} = 2$
 $x = 25 \quad x = 4$

Question 2

a) $Q(x)$ deg 3

b) i) $\Delta = (k-3)^2 - 4 \cdot 1 \cdot (2-k)$
 $= k^2 - 6k + 9 - 8 + 4k$
 $\Delta = k^2 - 2k + 1 = (k-1)^2$

ii)

since $\Delta \geq 0$ for all values of k

\therefore roots real

c) i) $(a+b)^2 = a^2 + b^2 + 2ab$
 $(a+b)^2 - (a^2 + b^2) = 2ab$
 $1 - 2 = 2ab$

$\therefore ab = -\frac{1}{2}$

ii)

$a^3 + b^3$

since $(a+b)^3 =$

$a(a^2 + 2ab + b^2) + b(a^2 + 2ab + b^2)$

$a^3 + 3a^2b + 3ab^2 + b^3$

$\therefore a^3 + b^3 = (a+b)^2 - 3a^2b - 3ab^2$

$= (a+b)^2 - 3ab(a+b)$

$= 1 - 3 \times \frac{-1}{2} \cdot 1$

$= 2\frac{1}{2}$

d) i) $x^{-1/2} = \frac{1}{\sqrt{x}}$

ii) $x^{1/2} + 10x^{-1/2} = 7$

$\sqrt{x} + \frac{10}{\sqrt{x}} = 7$ Let $u = \sqrt{x}$

$u + \frac{10}{u} = 7$

$u^2 + 10 = 7u$

$u^2 - 7u + 10 = 0$

$(u-5)(u-2) = 0$

$u = 5$

$u = 2$


Question 3

(2)

a) $y = \sqrt{x^2 - 4}$

$x^2 - 4 \geq 0$

$(x-2)(x+2) \geq 0$




i) D: $x \geq 2, x \leq -2$ (2)

ii) R: $y \geq 0$ (1)

b) i) $P(12t, 6t^2)$

$x = 12t \therefore t = \frac{x}{12}$ $y = 6\left(\frac{x}{12}\right)^2$

$\therefore y = \frac{x^2}{24}$ OR $x^2 = 24y$



find eqn of SQ

$m_{SQ} = \frac{4}{3}$

\therefore eqn SQ $y = \frac{4}{3}x + 6$

sub $P(12t, 6t^2)$

$6t^2 = \frac{4}{3} \cdot 12t + 6$

$6t^2 - 16t - 6 = 0$

$3t^2 - 8t - 3 = 0$

$(3t+1)(t-3) = 0$


$t = 3$ $t = -1/3$ (1)

$t = 3 \Rightarrow Q(36, 54)$

$\therefore t = -1/3 \Rightarrow P(12t, 6t^2)$

$P(-4, \frac{2}{3})$

(1)



A i) $\tan 30^\circ = \frac{h}{CG}$

$\therefore CG = \sqrt{3}h$ (1)

ii)

In $\triangle ABC \Rightarrow \cos \alpha = \frac{h^2 + 200^2 - h^2}{2 \cdot h \cdot 200}$

$\cos \alpha = \frac{200^2}{400h}$ (3)

$\cos \alpha = \frac{100}{h}$ (A)

In $\triangle ABC$

$\Rightarrow \cos(180 - \alpha) = \frac{h^2 + 200^2 - (\sqrt{3}h)^2}{2 \times h \times 200}$

$-\cos \alpha = \frac{20,000 - h^2}{200h}$ (E)

In $\triangle ACC \Rightarrow \cos \alpha = \frac{h^2 + 400^2 - (\sqrt{3}h)^2}{800h}$ (2)

$\cos \alpha = \frac{80,000 - h^2}{400h}$ (C)

any comb of A, B, C.

$-\frac{100}{h} = \frac{20,000 - h^2}{200h}$

$-20000 = 20,000 - h^2$

$h^2 = 40,000$


$h = 200 \text{ m}$ (1)

Question 4


a) i) $(\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$
 $= \alpha(\alpha + \beta + \gamma) + \beta(\alpha + \gamma + \beta) + \gamma(\alpha + \beta + \gamma)$
 $- 2\alpha\beta - 2\beta\gamma - 2\gamma\alpha$
 $= \underline{\underline{\alpha^2 + \beta^2 + \gamma^2}}$

ii) $\alpha^2 + \beta^2 + \gamma^2 = (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha)$ (roots of $ax^2 + bx + c = 0$)
 $a=1 \quad b=-4 \quad c=0 \quad d=-8$
 $= (4)^2 - 2(0)$
 $= \underline{\underline{16}}$

b) i) $LHS = \frac{1 + \cos^2 A - \sin^2 A}{2 \sin A \cdot \cos A}$
 $= \frac{2 \cos^2 A}{2 \sin A \cdot \cos A}$
 $= \frac{\cos A}{\sin A}$
 $= \cot A$
 $= \underline{\underline{RHS}}$



ii) $\cot 15^\circ = \frac{1 + \cos 30^\circ}{\sin 30^\circ}$
 $= (1 + \frac{\sqrt{3}}{2}) \div (\frac{1}{2})$
 $= (\frac{2 + \sqrt{3}}{2}) \times \frac{2}{1}$
 $= \underline{\underline{2 + \sqrt{3}}}$



i) $S(0,1) \xrightarrow[1:2]{\text{internally}} T(2t, t^2)$
 $P(\frac{0+2t}{3}, \frac{2+t^2}{3})$
 $\therefore P(\frac{2t}{3}, \frac{2+t^2}{3})$

ii) $T(2t, t^2)$
 $x = 2t \quad y = t^2$
 $\frac{x}{2} = t \quad \rightarrow \quad y = (\frac{x}{2})^2$
locus of T is $4y = x^2$

$P(\frac{2t}{3}, \frac{2+t^2}{3})$
 $x = \frac{2t}{3} \quad y = \frac{2+t^2}{3}$
 $\frac{3x}{2} = t \quad 3y = 2 + (\frac{3x}{2})^2$
 $4 \times 3y = 8 + 9x^2$
 $12y - 8 = 9x^2$

Question 5

a) i) $\alpha + \beta + \gamma = 6$
 $k\alpha + k\beta + k\gamma = -a$
 $k(\alpha + \beta + \gamma) = -a$
 $\therefore 6k = -a$

$\alpha\beta\gamma = -8$
 $k^3(\alpha\beta\gamma) = -512$
 $-8k^3 = -512$
 $k^3 = 64$
 $k = 4$ $\therefore a = -24$

$$\alpha\beta + \alpha\gamma + \beta\gamma = 5$$


$$k^2\alpha\beta + k^2\alpha\gamma + k^2\beta\gamma = b$$

$$k^2(\alpha\beta + \alpha\gamma + \beta\gamma) = b$$

$$k^2 \cdot 5 = b$$

$$16 \times 5 = b$$

$$\underline{\underline{b = 80}}$$



i) $m_{AP} = \frac{y-3}{x+2}$

ii) $\left(\frac{y-3}{x+2}\right) \cdot \left(\frac{y-5}{x-6}\right) = -1$

$$y^2 - 8y + 15 = -1(x^2 - 4x - 12)$$

$$y^2 - 8y + 15 = -x^2 + 4x + 12$$

$$x^2 + y^2 - 4x - 8y - 3 = 0$$

$$(x^2 - 4x + 4) + (y^2 - 8y + 16) = 17$$

$$(x-2)^2 + (y-4)^2 = \sqrt{17}^2$$

centre (2, 4) radius $\sqrt{17}$

c) i) $\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \cdot \tan B}$

ii) Root $\tan A, \tan B$

Sum $\tan A + \tan B = 2$

$\tan A \cdot \tan B = -1$


$$\tan(A+B) = \frac{2}{1-1}$$

$$\tan(A+B) = 1$$

$$\underline{\underline{A+B = 45^\circ}}$$

Question 6

a) i)



i) $x^2 = 4ay$

$$y = \frac{x^2}{4a}$$

$$\frac{dy}{dx} = \frac{2x}{4a} = \frac{x}{2a}$$

at P: $m_T = \frac{2at}{2a} = t$

∴ eqn tang at P $y - at^2 = t(x - 2a)$

$$\underline{\underline{y = tx - at^2}}$$

ii) $R(0, -at^2)$

iii) $PS = \sqrt{(at^2 - a)^2 + (2at - 0)^2}$

$$= \sqrt{a^2(t^2 - 1)^2 + 4a^2t^2}$$

$$= a\sqrt{t^4 - 2t^2 + 1 + 4t^2}$$

$$= a\sqrt{t^4 + 2t^2 + 1}$$


$$= a\sqrt{(t^2 + 1)^2}$$

$$\underline{\underline{PS = a(t^2 + 1)}}$$

iv) $SR = a + at^2$

$$= a(1 + t^2)$$

$SR = PS \therefore \Delta PSR$ isosceles



$\therefore \tan 60^\circ = t$

(grad of tang)

$$\underline{\underline{\therefore \sqrt{3} = t}}$$

b) $P(x) = 4x^3 + 9x - 4$

i) $P(\alpha+1) = 4(\alpha+1)^3 + 9(\alpha+1) - 4$

ii) α a root \therefore

$P(\alpha) = 0$

$4\alpha^3 + 9\alpha - 4 = 0$

$P(\alpha+1) = 4(\alpha^3 + 3\alpha^2 + 3\alpha + 1) + 9\alpha + 9 - 4$

$= 4\alpha^3 + 12\alpha^2 + 12\alpha + 4 + 9\alpha + 9 - 4$

$= 4\alpha^3 + 12\alpha^2 + 21\alpha + 9$

$= (4\alpha^3 + 9\alpha - 4) + 12\alpha^2 + 12\alpha + 13$

$\therefore P(\alpha+1) = 12\alpha^2 + 12\alpha + 13$

tve def
since $a > 0$ $\Delta < 0$

$\therefore P(\alpha+1) > 0$ for all α