

Name: ..... Maths Class: .....

# SYDNEY TECHNICAL HIGH SCHOOL



Year 11

## Mathematics Extension 1 HSC ASSESSMENT TASK ONE

December 2002

**TIME ALLOWED: 60 minutes**

***Instructions:***


- Write your name and class at the top of this page.
- Start each question on a new page
- At the end of the examination this examination paper must be attached to the front of your answers.
- The marks for each question are indicated on the question sheet
- **ALL** questions should be attempted
- All necessary working must be shown. Marks will be deducted for careless or badly arranged work.
- Marks indicated are a guide only and may be varied if necessary.

| Q 1 | Q 2 | Q 3 | Q 4 | Q 5 | Q 6 | TOTAL |
|-----|-----|-----|-----|-----|-----|-------|
| /7  | /8  | /8  | /9  | /7  | /6  | /45   |

**QUESTION ONE** ( 7 marks )

a) Find the value of  $x$

(2)



b) Evaluate  $6 + 11 + 16 + \dots + 426$

(3)

c) Find the value of  $\sum_{n=4}^7 n^2 + 2$

(2)

**QUESTION TWO** ( 8 marks )


a) If  $S_n = 3n^2 + 2n$ , find,

(4)

- i. The value of the second term
- ii. The  $n$ th term

b)

(4)



ABC is a triangle inscribed in a circle.  
 MAN is the tangent to the circle at A.  
 D is a point on AB and E is a point on AC such that  $DE \parallel MAN$ .

- i. Copy the diagram onto your answer page
- ii. Explain why  $\angle MAB = \angle ACB$ .
- iii. Hence show that BCED is a cyclic quadrilateral.

**QUESTION THREE** ( 8 marks )

a) The fourth term of a geometric sequence is 4 and the seventh term of the same sequence is 32 (4)

- i. Find the value of the first term and the common ratio.
- ii. Find the sum of the first 7 terms


b) If  $1^2 + 3^2 + \dots + (2k-1)^2 = \frac{k(2k-1)(2k+1)}{3}$  (4)

Prove that

$$1^2 + 3^2 + \dots + (2k-1)^2 + (2(k+1)-1)^2 = \frac{(k+1)(2k+1)(2k+3)}{3}$$

**QUESTION FOUR** ( 9 marks )

a)



(2)

ABCD are points on the circumference of a circle with centre O.  
EF is a tangent touching the circle at A.

$$\angle CAE = 50^\circ$$

Find, giving reasons,  $\angle ABC$

b) Prove, by mathematical induction, that  $9^{n+2} - 4^n$  is divisible by 5, for Integers  $n \geq 1$  (4)


c) A geometric series is given as (3)

$$1 + (2x+1) + (2x+1)^2 + \dots$$

- i. For what values of x does the series have a limiting sum ?
- ii. Is it possible for the series to have a limiting sum of -1. Explain.

**QUESTION FIVE** ( 7 marks )

a)



(3)

$TA$  is a tangent to a circle. Line  $ABDC$  intersects the circle at  $B$  and  $C$ .  
Line  $TD$  bisects angle  $BTC$ .

Prove  $AT = AD$

- b) Kermit invests \$2000 at the beginning of each year into an investment account earning 6% p.a. compounded monthly. (4)

Kermit begins his investment on January 1<sup>st</sup> 2002

- i. What is the value of the first investment at the end of 2032?
- ii. If Kermit makes his last investment on January 1<sup>st</sup> 2032, how much is in the account when he withdraws it all on December 31<sup>st</sup> 2032, immediately after the interest for the month has been added?

**QUESTION SIX** ( 6 marks )

(6)

Bert and Ernie have a small business account earning 9% p.a. compounded monthly

Into this account they invest the companies profits of \$5000 at the start of each month. At the end of each month, immediately after the interest has been paid, Bert and Ernie withdraw \$M for the coming month's expenses.

- i. How much is in the account , immediately before the first withdrawal?
- ii. Show that the amount in the account immediately after the second withdrawal is,

$$A_2 = 5000( 1.0075^2 + 1.0075 ) - M( 1.0075 + 1 )$$

- iii. Bert and Ernie hope to have saved \$100 000 by the end of three years, (immediately after the withdrawal for the coming months expenses)  
How much can they afford to withdraw for expenses each month?

### Question One

a)  $10^2 = x(x+8)$  ①

$$x^2 + 8x - 100 = 0$$

$$x = \frac{-8 \pm \sqrt{64 - 4 \times 1 \times -100}}{2}$$

$$= \frac{-8 \pm \sqrt{464}}{2}$$

$$= \frac{-8 \pm \sqrt{29} \cdot 4}{2}$$

$$= -4 + 2\sqrt{29}, -4 - 2\sqrt{29}$$

● as  $x > 0$

$$x = -4 + 2\sqrt{29}$$
 ①

b)  $6 + 11 + 16 + \dots + 426$

AP  $a=6$   $d=5$   $T_n=426$  ①

$$426 = 6 + (n-1)5$$

$$420 = 5n - 5$$

$$425 = 5n$$

$$n = 85$$
 ①

$$\therefore S_n = \frac{n}{2}(a+l)$$

$$= \frac{85}{2}(6 + 426)$$

$$= 18360$$
 ①

c)  $\sum_{n=4}^7 n^2 + 2$

$$= (4^2+2) + (5^2+2) + (6^2+2) + (7^2+2)$$
 ①

$$= 134$$
 ①

### Question Two

a)  $S_n = 3n^2 + 2n$

i.  $T_2 = S_2 - S_1$  ①

$$= (3 \times 4 + 4) - (3 \times 1 + 2)$$

$$= 11$$
 ①


ii.  $5, 11, 17, \dots$

AP  $a=5$   $d=6$

$$\therefore T_n = a + (n-1)d$$
 ①

$$= 5 + (n-1)6$$

$$= 6n - 1$$
 ①



ii.  $\angle MAB = \angle ACB$  ①

The angle between a tangent and a chord is equal to the angle in the alternate segment.

iii.  $\angle MAB = \angle ADE$  (alt L's,  $MN \parallel DE$ ) ①

$$\angle BDE = 180^\circ - \angle ADE$$
 (L's on a st. line add to  $180^\circ$ ) ①

However,  $\angle BDE = 180^\circ - \angle ACB$

(as  $\angle MAB = \angle ACB = \angle ADE$ )

$$\therefore \angle BDE + \angle ACB = 180^\circ$$

and BCED is a cyclic quad as opp. angles are supplementary

### Question three:

a) i.  $T_4 = ar^3 = 4$  — ①

$T_7 = ar^6 = 32$  — ②

② ÷ ①

$r^3 = 8$

$\therefore r = 2$  ✓①

$a = \frac{1}{2}$  ✓①

ii.  $n=7$   $S_n = \frac{a(r^n - 1)}{r - 1}$

$= \frac{\frac{1}{2}(2^7 - 1)}{1}$  ① ✓

$= 63.5$  ① ✓

b)

LHS =  $1^2 + 3^2 + \dots + (2k-1)^2 + (2k+1)^2$   
 $= \frac{k(2k-1)(2k+1)}{3} + (2k+1)^2$  ①

$= \frac{k(2k-1)(2k+1)}{3} + \frac{3}{3}(2k+1)^2$

$= \frac{(2k+1)}{3} [k(2k-1) + 3(2k+1)]$  ①

$= \frac{(2k+1)}{3} [2k^2 + 5k + 3]$


$= \frac{(2k+1)(k+1)(2k+3)}{3}$  ①

$= \frac{(k+1)(2k+1)(2k+3)}{3}$

$= \text{RHS}$

①  
 ↓  
 setting  
 out of  
 LHS =  
 = RHS.

### Question four:



$\angle CAE = 50^\circ$  (given)

$\angle CAE = \angle ADC$  ( $\angle$  between a tangent & a chord =  $\angle$  in the alt. segment) ①  
 $\therefore \angle ADC = 50^\circ$

$\angle ADC + \angle ABC = 180^\circ$

(opp  $\angle$ 's in a cyclic quad ADCB) ①

$\therefore \angle ABC = 130^\circ$

b)  $9^{n+2} - 4^n$  is  $\div$  by 5

Test  $n=1$

$9^3 - 4^1 = 725$

$= 5 \times 145$  which is  $\div$  by 5

$\therefore$  true for  $n=1$  ①

Assume true for  $n=k$

i.e.  $9^{k+2} - 4^k = 5M$  where  $M$  is an integer ①

Prove true for  $n=k+1$

$9^{k+1+2} - 4^{k+1}$

$= 9 \cdot 9^{k+2} - 4 \cdot 4^k$

$= 9(5M + 4^k) - 4 \cdot 4^k$  ①

$= 45M + 9 \cdot 4^k - 4 \cdot 4^k$

$= 45M + 5 \cdot 4^k$

$= 5(9M + 4^k)$  which is  $\div$  by 5

as  $9M + 4^k$  is an integer ①

$\therefore$  true for  $n=k+1$

If true  $n=k$  also true for  $n=k+1$

As true  $n=1$  also true for  $n=1+1=2, 3, 4, \dots$

Hence by M.I. true all true integer  $n$

Q4 c)

$$1 + (2x+1) + (2x+1)^2 + \dots$$

i.  $-1 < r < 1$

$$-1 < 2x+1 < 1$$

$$-2 < 2x < 0$$

$$-1 < x < 0 \quad \textcircled{1}$$

ii.  $S_{\infty} = \frac{a}{1-r}$

$$-1 = \frac{1}{1-(2x+1)}$$

$$-1 = \frac{1}{-2x}$$


$$2x = 1$$

$$x = \frac{1}{2} \textcircled{1} \text{ but as } -1 < x < 0$$

$$x \neq \frac{1}{2}$$

$$\therefore S_{\infty} \neq -1 \quad \textcircled{1}$$

Question five



Prove  $AT = AD$

let  $\angle CTD = x$

$$\therefore \angle DTB = x \text{ (given)}$$

let  $\angle BTA = y$

$$\therefore \angle TCA = y \text{ (alt. segment theorem)}$$

$$\angle TDA = \angle DTC + \angle TCD$$

(exterior  $\angle$  of  $\Delta TDC$ )

$$\therefore \angle TDA = x + y$$

$$\angle DTA = \angle DTB + \angle BTA$$

$$= x + y$$

$$\therefore \angle TDA = \angle DTA \text{ (} x + y \text{)}$$

$$\therefore AD = AT \text{ (sides opp. equal angles)}$$

b)  $6\% \text{ p.a} = \frac{1}{2}\% \text{ p. month}$

Annual Inv  $\rightarrow$  Comp monthly

$$1. \quad 2000(1.005)^{\frac{31}{30 \times 12}} = \$12045.15 \quad \textcircled{1}$$

$$\quad \quad \quad \$12788.07$$

$$11. \quad 2000(1.005)^{30 \times 12}$$

$$+ 2000(1.005)^{29 \times 12} \quad \textcircled{1}$$

$$+ 2000(1.005)^{28 \times 12}$$

$$\quad \quad \quad \vdots$$

$$2000(1.005)^{1 \times 12}$$

$$\therefore \text{Total} = 2000 \left[ 1.005^{12} + 1.005^{24} + \dots + 1.005^{31} \right]$$

$\overbrace{\hspace{2cm}}^{\text{GP}} \quad a = 1.005^{12}$

$$r = 1.005^{12}$$

$$n = 31$$

$$= 2000 \times \frac{1.005^{12} \left[ (1.005^{12})^{31} - 1 \right]}{1.005^{12} - 1} \quad \textcircled{1}$$

$$= \$172910.0466\dots$$

$$= \$172910 \text{ (nearest \$)} \quad \textcircled{1}$$

$$\$185698$$

### Question six

$$r = 1.0075$$

$$\begin{aligned} \text{i. } & 5000(1.0075) \quad \textcircled{1} \\ & = 5037.50 \end{aligned}$$

$$\text{ii. } A_1 = 5000(1.0075) - m$$

$$A_2 = [A_1 + 5000](1.0075) - m \quad \textcircled{1}$$

$$\bullet = 5000(1.0075)^2 + m(1.0075) + 5000(1.0075) - m \quad \textcircled{1}$$

$$= 5000[1.0075^2 + 1.0075] - m[1.0075 + 1]$$

$$\text{iii. } A_n = 5000[1.0075^n + 1.0075^{n-1} + \dots + 1.0075] - m[1.0075^{n-1} + 1.0075^{n-2} + \dots + 1]$$

$n = 36 \quad A_n = 100\,000$

$$100\,000 = 5000 \times \frac{1.0075(1.0075^{36} - 1)}{1.0075 - 1} \quad \textcircled{1} - m \left[ \frac{1 \times (1.0075^{36} - 1)}{1.0075 - 1} \right] \quad \textcircled{1}$$

$$\bullet \quad m = \$2607.53 \quad \textcircled{1}$$