#### Section A: Multiple Choice (Nos. 1-4, 1-mark each)

### Use the multiple choice answer sheet in the ANSWER BOOKLET

- **1.** A suitable catalyst for the synthesis of ammonia is:
  - (A) platinum
  - (B) iron
  - (C) concentrated H<sub>2</sub>SO<sub>4</sub>
  - (D) nitric acid
- 2. A dry precipitate of Mg(NH<sub>4</sub>)PO<sub>4</sub>.6H<sub>2</sub>O was obtained and weighed . It was found to weigh 6.47 g. How much of this precipitate is phosphorus?
  - (A) 0.82 g
  - (B) 1.46 g
  - (C) 0.03 g
  - (D) 1.64 g
- 3. Damage to the Earth's stratospheric ozone has mainly been due to a certain group of compounds. Which of the compounds given below is an example of this group of compounds?
  - (A) CClF<sub>3</sub>
  - (B) CCl<sub>2</sub>FH
  - (C)  $CF_2I_2$
  - (D) CF<sub>3</sub>H
- 4. Which of the following **IUPAC** names is correct for the compound given below?

- (A) 1,1,6-trifluoro-4,6-dichlorohexane
- (B) 4,6-dichloro-1,1,6-trifluoro-2-hexene
- (C) 1,1,6-trifluoro-4,6-dichlorohexene
- (D) 4,6-dichloro-1,1,6-trifluoro-1-hexene

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

Sample:  $2 + 4 = (A) \ 2 (B) \ 6 (C) \ 8 (D) \ 9$  $A \bigcirc B \bigcirc C \bigcirc D \bigcirc$ 

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word correct and drawing an arrow as follows.



#### **Section A**

### **Multiple Choice Answer Sheet**

- 1. AO BO CO DO
- $2. \hspace{0.5cm} A \hspace{0.1cm} O \hspace{0.5cm} B \hspace{0.1cm} O \hspace{0.5cm} C \hspace{0.1cm} O \hspace{0.5cm} D \hspace{0.1cm} O$
- 3. AO BO CO DO
- 4. AO BO CO DO

<b>Section B.</b> Answer the questions in the spaces provided.	Show all relevant
working in questions involving calculations	

MARKS

5. Describe	a chemical test and the result to identify	
(a) copper	ions	1
	ons	1
(c) calciur	m ions	.1
6. A studer 3.2 g of solution and weig	nt investigated the sulfate content of a fertiliser. Firstly, he dissolved fertiliser in distilled water. Then, he added barium chloride until no further precipitate formed. He, then filtered, washed, dried ghed the precipitate.  a net ionic equation for the formation of the precipitate.	1
	student recovered 5.6 g of barium sulfate. What percentage of this ser is sulfate ions?	Ž
	student assumed the original fertiliser consists of ammonium e only, what is the percentage of nitrogen in the fertiliser?	,

	Student No	••
	MAR	KS
7.	(a) List three different chemical occupations	1
	(b) (i) Choose one of these occupations and outline the role of the chemist.	1
	(ii) Explain a chemical principle used by this chemist.	1
8	Identify the origins of minerals in oceans.	2

<b>9.</b> (a) Use equations to show the destruction of ozone in the stratosphere by a CFC.	MARKS
(b) Explain the importance of the ozone layer to life on Earth.	1
10. Describe two methods which you would use, including details of the preparation which you would do to determine the dissolved solids content of a sample of river water.	3

		M	ARKS
11.	• (a) Draw the Lewis electron dot structures for the oxygen free radical.	e oxygen molecule and the	1
	oxygen molecule ox	xygen free radical	
(	(b) On the basis of molecular structure and bonding	g, explain the difference in:	
	(i) chemical reactivity of ozone and oxygen (C	$O_2$ )	1
			•••
			••••
			••••
	(ii) one physical property of ozone and oxygen	$\cup$ (O <sub>2</sub> )	1
			••••
			••••

Student No. .....

A END OF TEST

# **Section A:** *Multiple Choice* (Nos. 1-4, 1-mark each) MARKING GUIDELINES

Use the multiple choice answer sheet in the ANSWER BOOKLET

- **1.** A suitable catalyst for the synthesis of ammonia is:
  - (A) platinum
  - (B) iron
  - (C) concentrated H<sub>2</sub>SO<sub>4</sub>
  - (D) nitric acid
- 2. A dry precipitate of Mg(NH<sub>4</sub>)PO<sub>4</sub>.6H<sub>2</sub>O was obtained and weighed . It was found to weigh 6.47 g. How much of this precipitate is phosphorus?
  - (A) 0.82 g
  - (B) 1.46 g
  - (C) 0.03 g
  - (D) 1.64 g
- 3. Damage to the Earth's stratospheric ozone has mainly been due to a certain group of compounds. Which of the compounds given below is an example of this group of compounds?
  - (A) CClF<sub>3</sub>
  - (B) CCl<sub>2</sub>FH
  - (C)  $CF_2I_2$
  - (D) CF<sub>3</sub>H
- 4. Which of the following **IUPAC** names is correct for the compound given below?

- (A) 1,1,6-trifluoro-4,6-dichlorohexane
- (B) 4,6-dichloro-1,1,6-trifluoro-2-hexene
- (C) 1,1,6-trifluoro-4,6-dichlorohexene
- (D) 4,6-dichloro-1,1,6-trifluoro-1-hexene

## **Answer Booklet for Sections A and B INSTRUCTIONS**

Student No.....Possible Answer.

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

Sample:  $2 + 4 = (A) \ 2 (B) \ 6 (C) \ 8 (D) \ 9$  $A \bigcirc B \bigcirc C \bigcirc D \bigcirc$ 

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

A ● B 💥 C ○ D ○

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word correct and drawing an arrow as follows.



#### **Section A**

### **Multiple Choice Answer Sheet**

**Outcomes** 

H7	DO	СО	В І	ΑO	1.
H7	DO	СО	ВО	A <b>I</b>	2.
H7, H8, H9,H16	DO	СО	ВО	A <b>I</b>	3.
H13	D <b>I</b>	СО	ВО	ΑО	4.

#### Section B.

5. Describe a chemical test and the result to identify

Outcomes:H7,H13

(a) copper ions

Criteria	Mark
Add NaOH solution to form a blue precipitate, then NH <sub>3</sub>	1
solution to form a deep blue solution	

(b) lead ions

/	
Criteria	Mark
Add KI solution to form a yellow precipitate <b>or</b> add KCl	1
solution to form a white precipitate soluble in hot water.	

(c) calcium ions

Criteria	Mark
Add $Na_2C_2O_4$ solution to form a white precipitate or $NaF$	1
solution to form a white precipitate.	

**6.** A student investigated the sulfate content of a fertiliser. Firstly, he dissolved 3.2 g of fertiliser in distilled water. Then, he added barium chloride solution until no further precipitate formed. He, then filtered, washed, dried and weighed the precipitate.

#### Outcomes: H10, H11, H13

(a) Write a net ionic equation for the formation of the precipitate.

Criteria	Mark(s)
$Ba^{2+} + SO_4^{2-} \otimes BaSO_4(s)$	1

1

2

2

(b) If the student recovered 5.6 g of barium sulfate. What percentage of this fertiliser is sulfate ions?

$$mol \, BaSO_4 = \frac{mass}{mm} = \frac{5.6}{137.3 + 32.06 + 4(16.00)}$$

$$mol \, BaSO_4 = 0.02399 \quad \text{$>$ 0.024 = mol \, SO_4^{2^-}$}$$

$$mass \, SO_4^{2^-} = mol \, SO_4^{2^-} x \, molar \, mass \, SO_4^{2^-} = 0.024 \, x \, 96.06 = \textbf{2.3 g}$$

$$\%SO_4^{2^-} = (mass \, SO_4^{2^-} / mass \, fertilizer) \, x \, 100\% = \textbf{72\%}$$

(c) If the student assumed the original fertiliser consists of ammonium sulfate only, what is the percentage of nitrogen in the fertiliser?

$$(NH_4)_2SO_4$$
 ®  $2NH_4^+ + SO_4^{2-}$   
 $\setminus moles NH_4^+ = 2 \text{ x moles } SO_4^{2-}$   
 $mols NH_4^- - 2 \text{ x } 0.024 = 0.048 = moles N$   
 $\% N = ((moles N \text{ x } At. \text{ No. } N)/mass \text{ fertilizer}) \text{ x } 100\%$   
 $= \frac{0.048 \text{x} 14.01}{3.2} \text{ x } 100\% = 21\%$ 

#### **OUTCOMES: H15**

### 7. (a) List three different chemical occupations

Criteria	Mark
Any of the following or other reasonable types: <i>polymer</i> ,	1
organic, physical, analytical, environmental, industrial, cereal,	
etc.	

### (b) (i) Choose one of these occupations and outline the role of the chemist.,

#### **OUTCOMES:H13**

Criteria	Mark(s)
Any appropriate answer, for example:	1
An analytical chemist in a pharmaceutical company may	
monitor the acetylsalicylic acid content of tablets or	
preparations	

# (ii) Explain a chemical principle used by this chemist. OUTCOMES:H13

Criteria	Mark(s)
Any reasonable answer for example: The chemist may be using	
colorimetric methods to quantitate the aspirin, in which case, the	
aspirin is hydrolysed, allowed to react with $Fe^{3+}$ and the colour	
that develops is measured in a colorimeter. The principle	
behind this is Beer's law or the fact that the extent of absorption	1
of light depends on the number of particles absorbing the the	
light and hence for a given volume of solution, depends on the	
concentration.	

#### **8.** Identify the origins of minerals in oceans. **H8**

Criteria	Mark(s)
1 mark for each identified source	
Minerals may come from runofffs as water percolates through soil,	1
dissolving minerals on the way to the ocean	
Hydrothermal vents in midocean ridges is a conduit for hot water	
(heated up by molten magma) allowing it to percolate through	1b
rocks, dissolving minerals and at the same time allowing the	
diffusion of the dissolved minerals to the greater bulk of ocean	
water	

**9.** (a) Use equations to show the destruction of ozone in the stratosphere by a CFC.

**OUTCOMES: H16,H7** 

Criteria	Mark(s)
$CCl_3F + uv \ light \rightarrow Cl + CCl_2F$	1
$Cl + O_3 \rightarrow ClO + O_2$	1
$ClO + O \rightarrow O_2 + Cl$	1

(b) Explain the importance of the ozone layer to life on Earth. OUTCOMES: H16, H7, H9

Criteria	Mark(s)
Because of its ability to initate bond breakage and bond	
formation in substances such DNA of the cell, UV rays damage,	
alter or even kill organisms in various ways. The ozone layer	1
filters off most of these harmful UV rays, protecting the life	
forms on Earth.	

**10**. Describe two methods which you would use, including details of the preparation which you would do to determine the dissolved solids content of a sample of river water.

**OUTCOMES: H13,H14, H11** 

Criteria	Marks
Each method explained earns 1 mark while preparations for the analysis	
such as filtering and cleaning the electrodes earn 1 mark	
In the evaporation method, the water is first filtered, then	
weighed and evaporated to dryness in a weighed container.	
The residue, completely devoid of water is weighed and the	
mass of the container subtracted to get the mass of the residue.	
The concentration in the original sample is calculated by	3
dividing the mass of the residue with the mass of the water.	
In the conductivity technique, the electrodes of the conductivity	
meter are immersed in the water sample and the reading taken.	
Readings obtained are in ppm. The liquid being measured	
should freely come into contact with the electrodes of the meter.	

**11.** (a) Draw the Lewis electron dot structures for the oxygen molecule and the oxygen free radical. **OUTCOMES:** H13, H7.H6

00100MES. 1113, 117,110	
oxygen molecule	oxygen free radical

Criteria	Mark(s)
Corrrect structures	1

(b) On the basis of molecular structure and bonding, explain the difference in:

(i) chemical reactivity of ozone and oxygen (O<sub>2</sub>)

Criteria	Mark(s)
$O_3$ is more reactive than $O_2$ because the single (coordinate)	1
covalent bond in $O_3$ requires less energy to break than the	
double covalent bond in the $O_2$ molecule.	

(ii) one physical property of ozone and oxygen (O<sub>2</sub>)

Criteria	Mark(s)
Higher solubility of ozone in water: Due to the unsymmetrical distribution of the electrons in the ozone molecule, ozone is slightly more polar than oxygen and hence more soluble in water, interacting via dipole-dipole interaction.	1

#### A END OF TEST