SYDNEY GRAMMAR SCHOOL



2014 Half-Yearly Examination

# FORM VI MATHEMATICS 2 UNIT

Wednesday 26th February 2014

## General Instructions

- Writing time 2 hours
- Write using black or blue pen.
- Board-approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

#### Total - 85 Marks

• All questions may be attempted.

#### Section I – 10 Marks

- Questions 1–10 are of equal value.
- Record your solutions to the multiple choice on the sheet provided.

## Section II – 75 Marks

- Questions 11–15 are of equal value.
- All necessary working should be shown.
- Start each question in a new booklet.

## Collection

- Write your candidate number on each booklet and on your multiple choice answer sheet.
- Hand in the booklets in a single wellordered pile.
- Hand in a booklet for each question in Section II, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Place your multiple choice answer sheet inside the answer booklet for Question Eleven.
- Write your candidate number on this question paper and submit it with your answers.

## Checklist

- SGS booklets 5 per boy
- Multiple choice answer sheet
- Candidature 90 boys

Examiner GMC

# **SECTION I - Multiple Choice**

Answers for this section should be recorded on the separate answer sheet handed out with this examination paper.

#### QUESTION ONE

Given that  $y = \frac{1}{x}$ , which of the following statements is true?


(A)  $\frac{dy}{dx} = \frac{1}{x^2}$ 

(B) 
$$\frac{dy}{dx} = -\frac{1}{x^2}$$

(C) 
$$\frac{dy}{dx} = \frac{2}{x^2}$$

(D) 
$$\frac{dy}{dx} = -\frac{2}{x^2}$$

#### QUESTION TWO



The graph of y = f(x) is shown above. Which of the labelled points satisfies f(x) < 0 and f''(x) > 0?

- $(A) \quad A$
- (B) B
- $(C) \quad C$
- $(D) \quad D$

## **QUESTION THREE**


The value of the limit  $\lim_{x\to 3} \frac{x-3}{x^2-9}$  is given by:

(A) 0 (B)  $\frac{1}{3}$ 

(C) 
$$\frac{1}{6}$$

(D) None of the above.

## **QUESTION FOUR**



A linear function y = f(x) is graphed above. Which of the following expressions represents the area of the shaded region?

(A) 
$$\int_{0}^{4} f(x) dx$$
  
(B) 
$$-\int_{0}^{4} f(x) dx$$
  
(C) 
$$2\int_{0}^{2} f(x) dx$$
  
(D) 
$$2\int_{2}^{4} f(x) dx$$


Exam continues overleaf ...

## **QUESTION FIVE**

What are the coordinates of the focus of the parabola  $x^2 = -4y$ ?

- (A) (0,1)
- (B) (1,0)
- (C) (0, -1)
- (D) (-1, 0)

## QUESTION SIX



The diagram above is a graph of y = f(x). For what values of x is the function y = f'(x) positive?

- (A) x > 0
- (B) x < b or x > d
- (C) x > a
- (D) b < x < d


## **QUESTION SEVEN**

What is the value of the definite integral  $\int_0^1 e^{-x} dx$ ?

(A)  $\frac{e-1}{e}$ (B)  $\frac{1-e}{e}$ (C)  $\frac{e+1}{e}$ (D)  $\frac{1}{e}$ 

#### **QUESTION EIGHT**


The graph of  $f(x) = e^x$  is shown below.



Which one of the following statements is false?

- (A) f(x) > 0
- $(B) \quad f'(x) > 0$
- (C) f(x) = f'(x)
- (D) f(-x) = -f(x)

#### **QUESTION NINE**



A parabola with focus S(0, 2) and directrix x = 2 is shown above. Which of the following is the equation of the parabola?

> (A)  $(y-2)^2 = 4(x-1)$ (B)  $(y-2)^2 = -4(x-1)$ (C)  $(y-2)^2 = 8x$ (D)  $(y-2)^2 = -8x$

## QUESTION TEN

The continuous function y = f(x) has the properties that  $\int_{a}^{c} f(x) dx = 7$  and  $\int_{b}^{c} f(x) dx = -4$ . Given that a < b < c, what is the value of  $\int_{a}^{b} f(x) dx$ ? (A) 11 (B) -11 (C) 3 (D) -3

End of Section I

#### **SECTION II - Written Response**

Answers for this section should be recorded in the booklets provided.

Show all necessary working.

Start a new booklet for each question.

#### **QUESTION ELEVEN** (15 marks) Use a separate writing booklet.

- (a) Calculate  $3e^{-2}$  correct to 2 decimal places.
- (b) Differentiate the following with respect to x:
  - (i)  $2x^2 + 5$
  - (ii)  $x^{\frac{1}{3}}$
  - (iii)  $(4x+1)^6$
- (c) Find a primitive for each of the following:
  - (i)  $3x^5$
  - (ii)  $e^{-2x}$
  - (iii)  $\sqrt{x}$
- (d) Write down the equation of the locus of the point P(x, y) that is:
  - (i) 3 units from the point (-2, 1),
  - (ii) 4 units below the line y = 1.
- (e) Sketch a graph of  $y = e^{-x} + 2$  clearly showing the asymptote and y-intercept.
- (f) Consider a curve whose first derivative is given by y' = 3x + 2. For what value of x is the curve stationary?
- (g) Consider a curve whose second derivative is given by y'' = 2x + 4. For what values of 1 x is the curve concave up?

1
1
<b>2</b>

Marks

1

1

1

 $\mathbf{2}$ 

L	T	
	1	
I	<b>2</b>	
	2	]

Exam continues overleaf ...

**QUESTION TWELVE** (15 marks) Use a separate writing booklet.

(a) Simplify 
$$\frac{e^{3x+2}}{e^x}$$
.

Marks

 $\mathbf{2}$ 

 $\mathbf{2}$ 

1

1

1

1

1

1


(b) Differentiate the following with respect to x:

(i) 
$$y = (2x - 1)(3x + 2)$$
  
(ii)  $y = \frac{x^2 - 2}{x}$ 

(iii) 
$$y = \sqrt{x^2 + 1}$$

(c) Evaluate 
$$\int_{-1}^{2} (4 - x^2) dx$$
.

- (d) A parabola has equation  $x^2 = 16y$ .
  - (i) Write down the coordinates of the vertex.
  - (ii) Find the coordinates of the focus.
  - (iii) Find the equation of the directrix.
  - (iv) Sketch the parabola clearly showing the vertex, focus and directrix.



The diagram above shows the graph of y = f(x).

(i) Copy and complete the table below.

x	0	25	50	75	100
f(x)					




**QUESTION THIRTEEN** (15 marks) Use a separate writing booklet.

(a) Consider the function  $y = x^3 - 6x^2 + 9x - 1$ .

(i) Show that 
$$\frac{dy}{dx} = 3(x-1)(x-3)$$
 and find  $\frac{d^2y}{dx^2}$ .

- (ii) Find the coordinates of any stationary points and determine their nature.
- (iii) Find the coordinates of the point of inflexion. You must show that it is a point of inflexion.
- (iv) Sketch the graph of the function, clearly showing all stationary points, the point of inflexion and the *y*-intercept. Do NOT attempt to find any *x*-intercepts.



(b)

The region bounded by  $y = \frac{2}{x}$ , the x-axis and the lines x = 1 and x = 4 is shown above.

Find the volume of the solid generated when this region is rotated about the x-axis.

(c) A curve has gradient function  $\frac{dy}{dx} = 6x^2 - 3$  and passes through the point (1,5). Find the equation of the curve.

(d) Find the value of k if 
$$\int_{2}^{k} (x-1) dx = 4$$
 and  $k > 2$ .

Marks

2 2

 $\mathbf{2}$ 

 $\mathbf{2}$ 


 $\mathbf{2}$ 

 $\mathbf{2}$ 

3

#### **QUESTION FOURTEEN** (15 marks) Use a separate writing booklet.

- (a) Find the equation of the tangent to the curve  $y = 1 e^{-x}$  when x = 1.
- (b) (i) Express the equation  $y^2 + 4y 3x 5 = 0$  in the form  $(y k)^2 = 4a(x h)$ .
  - (ii) Hence find the coordinates of the focus and the equation of the directrix of the parabola  $y^2 + 4y 3x 5 = 0$ .
- (c) Using the quotient rule, or otherwise, find the derivative of  $y = \frac{2x^2}{e^{3x}}$ . Express your **2** answer in simplest form.
- (d)



The diagram above shows the curves  $y = -x^2 + 4x - 1$  and  $y = x^2 - 6x + 7$ .

- (i) By solving a pair of simultaneous equations, show that the points of intersection of the two curves are (1, 2) and (4, -1).
- (ii) Hence calculate the shaded area between the curves.
- (e) (i) Differentiate  $y = e^{x^2}$ .

(ii) Hence evaluate 
$$\int_0^1 6x e^{x^2} dx$$

<b>2</b>	

 $\mathbf{2}$ 

3

Marks




1

**QUESTION FIFTEEN** (15 marks) Use a separate writing booklet.

- (a) Consider the exponential function  $y = e^{-3x}$ .
  - (i) Find y' and y''.
  - (ii) Show that  $y = e^{-3x}$  satisfies the equation 3y = 5y' + 2y''.
- (b) The locus of a point P(x, y) is a circle. The distance of P from A(8, -16) is three times the distance of P from the origin.
  - (i) Find the equation of the locus of P.
  - (ii) Hence write down the centre and radius of the circle described by P.





A closed metal box is in the shape of a prism with a right-angled triangular cross section. The surface area of the box is  $240 \text{ cm}^2$  and the perpendicular sides of the triangular cross section are in the ratio 3 : 4. Let the dimensions of the box be 3x, 4x and h as shown in the diagram above.

- (i) Show that the surface area, S, of the box is given by  $S = 12x^2 + 12xh$ .
- (ii) Show that the volume of the box, V, is given by  $V = 120x 6x^3$ .
- (iii) Hence find the greatest possible volume of the box in exact form.

#### QUESTION CONTINUES ON THE NEXT PAGE

2

1


Marks

	<b>2</b>	
Γ	1	

1
<b>2</b>
<b>2</b>

**QUESTION FIFTEEN** (Continued)

(d)



The curve x = f(y) passes through the points (1,0), (2,2), (4,4) and (8,6) as shown in the diagram above.

(i) Using the trapezoidal rule with four function values, estimate the volume formed by rotating the region bounded by x = f(y), the x-axis, the y-axis and y = 6 about the y-axis.

3

1

(ii) Does the trapezoidal rule under-estimate or over-estimate the volume of the solid formed in part (i)? Justify your answer.

End of Section II

## END OF EXAMINATION

# BLANK PAGE

The following list of standard integrals may be used:

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax \, dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax \, dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax \, dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec^2 ax \, dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} \, dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} \, dx = \ln \left(x + \sqrt{x^2 - a^2}\right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} \, dx = \ln \left(x + \sqrt{x^2 + a^2}\right)$$

NOTE : 
$$\ln x = \log_e x, x > 0$$

SYDNEY GRAMMAR SCHOOL



2014 Half-Yearly Examination FORM VI MATHEMATICS 2 UNIT Wednesday 26th February 2014

- Record your multiple choice answers by filling in the circle corresponding to your choice for each question.
- Fill in the circle completely.
- Each question has only one correct answer.

Question One			
A 🔿	В ()	С ()	D ()
Question '	Γwo		
A 🔿	В ()	С ()	D ()
Question '	Three		
A 🔿	В ()	С ()	D ()
Question 1	Four		
A 🔿	В ()	С ()	D ()
Question 1	Five		
A 🔿	В ()	С ()	D ()
Question Six			
A ()	В ()	С ()	D ()
Question 8	Seven		
A ()	В ()	С ()	D ()
Question 1	Eight		
A 🔿	В ()	С ()	D ()
Question Nine			
A ()	В ()	С ()	D ()
Question '	Ten		
A 🔾	В ()	С ()	D ()

CANDIDATE NUMBER: .....

FORM VI HALF YEARLY 2014 2 UNIT.

B 6. B 1. A 7. B 2. 8. D C 3. 9. B D 4. 10 <u>10.</u> A ٢. C a) 3e<sup>2</sup> = 0.41 (2d.p.) must show 11. correct to 2d.p. b) :)  $ddx(2x^{2}+5) = 4x$  $\checkmark$  $ii) dlan(x') = \frac{1}{3}x'$ or 32<sup>2/3</sup> one mark for 6(4xr1)5 iii)  $d/dn((4x+1)^{6}) = 24(4x+1)^{5}$  $\int \int$ c) i)  $\int 3\pi^{5} dn = \frac{\pi^{6}}{2} + C$ NB: Constant of ./ Integration not ii) (e-2x dn = -2e + c  $\checkmark$ required here.  $\ddot{u} \int x^{\prime / 2} dn = \frac{3}{3} x^{2} + c$ [] d) i)  $(x+2)^{2} + (y-1)^{2} = 9$ ./ ii)  $\gamma = -3$ e) // ^ ' y= e+2  $\sqrt{}$ -1 for incorrect shape, asymptote or missing y-- y=2 \_\_\_\_\_ interept. Do not reed a シメ separah point.

f) let y'= 0 32+2=0  $n = -\frac{2}{3}$ So the curve is stationary when x = - 3/3 V g) let y">0 2x+4>0 2x>-4 x > - 2 So the curve is concare of when x>-2 a)  $\frac{3x+2}{p^{\chi}} = \frac{2x+2}{e}$ 12. b) i)  $d_{d_{\mathcal{H}}}((2x-1)(3x+2))$ =  $d_{d_{\mathcal{H}}}(6x^2+x-2)$  $\sqrt{}$ = 12x + 1ii)  $\frac{d}{du}\left(\frac{x^2-2}{x}\right)$ =  $\frac{d}{du}\left(x-\frac{2}{x}\right)$  $\sqrt{}$  or  $1+2\pi^{-2}$  $= 1 + \frac{2}{2}$  $\begin{array}{rcl} & & & \\ & & & \\ & & & \\ & & = & \frac{d}{du}\left(\left(\chi^{2}+1\right)\right) & & \\ & & & \\ & & = & \frac{1}{2}\left(\chi^{2}+1\right)^{-1/2} \times 2\pi & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & \\$ 

c)  $\int (4-x^2) dx = \left[ 4x - \frac{x^3}{3} \right]^2 \sqrt{\frac{x^3}{3}}$  $= \left(8 - \frac{8}{3}\right) - \left(-4 + \frac{1}{3}\right)$ = 9 i) Vertex(0,0)d) ii) Focus (0,4) iii) Directrix : y=-4 / x2 = 16y 5(0,4) V(0,0) 4--4 i)  $\frac{\pi}{(n)}$   $\frac{25}{50}$   $\frac{50}{15}$   $\frac{15}{15}$   $\frac{15}{15}$ e) 75 100 0 45  $ii) \int f(x) dx = \frac{50-0}{6} \frac{30+4x15+45}{6} + \frac{100-50}{6} \frac{45+4x0+45}{6}$ = 1875

a) y=x3-6x2\*9x-1 13. i)  $\frac{dy}{dn} = 3x^2 - 12x + 9$ = 3(x2-4x+3) = 3 (x-1)(x-3) as regd. ... du2 = 6x - 12 in) let den =0 3(n-1)(x-3) = 0 x = 1, x = 3 7 Statpts at (1,3) and (3,-1)  $\sqrt{y = 3}, y = -1$  $y = 3, \ y = -6$ when  $x = 1, \ \frac{d^2y}{du^2} = -6$ when  $x = 3, \ \frac{d^2y}{du^2} = 6$  y = -6 y = -7 y = -7 y = -6 y = -7 y =So (1,3) is a max iii) let du = 0 6n-12=0  $\frac{\chi = \alpha}{4},$   $\frac{\chi + 1}{2} = \frac{3}{6}$   $\frac{d^{1}}{du^{2}} = -6 \quad 0 \quad 6$   $\frac{\chi}{du^{2}} = -6 \quad \sqrt{6}$   $\frac{\chi}{du^{2}} = -6 \quad \sqrt{6}$   $\frac{\chi}{du^{2}} = -6 \quad \sqrt{6}$   $\frac{\chi}{du^{2}} = -6 \quad \sqrt{6}$ x=d., y=1 So (2,1) is a point of inflexion

y= x3-6x2+9x-1 (1,3) iv) -1 for any missing point or y-intercept. (2,1) ۱ (3,-1) b)  $V = \pi \int y^2 du$ = TTS 422 du  $= -4\pi \left[ x^{-1} \right]_{1}^{4}$  $= -4\pi \left( \frac{1}{4} - 1 \right)$ = 311 So volume is 3TT units? c)  $dn = 6x^2 - 3$ y = 2x3-3x+C must have 2 c um x=1, y=5 S = 2-3+C C=6  $y = 2x^3 - 3x + 6$ d)  $\int_2^k (x-i) dx = 4$  $\left[\frac{3^{2}}{2}-n\right]_{2}^{k}=4$  $(\frac{k^{2}}{2}-k)-(\frac{4}{2}-2)=4$  $k^{2}-2k-8=0$ (k-4)(k+2)=0k=4, k=-2 but k>2 So k=4

. . . . . . .

i) y=1-e-x 14. a)  $dy = e^{-\chi}$   $du = e^{-\chi}$ when  $\chi = 1$ ,  $du = e^{-\chi}$ tangent has grad te passes through (1, 1-te) /  $y - y_1 = m(x - x_1)$   $y - (1 - \frac{1}{2}) = \frac{1}{2}(x - 1)$   $y - 1 + \frac{1}{2} = \frac{1}{2}x - \frac{1}{2}$   $y = \frac{1}{2}x + 1 - \frac{2}{2}$  or x - ey + e - 2 = 0 $y^{2} + 4y - 3x - 5 = 0$ b) i) y + 4y + 4 = 3x + 5 + 4  $(y + 2)^2 = 3(x + 3)$  / for completing the  $(y + 2)^2 = 4x = \frac{3}{4}(x + 3)$  / either of last two lines of. focal length: 3/4 ii) focus:  $(-2^{\frac{1}{4}}, -2)$  directrix:  $x = -3^{\frac{3}{4}}$  $y = \frac{2\chi^2}{e^{3\chi}}$ c) OK to use  $e^{3\pi} \cdot 4\pi - 2\pi$  $(e^{3\pi})^{2}$ dis du =  $2n^2 \cdot 3e^{3x}$ product role. or (4n-6n²)e<sup>-3n</sup>  $\frac{4x-6x^2}{e^{3n}}$  $\frac{\mathcal{R} \times (2-3x)}{\sigma^{3n}}$ 

d) i) let  $-x^{2}+4x-1 = x^{2}-6x+7$ 2x2-10x+8=0  $\chi^2 - 5\kappa + 4 = 0$ (x-4)(x-1)=0x = 1, x = 4So points of intersection are (1,2) and (4,-1) ii) Area = ((-x2+4x-1)-(x2-6x+7)) du =  $\int (-2\kappa^2 + 10\kappa - 8) d\kappa \sqrt{}$  $= \left( \frac{-2x^{3}}{2} + 5x^{2} - 8x \right)^{2}$  $= \left(\frac{-128}{2} + 80 - 32\right) - \left(\frac{-2}{3} + 5 - 8\right)$ = 9 / So Area = 9 units<sup>2</sup>  $d/du(e^{x^{-}}) = 2xe^{x^{+}}$ e)  $\mathbf{i}$ ii) j6xerdu = 3j2xerdu  $= 3[e^{\chi}]_{0}$ = 3(e-i)

ü) V = = = 3x + 4x + h  $= 6 x^{th}$ from i) 240 = 12x2 + 12x4  $20 = \chi^2 + \chi h$  $h = \frac{20 - x^2}{x} /$ So  $V = 6x^2 \times 20 - x^2$  $= 6 \times (20 - \chi^2)$ = 120 x - 6x<sup>3</sup> as required  $\frac{dv}{dn} = \frac{120 - 18n^2}{2}$ let du = 0  $120 - 18x^2 = 0$  $\chi^2 = \frac{20}{3}$ x=+255 but x>0  $\frac{d^2 V}{du^2} = -36\pi$   $\frac{du^2}{ulm} = \frac{25}{\sqrt{3}}, \frac{d^2 V}{du^2} < 0$   $\frac{-1}{\sqrt{3}} \text{ for } not \text{ lesting } not \text{ lesting } not \text{ lesting } not \text{ restriction}$  $V = 120 \times \frac{2\sqrt{5}}{\sqrt{2}} - 6 \times \left(\frac{2\sqrt{5}}{\sqrt{2}}\right)^3$ = 24055 - 2405553 - 353 $= \frac{160\sqrt{5}}{\sqrt{2}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{160\sqrt{5}}{3}$ So marinum volume is 160515 cm.

d) i)  $V = Ti \int x^2 dy$ y 0 2 4 x 1 2 4 x 1 4 16 6 8 64  $\int \chi^2 dy = \frac{2 - 0}{2} \left\{ 1 + 2 \times Y + 2 \times 16 \times 6 + \right\} \sqrt{2}$ = 105 So Volume  $\doteq$  105 TT units<sup>3</sup> ii) The estimation in part i) is an over-estimation as from the perspective of the y-axis, the curve is close to the y axis than the straight line segments formed by joining the points. This relationship is preserved when the x values are squared in order to estimate the volume integral as the x values are larger than or equal to one. v or similar.