



**SYDNEY BOYS HIGH  
SCHOOL  
MOORE PARK, SURRY HILLS**

**2005**

**YEAR 12**

**ASSESSMENT TASK #2**

# Mathematics Extension 1

## General Instructions

- Working time – 90 minutes.
- Reading Time – 5 minutes.
- Write using black or blue pen.
- Board approved calculators may be used.
- All necessary working should be shown in every question if full marks are to be awarded.
- Marks may not be awarded for messy or badly arranged work
- Hand in your answer booklets in 3 sections. Section A (Question 1), Section B (Question 2) and Section C (Question 3)

## Total Marks - 76

- Attempt questions 1 – 3
- All sections are **NOT** of equal value.

Examiner: *A. Fuller*

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate

Total marks - 76

Attempt Questions 1 - 3

All questions are NOT of equal value

Answer each SECTION in a SEPARATE writing booklet.

---

Section A

Marks

Question 1 (22 marks)

(a) Differentiate the following:

i)  $\tan 2x$  1

ii)  $\frac{1}{\frac{x}{e^2}}$  2

iii)  $\ln(1-2x)^2$  2

iv)  $\sin^2(3x)$  2

(b) Find the following:

i)  $\int \cos \frac{x}{2} dx$  1

ii)  $\int e^{1-4x} dx$  1

iii)  $\int \cot x dx$  2

iv)  $\int 3x.e^{x^2} dx$  2

(c) Find  $\lim_{x \rightarrow 0} \frac{\sin 2x}{x}$  1

(d) i) Show that the equation  $x \cdot \ln x - 1 = 0$  has a root between  $x = 1$  and  $x = 2$ . 2

ii) Using  $x = 2$  as the first approximation, apply Newton's method once to obtain a better approximation of this root. (correct to two decimal places) 2

(e) i) Show that  $\frac{2x+1}{x+2} = 2 - \frac{3}{x+2}$  1

ii) Hence or otherwise, find the exact value of  $\int_0^1 \frac{2x+1}{x+2} dx$ . 3

**End of Section A**


**Section B** (Use a SEPARATE writing booklet)

**Marks**

Question 2 (28 marks)

- (a) Consider the function  $f(x) = \log_e(3x-6)$
- State the largest possible domain of  $f(x)$ . 1
  - Sketch the curve  $y = f(x)$ . 2
  - Find the equation of the normal to the curve at the point where  $x = 4$ . 3

(b)




Given that the equation of the above curve is of the form  $y = a \cos(bx)$  find the values of  $a$  and  $b$ . 2

- (c) Consider the function  $f(x) = xe^{-x}$
- Find the coordinates of any stationary points and determine their nature. 2
  - Show that there is a point of inflection at  $x = 2$ . 2
  - Hence sketch the graph of  $y = f(x)$  showing all essential features. 3

- (d) i) Evaluate  $\int_0^{\frac{\pi}{8}} \sin^2 3x dx$  3
- ii) Show that  $\lim_{h \rightarrow 0} \frac{1 - \cos 2h}{h^2} = 2$  2

- (e) MN is a tangent at R and PQRS is a cyclic quadrilateral.  
PS is parallel to QR and QT is parallel to SR



Prove that:

- i) QR bisects  $\angle PQT$  2
- ii)  $MN \parallel PT$  2
- (f) Prove by Mathematical Induction that  $2^{3n-1} + 3$  is divisible by 7 for all positive integers  $n \geq 1$ . 4

**End of Section B**

Section C (Use a SEPARATE writing booklet)

Marks

Question 3 (26 marks)

(a) An eight-person committee is to be formed from a group of 10 women and 15 men. In how many ways can the committee be chosen if the committee must contain:

- |      |                     |   |
|------|---------------------|---|
| i)   | 4 men and 4 women   | 1 |
| ii)  | more women than men | 2 |
| iii) | at least 2 women    | 2 |

(b) Using the results

$$\frac{d(e^x \sin x)}{dx} = e^x \cos x + e^x \sin x$$

$$\text{and } \frac{d(e^x \cos x)}{dx} = e^x \cos x - e^x \sin x$$


evaluate  $\int_0^{\frac{\pi}{2}} e^x \cos x dx$  2

(c) A car can hold six people, three in the front and three in the back. Only two of the six people can drive. In how many different seating arrangements can they complete the journey? 2

(d) The portion of the curve  $y = \sin x + \cos x$  between  $x = 0$  and  $x = \frac{\pi}{2}$  is rotated about the  $x$ -axis. Show that the volume of the solid of revolution generated is  $\frac{\pi}{2}(\pi + 2)$  cubic units. 3

(e) Using a first approximation  $x_1 \neq 0$ , show that the second approximation,  $x_2$ , is such that  $|x_2| > |x_1|$  when using Newton's method to obtain the zero of  $\sqrt[3]{x}$  2

- (f) Prove by Mathematical Induction that 4  
 $\cos x + \cos 3x + \cos 5x + \dots + \cos(2n-1)x = \frac{\sin 2nx}{2\sin x}$ ,  
 for  $n \geq 1$  (where  $n$  is an integer).
- (g) If  $y = \frac{1}{2}(e^x - e^{-x})$ , prove that  $x = \log_e(y + \sqrt{y^2 + 1})$  3
- (h)



ABCD is a cyclic quadrilateral. The diagonals AC and BD intersect at right angles at X. M is the midpoint of BC. MX produced meets AD at N. Prove that:

- i)  $\angle MBX = \angle MXB$  2
- ii) MN is perpendicular to AD 3

**End of paper**



**SYDNEY BOYS HIGH  
SCHOOL  
MOORE PARK, SURRY HILLS**

**2005**

**YEAR 12**

**ASSESSMENT TASK #2**

# Mathematics Extension 1

## Sample Solutions

<b>Section</b>	<b>Marker</b>
<b>A</b>	<b>AMG</b>
<b>B</b>	<b>FN</b>
<b>C</b>	<b>EC</b>



### Question 1

$$(a) (i) \frac{d}{dx} \tan 2x = 2 \sec^2 2x \quad (1)$$

$$(ii) \frac{d}{dx} \left( \frac{1}{e^{x/2}} \right) = \frac{d}{dx} e^{-x/2}$$
$$= -\frac{1}{2} e^{-x/2}$$
$$= \frac{-1}{2e^{x/2}} \quad (2)$$

$$(iii) \frac{d}{dx} \ln(1-2x)^2$$
$$= \frac{1}{(1-2x)^2} \times \frac{d}{dx} (1-2x)^2$$
$$= \frac{2(1-2x) \times -2}{(1-2x)^2}$$
$$= \frac{-4}{1-2x} \quad (2)$$

$$(iv) \frac{d}{dx} \sin^2(3x)$$
$$= 2 \cos(3x) \sin(3x) \times 3$$
$$= 3 \sin(6x) \quad (2)$$

$$(b) (i) \int \cos \frac{x}{2} dx$$
$$= 2 \int \cos \frac{x}{2} dx \quad (1)$$
$$= 2 \sin \frac{x}{2} + C$$

$$(ii) \int e^{1-4x} dx = -\frac{1}{4} e^{1-4x} + C \quad (1)$$

$$(iii) \int \cot x dx = \int \frac{\cos x}{\sin x} dx$$
$$= \ln |\sin x| + C \quad (2)$$

$$(iv) \int 3x \cdot e^{x^2} dx = \frac{3}{2} \int 2x e^{x^2} dx$$
$$= \frac{3}{2} e^{x^2} + C \quad (2)$$

$$(c) \lim_{x \rightarrow 0} \frac{\sin 2x}{x} = 2 \lim_{x \rightarrow 0} \frac{\sin 2x}{2x}$$
$$= 2 \lim_{u \rightarrow 0} \frac{\sin u}{u}$$

(Since if  $u=2x$ , then  $u \rightarrow 0$  as  $x \rightarrow 0$ )

$$= 2 \times 1$$
$$= 2 \quad (1)$$

Qn 1 contd

$$(d)(i) x \cdot \ln x - 1 = 0$$

$$\begin{aligned} \text{When } x=1 \quad \text{LHS} &= 1 \cdot \ln 1 - 1 \\ &= 1 \times 0 - 1 \\ &= -1 \end{aligned}$$

$$\begin{aligned} \text{When } x=2 \quad \text{LHS} &= 2 \ln 2 - 1 \\ &\doteq 0.386 \end{aligned}$$

Since curve is continuous, and sign changes,  $\therefore$  there is a root in  $[1, 2]$  2

$$(ii) f(x) = x \ln x - 1$$

$$\begin{aligned} f'(x) &= x \cdot \frac{1}{x} + 1 \cdot \ln x \\ &= 1 + \ln x \end{aligned}$$

$$\text{Now } z_2 = z_1 - \frac{f(z_1)}{f'(z_1)}$$

$$\text{let } z_1 = 2$$

$$\begin{aligned} z_2 &= 2 - \frac{f(2)}{f'(2)} \\ &= 2 - \frac{2 \ln 2 - 1}{1 + \ln 2} \\ &\doteq 1.77 \end{aligned}$$

2

(2)

$$(e)(i) \frac{2x+1}{x+2} = \frac{2x+4-3}{x+2}$$

$$= \frac{2(x+2)-3}{x+2}$$

$$= 2 - \frac{3}{x+2} \quad \boxed{1}$$

$$(ii) \int_0^1 \frac{2x+1}{x+2} dx = \int_0^1 \left[ 2 - \frac{3}{x+2} \right] dx$$

$$= \left[ 2x - 3 \ln|x+2| \right]$$

$$= (2 - 3 \ln 3) - (0 - 3 \ln 2)$$


$$= 2 + 3 \ln 3 - 3 \ln 2$$

$$= 2 + 3 \ln \frac{3}{2}$$

3

**QUESTION 2**

(a) (i)  $3x - 6 > 0$   
 $x > 2$



(iii)  $\frac{dy}{dx} = \frac{3}{3x-6}$

gradient of tangent =  $\frac{1}{x-2}$

gradient of normal =  $-x+2$

at  $x=4$  grad of normal =  $-2$  and  $y = \log_6 6$

eqn. of normal is  $y - \log_6 6 = -2(x-4)$

$2x + y - 8 - \ln 6 = 0$

(b) period =  $\frac{2\pi}{b} = 8$

$b = \frac{\pi}{4}$

$3 = a \cos(b \times 1)$

$3 = a \cos \frac{\pi}{4}$

$3 = a \times \frac{1}{\sqrt{2}}$

$a = 3\sqrt{2}$

(c) (i)

$y = xe^{-x}$

$y' = e^{-x} - xe^{-x}$

$y' = e^{-x}(1-x)$

$y'' = e^{-x} - e^{-x} + xe^{-x}$

$y'' = e^{-x}(-1-1+x)$

$y'' = e^{-x}(-2+x)$

$y' = 0$  when  $x = 1$  and at  $x = 1$   $y' = \frac{1}{e}(-1)$ ,  $y'' < 0$


$\therefore$  max. turning point at  $x=1$  (2)

(ii)  $y'' = 0$  when  $x = 2$ ,

$y'' < 0$  when  $x = 1$  and  $> 0$  when  $x = 3$   $\left(\frac{1}{e} \times 1\right)$

$\therefore$  change in concavity- point of inflexion at  $x=2$  (2)

(iii)



(d) (i)  $\int_0^{\frac{\pi}{8}} \sin^2 3x dx = \frac{1}{2} \int_0^{\frac{\pi}{8}} 1 - \cos 6x dx$

$= \frac{1}{2} \left[ x - \frac{1}{6} \sin 6x \right]_0^{\frac{\pi}{8}}$

$= \frac{1}{2} \left[ \frac{\pi}{8} - \frac{1}{6} \sin \frac{3\pi}{4} \right] - \frac{1}{2} [0 - 0]$

$= \frac{1}{2} \left[ \frac{\pi}{8} - \frac{1}{6\sqrt{2}} \right]$

$= \frac{\pi}{16} - \frac{\sqrt{2}}{24}$

(3)

(ii)  $\lim_{h \rightarrow 0} \frac{1 - \cos 2h}{h^2}$

$= \lim_{h \rightarrow 0} \frac{2 \sin^2 h}{h^2}$

$= 2 \times \lim_{h \rightarrow 0} \frac{\sin h}{h} \times \frac{\sin h}{h}$

$= 2 \times 1 \times 1$

$= 2$

(2)

(e) (i) Let  $\angle TQR = a$

$\angle QRS = a$  (alternate  $\angle$ )

$\angle PSR = \pi - a$  (cointerior)

$\angle PQR = a$  (opp.  $\angle$  in cyclic quad)

so PQ bisects  $\angle PQT$

(2)

(ii) Join PT

from (i)  $\angle PQR = \angle PTR = a$  (angles on arc PR)

$\angle RQT = \angle TRN = a$  ( $\angle$  in opp. segment)

so  $PT \parallel MN$  as  $\angle PTR = \angle TRN$  (alt.  $\angle$ s)

(2)

(f) When  $n = 1$ ,  $2^{3 \times 1 - 1} + 3 = 7$

(4)

Statement true for  $n = 1$

Assume statement true for  $n = k$

$2^{3k-1} + 3 = 7A$  where A is a positive integer.

If  $n = k + 1$

$2^{3(k+1)-1} + 3 = 2^{3k-1} \times 2^3 + 3$

$= 8(7A - 3) + 3$

$= 56A - 21$  which is divisible by 7

$2^{3(k+1)-1} + 3$  is divisible by 7

If the statement is true for  $n=k$ , it is true for  $n=k+1$ . So, by Mathematical Induction it is true for any integer  $n \geq 1$

[ Section C. ]

Question (3) (a)

(i)

M (15)	W (10)

(i)  $\binom{15}{4} \times \binom{10}{4} = 1365 \times 210 = 286650$

(ii)

5W 3M  $\binom{15}{3} \binom{10}{5} = 455 \times 252 = 114660$

6W 2M  $\binom{15}{2} \binom{10}{6} = 22050$

7W 1M  $\binom{15}{1} \binom{10}{7} = 1800$

8W  $\binom{10}{8}$  Total = 138555

(iii) 6M 2W, 5M 3W,  
4M 4W, 3M 5W, 2M 6W  
1M 7W, 8W.

(b)  $\binom{15}{6} \binom{10}{2} + \binom{15}{5} \binom{10}{3} + \binom{15}{4} \binom{10}{4}$


$+ \binom{15}{3} \binom{10}{5} + \binom{15}{2} \binom{10}{6} + \binom{15}{1} \binom{10}{7}$

$+ \binom{10}{8}$

$= 225225 + \dots + 45$

$= 1010790$

(c)



2 choices for the driver and the rest of the passengers can arrange themselves

ii  $5!$  ways

(i.e)  $= 2 \times 5! = 240$

$\int (e^{x \cos x} + e^x \sin x) = e^x \sin x$

$\int (e^{x \cos x} - e^x \sin x) = e^{x \cos x}$

$\therefore \int_0^{\pi/2} e^{x \cos x} dx = \left[ \frac{e^x}{2} (\sin x + \cos x) \right]_0^{\pi/2}$

$= \frac{e^{\pi/2}}{2} - \frac{1}{2}$

$= \frac{1}{2} (e^{\pi/2} - 1)$

(d)



$V = \pi \int_0^{\pi/2} y^2 dx = \pi \int_0^{\pi/2} (1 + \sin 2x)^2 dx$

$= \pi \left[ x - \frac{\cos 2x}{2} \right]_0^{\pi/2}$

$= \left( \frac{\pi}{2} + \frac{1}{2} \right) - \left( -\frac{1}{2} \right)$

$= \frac{\pi}{2} (\pi + 2)$

Question (3).

$$(e) f(x) = x^{1/3}$$

$$f'(x) = \frac{1}{3} x^{-2/3}$$

If  $x_1$  is the 1<sup>st</sup> approx.  
then a better approx.  
 $x_2$  is given by

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$= x_1 - \frac{x_1^{1/3}}{(1/3)x_1^{-2/3}}$$

$$= x_1 - 3x_1 = -2x_1$$

$$\therefore |x_2| = |-2x_1| = 2|x_1|$$

i.e.  $|x_2| > |x_1|$ .

(f) Let  $S(n)$  be the statement that

$$\cos x + \cos 3x + \dots + \cos(2n-1)x$$

$$= \frac{\sin 2nx}{2 \sin x}$$

For  $n=1$ ,

$$LHS = \cos x$$

$$RHS = \frac{\sin 2x}{2 \sin x} = \frac{2 \sin x \cos x}{2 \sin x}$$

$$= \cos x$$

$\therefore S(1)$  is true

Assume  $S(k)$  is true

Consider  $n = k+1$

Now  $S(k+1) =$

$$\cos x + \cos 3x + \dots + \cos(2k+1)x$$

$$= \frac{\sin 2kx}{2 \sin x} + \cos(2k+1)x$$

$$= \frac{\sin 2kx + 2 \sin x \cos(2k+1)x}{2 \sin x}$$

$$= \frac{\sin 2kx + 2 \sin x [\cos 2kx \cos x - \sin 2kx \sin x]}{2 \sin x}$$

therefore  $B, C, X$  are concyclic pts.

$$= \frac{\sin 2kx (1 - 2 \sin^2 x) + \cos 2kx \sin 2x}{2 \sin x}$$

$$= \frac{\sin 2kx (\cos 2x) + \cos(2kx) \sin 2x}{2 \sin x}$$

$$= \frac{\sin(2kx + 2x)}{2 \sin x}$$

$$= \frac{\sin 2(k+1)x}{2 \sin x}$$

$S(k+1)$  is true if  $S(k)$  is true.  $\therefore$  By M.I.

$S(n)$  is true  $\forall n \geq 1$ .  
where  $n \in \mathbb{Z}^+$

$$(g) 2y = e^x - \frac{1}{e^x}$$

$$2y e^x = e^{2x} - 1$$

$$\therefore (e^x)^2 - (2y)e^x - 1 = 0$$

$$\Rightarrow e^x = \frac{2y \pm \sqrt{4y^2 + 4}}{2}$$

$$e^x = y \pm \sqrt{y^2 + 1}$$

$$\because e^x > 0 \therefore e^x = y + \sqrt{y^2 + 1}$$

$$\therefore x = \ln(y + \sqrt{y^2 + 1})$$

(h)  $\triangle C X B$  is a rt.  $\triangle$ .

and  $\therefore BC$  subtends a right  $\angle \Rightarrow BC$  is a diameter.

$$\therefore MC = MB$$

$\therefore M$  is the centre

and  $MX$  is also a radian.

i.e.  $\triangle MBX$  is isosceles.

$$\therefore \angle MBX = \angle MXB.$$

(ii) In  $\triangle BCX$  Let  $\angle B = \alpha$ .

$$\therefore \angle BCA = 90 - \alpha \text{ (}\angle \text{ sum of } \triangle \text{)}$$

and  $\angle BCA = \angle BDA = 90 - \alpha$   
( $\angle$  in same seg). Also

$$\angle BXM = \angle DXN = \alpha \text{ (vert. opp)}$$

$$\therefore \angle XND = 90^\circ \Rightarrow MN \perp AD.$$