

# SYDNEY TECHNICAL HIGH SCHOOL

## YEAR 12 HSC ASSESSMENT TASK 2

MARCH 2008

MATHEMATICS

Extension 1

**Time Allowed:** 70 minutes

**Instructions:**

- Attempt all questions
- Start each question on a new page
- Show all necessary working
- The marks for each question are indicated next to the question
- Marks may be deducted for careless or badly arranged work
- Marks indicated are a guide only and may be varied if necessary

Name: \_\_\_\_\_ Teacher: \_\_\_\_\_


| Question 1 | Question 2 | Question 3 | Question 4 | Question 5 | Total |
|------------|------------|------------|------------|------------|-------|
|            |            |            |            |            |       |

**QUESTION 1** (10 Marks)


- a) Find the gradient of the tangent to the curve  $y = \cos^3 x$  at  $x = \frac{\pi}{6}$  **2**
- b) Evaluate  $\lim_{x \rightarrow 0} \frac{\sin 5x}{4x}$  **1**
- c) Write a primitive for  $(5 - 2x)^4$  **2**
- d) Find  $\int \frac{x dx}{(1+x^2)^2}$  by first differentiating  $\frac{x^2}{1+x^2}$  **2**
- e) Evaluate  $\int_0^{\sqrt{3}} \frac{x dx}{\sqrt{1+x^2}}$  using the substitution  $u = 1+x^2$  **3**

**QUESTION 2 (10 Marks)**

- a) The points  $O$  and  $P$  in the plane are  $d$  cm apart. A circle centre  $O$  is drawn to pass through  $P$ , and another circle centre  $P$  is drawn to pass through  $O$ . The two circles meet at  $S$  and  $T$ , as in the diagram.




- i) Explain why angle  $SOT$  is  $\frac{2\pi}{3}$  2
- ii) Hence find the exact area of the shaded region in terms of  $d$  2
- b) The diagram shows the curve  $x = 6y - 5 - y^2$  and the line  $x + y = 5$  3  
The two graphs intersect at  $(0, 5)$  and  $(3, 2)$



Determine the magnitude of the shaded area

- c) For the curve  $y = x^5 - 80x$ ,  $\frac{d^2y}{dx^2} = 0$  at  $(0, 0)$ . Is  $(0, 0)$  a point of inflexion? Justify your answer. 3

**QUESTION 3 (10 Marks)**



This is the graph of  $y = f''(x)$

- i) Find the equation of  $f'(x)$  if there is a stationary point at  $(1,4)$  **2**
- ii) What is the nature of the stationary point at  $(1,4)$ ? Give a reason. **1**
- b)
- i) Sketch on the same diagram the graphs of  $y = 2 \sin x$  and  $y = \cos 2x$  for  $0 \leq x \leq 2\pi$  **3**
- ii) Use your graph or otherwise determine a value for  $d$ , where  $d$  is an integer, so that the equation  $2 \sin x - \cos 2x = d$  has 4 solutions in the interval  $0 \leq x \leq 2\pi$  **1**
- c) Find  $\int x^3 (x^2 + 1)^2 dx$  by using the substitution  $u = x^2 + 1$  **3**

### QUESTION 4 (10 Marks)

a) For the curve  $y = \frac{x^2}{1+x}$


i) Find the co-ordinates of the stationary points and determine their nature. 3

ii) Given  $y = \frac{x^2}{1+x}$  can be written as  $y = x - 1 + \frac{1}{x+1}$

Write down the equations of any asymptotes 2

iii) Sketch the curve showing the stationary points and the asymptotes 2


b) 3



The shaded region in the diagram is bounded by the curve  $y = x - \frac{1}{x}$ , the  $x$ -axis and the line  $x = 4$ . Find the volume of the solid of revolution formed when the shaded region is rotated about the  $x$ -axis.

**QUESTION 5 (10 Marks)**

a)



In a quadrilateral  $ABCD$ ,  $BC$  is parallel to  $AD$ , the sides  $AB$ ,  $BC$ ,  $CD$  are each 5m long and the angles  $BAD$ ,  $ADC$  each have size  $\theta$ , as shown in the diagram:


- i) Show that the area of the trapezium is given by the formula 2

$$\text{Area} = 25 \sin \theta (1 + \cos \theta)$$

- ii) Hence find the value of  $\theta$  for which this area is a maximum 4

- b) A hemi-spherical bowl is formed by rotating the semi-circle 4

$y = r - \sqrt{r^2 - x^2}$  about the  $y$  - axis. The bowl contains water up to the height  $h$  where  $0 < h < r$ .



Show that the volume of water in the bowl is  $\frac{\pi h^2(3r-h)}{3}$

Questões

a)  $y = \cos^3 x$   
 $y' = -3 \sin x \cdot \cos^2 x$

at  $x = \pi/6$   
 $y' = -3 \cdot \frac{1}{2} \cdot \left(\frac{\sqrt{3}}{2}\right)^2$   
 $= -\frac{9}{8}$

b)  $\lim_{x \rightarrow 0} \frac{\sin 5x}{4x} = \frac{5}{4}$

c)  $\frac{-(5-2x)^5 + C}{10}$

d)  $\frac{d}{dx} \left( \frac{x^2}{1+x^2} \right) = \frac{(1+x^2) \cdot 2x - x^2 \cdot 2x}{(1+x^2)^2}$   
 $= \frac{2x}{(1+x^2)^2}$

$\therefore \int \frac{x}{(1+x^2)^2} dx = \frac{1}{2} \frac{x^2}{1+x^2} + C$

e)  $u = 1+x^2 \rightarrow du = 2x dx$   
 $x = \sqrt{3} \quad u = 4$   
 $x = 0 \quad u = 1$

$\therefore = \frac{1}{2} \int_0^4 \frac{du}{u^{1/2}}$   
 $= \frac{1}{2} \int_0^4 u^{-1/2} du$   
 $= \frac{1}{2} \left[ 2u^{1/2} \right]_1^4$   
 $= 1$

Question 1

$$\begin{aligned} \text{a) } y &= 6x - 5 - x^2 \\ \text{Area} &= \int_2^5 (6x - 5 - x^2) dx \\ &= \left[ 3x^2 - 5x - \frac{x^3}{3} \right]_2^5 \\ &= \left[ 75 - 25 - \frac{125}{3} \right] - \left[ 12 - 10 - \frac{8}{3} \right] \\ &= 8\frac{1}{3} - \left(-\frac{2}{3}\right) \\ &= 4\frac{1}{2} \end{aligned}$$

$$\begin{aligned} \text{b) Area} &= \int_2^5 (6x - 5 - x^2) dx \\ &= \left[ 3x^2 - 5x - \frac{x^3}{3} \right]_2^5 - \frac{9}{2} \\ &= \left[ 75 - 25 - \frac{125}{3} \right] - \left[ 12 - 10 - \frac{8}{3} \right] - \frac{9}{2} \\ &= 8\frac{1}{3} - \left(-\frac{2}{3}\right) - \frac{9}{2} \\ &= 4\frac{1}{2} \end{aligned}$$

$$\begin{aligned} \text{c) } y' &= 5x^4 - 80 \\ y' &= 0 \text{ for stationary pt.} \end{aligned}$$

$$\therefore 5x^4 - 80 = 0$$

$$5[x^4 - 16] = 0$$

$$\therefore x = \pm 2$$

$\therefore$  stationary pts at  $x = \pm 2$

$\therefore (0,0)$  is a point of inflection as it is between two stationary points on a continuous curve.

$$y'' = 20x^3$$

|     |    |   |   |
|-----|----|---|---|
| x   | -1 | 0 | 1 |
| y'' | -  | 0 | + |

curve changes concavity at either side of  $(0,0)$



$$= r^2 - [r^2 - 2ry + y^2]$$

$$= 2ry - y^2$$

$$\text{Volume} = \pi \int_0^h (2ry - y^2) dy$$

$$= \pi \left[ ry^2 - \frac{y^3}{3} \right]_0^h$$

$$= \pi \left[ rh^2 - \frac{h^3}{3} \right]$$

$$= \pi \left[ \frac{3rh^2 - h^3}{3} \right]$$

$$= \frac{\pi h^2 [3r - h]}{3}$$

Teacher's Name

Student's Name

Question 3

a) i)  $f(x) = 2x^2 + 4x - 3$

$$f'(x) = 4x + 4$$


at  $(1, 4)$   $0 = 4 + 4 + c$

$$c = -8$$

$$\therefore f'(x) = -4x^2 + 8x - 8$$

ii) at  $x = 1$   $f''(x) > 0$

$\therefore$  minimum turning point



ii)  $2 \sin x = \cos 2x + c$   
 $\therefore c = -1$

c)  $u = x^2 + 1 \rightarrow du = 2x dx$   
 $x^2 = u - 1$

$$\begin{aligned} \therefore \int x^3 (x^2 + 1)^2 dx &= \int x^2 \cdot x \cdot (x^2 + 1)^2 dx \\ &= \int (u - 1) u^2 \cdot \frac{du}{2} \\ &= \frac{1}{2} \int u^3 - u du \\ &= \frac{1}{2} \left[ \frac{u^4}{4} - \frac{u^3}{3} \right] + c \\ &= \frac{(x^2 + 1)^4}{8} - \frac{(x^2 + 1)^3}{6} + c \end{aligned}$$

## Question 4

$$a) i) y' = \frac{(1+x) \cdot 2x}{(1+x)^2}$$

$$= \frac{2x + 2x^2}{(1+x)^2}$$

$y' = 0$  for stationary point

$$\therefore 2x + 2x^2 = 0$$


$$x(2 + 2x) = 0$$

$$x = -2 \text{ or } 0$$

|    |    |    |      |    |   |   |
|----|----|----|------|----|---|---|
| x  | -3 | -2 | -1.5 | -1 | 0 | 1 |
| y' | +  | 0  | -    | -  | 0 | + |

$\therefore$  maximum at  $(-2, -4)$  minimum at  $(0, 0)$

ii)  $y = x - 1$  at  $x = -1$



$$b) \text{ Volume} = \pi \int_1^4 \left(x - \frac{1}{x}\right)^2 dx$$

$$= \pi \int_1^4 \left(x^2 - 2 + \frac{1}{x^2}\right) dx$$

$$= \pi \left[ \frac{x^3}{3} - 2x - \frac{1}{x} \right]_1^4$$

$$= \pi \left[ \left(\frac{64}{3} - 8 - \frac{1}{4}\right) - \left(\frac{1}{3} - 2 - 1\right) \right]$$

$$= \frac{63\pi}{4}$$

## Question 5

a)



$$h = 5 \sin \theta \text{ and } d = 5 \cos \theta$$

$$\therefore \text{Area} = \frac{5 \sin \theta}{2} [5 + 5 + 10 \cos \theta]$$

$$= 25 \sin \theta [1 + \cos \theta]$$

$$iii) A' = 25 [ (1 + \cos \theta) \cos \theta + \sin \theta \cdot \sin \theta ]$$

$$= 25 [ \cos \theta + \cos^2 \theta - \sin^2 \theta ]$$

$$= 25 [ \cos \theta + \cos^2 \theta - [1 - \cos^2 \theta] ]$$

$$= 25 [ 2 \cos^2 \theta + \cos \theta - 1 ]$$

For maximum  $A' = 0$

$$\therefore 25 [ 2 \cos^2 \theta + \cos \theta - 1 ] = 0$$

$$25 [ (2 \cos \theta - 1)(\cos \theta + 1) ] = 0$$

$$\therefore \cos \theta = \frac{1}{2} \text{ or } \cos \theta = -1 \text{ (out of domain)}$$

$$\theta = \frac{\pi}{3}$$

|          |                 |                 |                 |
|----------|-----------------|-----------------|-----------------|
| $\theta$ | $\frac{\pi}{6}$ | $\frac{\pi}{3}$ | $\frac{\pi}{2}$ |
| $A'$     | +               | 0               | +               |

$\therefore$  max area when  $\theta = \frac{\pi}{3}$