



**PETRUS KY
COLLEGE**
NEW SOUTH WALES

in partnership
with



**VIETNAMESE COMMUNITY
IN AUSTRALIA**
NSW CHAPTER

JULY 2006

MATHEMATICS EXTENSION 2

PRE-TRIAL TEST

HIGHER SCHOOL CERTIFICATE (HSC)

Student Number:

--	--	--	--

Student Name:

General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question
- Write your Student Number and your Name on all working booklets

Total marks – 96

- Attempt Questions 1–8
- Question 8 is optional
- All questions are of equal value

Blank Page

Marks

Question 1

12

(A) Integrate the following,

(i) $\int \frac{\sin 2x}{\sqrt{1 - \cos 2x}} dx$ 1

(ii) $\int \frac{dx}{x\sqrt{x^6 - 4}}$ (Let $x^3 = 2 \sec u$) 2

(iii) $\int_{-1}^1 \frac{2x}{(x^2 + 2x + 5)^2} dx$ 2

(iv) $\int \frac{dx}{e^x \sqrt{1 - e^{-2x}}}$ 2

(B) By substituting $x = a - y$, show that 1

$$\int_0^a f(x) dx = \int_0^a f(a - x) dx$$

Hence use this result to evaluate.

(i) $\int_0^1 x(1-x)^{12} dx$ 2

(ii) $\int_0^{\frac{\pi}{2}} \frac{\cos x - \sin x}{1 + \sin 2x} dx$ 2

Question 2**12**

(A) Given $Z_1 = i\sqrt{2}$ and $Z_2 = \frac{2}{1-i}$

4

(i) Express Z_1 and Z_2 in the modulus/argument form.

(ii) If $Z_1 = w.Z_2$ express w in the modulus/argument form.

(iii) Show Z_1 , Z_2 and $Z_1 + Z_2$ on an Argand diagram.

Hence show that

$$\text{Arg}(Z_1 + Z_2) = \frac{3\pi}{8}$$

Use the diagram to find the exact value of $\tan \frac{3\pi}{8}$

(B) If Z_1, Z_2 are complex numbers, prove that

4

$$\frac{|Z_1|}{|Z_2|} = \frac{|Z_1|}{|Z_2|}$$

Given a complex number $Z = \frac{c+2i}{c-2i}$ where c is real.

Find $|Z|$ and hence describe the exact locus of Z if c varies from -1 to 1 .

(C) If $w = 2\sqrt{3}i - 2$, find $|w|$ and $\arg w$, then indicate on an Argand diagram the complex number w , \bar{w} , iw , $\frac{1}{w}$, $-w$.

4

Show that $w^2 = 4\bar{w}$

Prove that w is a root of the equation $Z^3 - 64 = 0$. Find other roots.

Question 3**12**

(A) P is the point $(3\cos \theta, 2\sin \theta)$ and Q is the point $(3\sec \theta, 2\tan \theta)$.

Sketch the curves which are the loci, as θ varies, of P and Q, marking on them the range of positions occupied by P and Q respectively as θ varies from $\frac{\pi}{2}$ to π .

(i) Prove that for any value of θ , the line PQ passes through one of the common points of the 2 curves. 2

(ii) Show that the tangent at P to the first curve meets the tangent at Q to the second curve in a point which lies on the common tangent to the two curves at their other common point. 2

(iii) Prove that the two curves have the same length of the latus rectum. 2

(B) Show that the condition for a straight line $y = mx + c$ to touch the ellipse E of 3

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad \text{is} \quad c^2 = b^2 + a^2 m^2$$

Hence show that the locus of the point P(x, y) from which the 2 tangents to the ellipse E

$\frac{x^2}{16} + \frac{y^2}{9} = 1$ are perpendicular together, is a curve with the centre at the origin and a radius of 5.

Question 4

12

(A) A function is defined with the polar equation as follows:

$$\begin{cases} x = 8 \cos^3 \theta \\ y = 8 \sin^3 \theta \end{cases} \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

(i) Find $\frac{dy}{dx}$ in term of θ and show that the graph of this function touches the x and y axis. Sketch the curve. 3

(ii) Show that the Cartesian equation of that function is 3

$$x^{\frac{2}{3}} + y^{\frac{2}{3}} = 4$$

Show that the equation of the tangent to the curve at the point P(x_0 , y_0) is:

$$y_0^{1/3} x + x_0^{1/3} y = 4x_0^{1/3} \cdot y_0^{1/3}$$

Prove that the segment intercepted on this tangent by the coordinate axis is independent of the position of P on the curve.

(B) Consider the function $f(x) = 2 - \frac{4x}{x^2 + 1}$

(i) Show that the function is always positive for any value of x. 1

(ii) Find the asymptote (if any) and the stationary point of that curve. 1

(iii) Sketch the curve $y = f(x)$. 1

(iv) On a separate diagram, sketch the relating curves:

a) $y = f(|x|)$ 1

b) $y = \frac{1}{f(x)}$ 1


c) $y = \ln f(x)$ 1

Question 5

12

(A) The area bounded by the curve $y=\sin x$, the two lines $y=-x$ and $x=\pi$ is rotated about the line $y=-x$. Find the volume of the solid shape of that formation. 6


(B) A cylindrical timber is chopped at two ends by 2 planes which are inclined 60° and 45° respectively. If the two ends are tilt toward each other and the shortest length of 2 ends is 120cm. Find the volume of that timber. Give the radius of timer is 10cm. 6



Question 6

12

(A)



Two circles intersect at A and D. P is the point on the major arc of one circle. The other circle has the radius r . PA produced and PD produced meet the other circle at B and C respectively. Let $\angle APD = \alpha$ and $\angle ACD = \beta$.

(i) Show that $BC = 2r \sin(\alpha + \beta)$. 2

(ii) As P moves along the major arc AD on its circle, show that the length of the chord BC is independent of the position of P. 2

(iii) If the 2 circles have equal radii, show that $BC = 2 \cos \alpha \cdot AD$ 2

(B) P is any point $(ct, c/t)$ on the Hyperbola $xy = c^2$, whose centre is O.

(i) M and N are perpendicular roots of P to the 2 asymptotes. Prove that PM.PN is constant. 1

(ii) Find the equation of tangent at P, and show that OP and this tangent are equally inclined to the asymptotes. 2

(iii) If the tangent at P meet the asymptotes at A and B, and Q is the fourth vertex of the rectangle OAQB, find the locus of Q. 1

(iv) Show that $PA=PB$ and hence conclude that the area of ΔOAB is independent of position of P. 2

Question 7**12**

- (A) If the polynomial $P(x) = x^4 - 4x^3 + 11x^2 - 14x + 10$ has two zeros $(a + ib)$ and $(a - 2ib)$ where a and b are real, then find the values of a and b . 4

Hence find the zeros of $P(x)$ over the complex field C , and express $P(x)$ as the product of 2 quadratic factors with rational coefficients.

- (B) Show that if the polynomial $P(x) = 0$ has a root α of multiplicity m , then $P'(x) = 0$ has a root α of multiplicity $(m - 1)$. 4

Given that $P(x) = x^4 + x^3 - 3x^2 - 5x - 2 = 0$ has a 3-fold root, find all the roots of $P(x)$.

- (C) Find the cubic roots of unity and express them in the form $r(\cos \theta + i \sin \theta)$.

Show these roots on an Argand diagram.

If w is one of the complex roots, prove that the other root is w^2 and show that

$$1 + w + w^2 = 0.$$

- (i) Prove that if n is a positive integer, then $1 + w^n + w^{2n} = 3$ or 0 depending on whether n is or is not a multiple of 3. 2

- (ii) If $x = a + b$, $y = aw + bw^2$ and $z = aw^2 + bw$, show that 2
- $$z^2 + y^2 + x^2 = 6ab$$

Question 8 - Optional**12**

(A) A particle is projected vertically upward with initial speed u . The air resistance is proportional to the speed of the particle.

(a) If $\ddot{x} = -(g + kv)$ with k is the constant, then find the maximum height reached by the particle and the time to do so. 2

(b) Set up the differential equation for the downward motion. 2

(c) Show that the particle returns to its point of projection with speed v given by 2

$$k(u + v) = g \log_e \left[\frac{-g + ku}{-g - kV} \right]$$

(B) Show that for $n \geq 1$ 3

$$1. \ln \frac{2}{1} + 2 \ln \frac{3}{2} + 3 \ln \frac{4}{3} + \dots + n \ln \left(\frac{n+1}{n} \right) = \ln \left(\frac{(n+1)^n}{n!} \right)$$

(C) By using the induction method, prove that $(35)^n + 3 \times 7^n + 3 \times 5^n + 6$ is divisible by 12 for $n \geq 1$ 3

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1}x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a}e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$

Blank Page



**PETRUS KY
COLLEGE**
NEW SOUTH WALES

in partnership
with



**VIETNAMESE COMMUNITY
IN AUSTRALIA**
NSW CHAPTER

JULY 2006

MATHEMATICS EXTENSION 2

PRE-TRIAL TEST

SOLUTION

HIGHER SCHOOL CERTIFICATE (HSC)

Student Number:

--	--	--	--

Student Name: _____

General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question
- Write your Student Number and your Name on all working booklets

Total marks – 96

- Attempt Questions 1–8
- Question 8 is optional
- All questions are of equal value

Question 1

12

(A) Integrate the following,

(i) $\int \frac{\sin 2x}{\sqrt{1-\cos 2x}} dx$

1

Let $u = 1 - \cos 2x$, $\frac{du}{dx} = 2\sin 2x$.

$$\begin{aligned} \int \frac{\sin 2x \cdot dx}{\sqrt{1-\cos 2x}} &= \frac{1}{2} \int \frac{du}{\sqrt{u}} = \sqrt{u} + C \\ &= \frac{1}{2} \sqrt{1-\cos 2x} + C \end{aligned}$$

(ii) $\int \frac{dx}{x\sqrt{x^6-4}}$ (Let $x^3 = 2 \sec u$)

2

Let $x^3 = 2 \sec u \longrightarrow u = \cos^{-1} \left(\frac{2}{x^3} \right)$

$x = (2 \sec u)^{1/3}$

$\frac{dx}{du} = \frac{2 \sec u \cdot \tan u \cdot (2 \sec u)^{-2/3}}{3}$

$= \frac{\tan u \cdot (2 \sec u)^{1/3}}{3}$

$\therefore \int \frac{dx}{x\sqrt{x^6-4}} = \frac{1}{3} \int \frac{\tan u \cdot (2 \sec u)^{1/3} du}{(2 \sec u)^{1/3} \sqrt{4 \sec^2 u - 4}}$

$= \frac{1}{3} \int \frac{\tan u \cdot du}{2 \sqrt{\sec^2 u - 1}}$

$= \frac{1}{6} \int du = \frac{1}{6} u + C$

$\therefore \int \frac{dx}{x\sqrt{x^6-4}} = \frac{1}{6} \cos^{-1} \left(\frac{2}{x^3} \right) + C$

$$(iii) \int_{-1}^1 \frac{2x}{(x^2+2x+5)^2} dx$$

$$\int_{-1}^1 \frac{2x \cdot dx}{(x^2+2x+5)^2} = \int_{-1}^1 \frac{2x dx}{((x+1)^2+4)^2}$$

• Let $x+1 = 2 \tan \theta \longrightarrow x = 2 \tan \theta - 1$
 $\frac{dx}{d\theta} = 2 \sec^2 \theta \cdot d\theta$

• Change the range

- When $x = -1$, $\tan \theta = 0$, $\theta = 0$

- When $x = 1$, $\tan \theta = 1$, $\theta = \frac{\pi}{4}$

$$\begin{aligned} \therefore \int_{-1}^1 \frac{2x dx}{(x^2+2x+5)^2} &= \int_0^{\pi/4} \frac{2(2 \tan \theta - 1) \cdot 2 \sec^2 \theta \cdot d\theta}{(4 \tan^2 \theta + 4)^2} \\ &= \frac{4}{16} \int_0^{\pi/4} \frac{(2 \tan \theta - 1) \sec^2 \theta d\theta}{\sec^4 \theta d\theta} \\ &= \frac{1}{4} \int_0^{\pi/4} 2 \tan \theta \times \cos^2 \theta - \cos^2 \theta d\theta \\ &= \frac{1}{4} \int_0^{\pi/4} 2 \sin \theta \cdot \cos \theta - \frac{1}{2} (1 + \cos 2\theta) d\theta \\ &= \frac{1}{4} \int_0^{\pi/4} \sin 2\theta - \frac{1}{2} - \frac{1}{2} \cos 2\theta d\theta \\ &= -\frac{1}{4} \left[\frac{1}{2} \cos 2\theta + \frac{1}{2} \theta + \frac{1}{4} \sin 2\theta \right]_0^{\pi/4} \\ &= -\frac{1}{4} \left[\left(\theta + \frac{\pi}{8} + \frac{1}{4} - \frac{1}{2} \right) \right] \\ \underline{\text{Answer}} &= \frac{1}{16} - \frac{\pi}{32} \end{aligned}$$

$$(iv) \int \frac{dx}{e^x \sqrt{1-e^{-2x}}}$$

$$\int \frac{dx}{e^x \sqrt{1-e^{-2x}}}$$

$$\text{Let } u = e^{-x}$$

$$\frac{du}{dx} = -e^{-x} \quad , \quad -du = \frac{dx}{e^x}$$

$$\therefore \int \frac{dx}{e^x \sqrt{1-e^{-2x}}} = - \int \frac{du}{\sqrt{1-u^2}} = \cos^{-1} u + C$$

$$= \cos^{-1} \left(\frac{1}{e^x} \right) + C$$

(B) By substituting $x = a - y$, show that

1

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

Hence use this result to evaluate.

Definite integral

$$\text{Show that } \int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$\text{Let } x = a - y \quad \Bigg| \quad \begin{array}{l} \text{when } x = 0, y = a \\ x = a, y = 0 \end{array}$$

$$\frac{dx}{dy} = -1$$

$$\therefore \int_0^a f(x) dx = - \int_a^0 f(a-y) dy$$

$$= \int_0^a f(a-y) dy$$

$$= \int_0^a f(a-x) dx$$

(i) $\int_0^1 x(1-x)^{12} dx$

2

$$\begin{aligned}\int_0^1 x(1-x)^{12} dx &= \int_0^1 (1-x)(1-(1-x))^{12} dx \\ &= \int_0^1 (1-x)x^{12} dx \\ &= \int_0^1 x^{12} - x^{13} dx \\ &= \left[\frac{x^{13}}{13} - \frac{x^{14}}{14} \right]_0^1 \\ &= \frac{1}{13} - \frac{1}{14} = \frac{1}{182}\end{aligned}$$

(ii) $\int_0^{\pi/2} \frac{\cos x - \sin x}{1 + \sin 2x} dx$

2

$$\begin{aligned}\int_0^{\pi/2} \frac{\cos x - \sin x}{1 + \sin 2x} dx &= \int_0^{\pi/2} \frac{\cos(\frac{\pi}{2} - x) - \sin(\frac{\pi}{2} - x)}{1 + \sin 2(\frac{\pi}{2} - x)} dx \\ &= \int_0^{\pi/2} \frac{\sin x - \cos x}{1 + \sin(\pi - 2x)} dx \\ &= \int_0^{\pi/2} \frac{\sin x - \cos x}{1 + \sin 2x} dx\end{aligned}$$

$$\therefore 2 \int_0^{\pi/2} \frac{\cos x - \sin x}{1 + \sin 2x} dx = 0$$

Answer = 0

Question 2

12

(A) Given $Z_1 = i\sqrt{2}$ and $Z_2 = \frac{2}{1-i}$

4

(i) Express Z_1 and Z_2 in the modulus/argument form.

$$\bullet \quad z_1 = \sqrt{2} \operatorname{cis} \frac{\pi}{2}$$

$$z_2 = \frac{2}{1-i} \times \frac{1+i}{1+i} = \frac{2(1+i)}{2} = 1+i$$

$$\bullet \quad z_2 = \sqrt{2} \operatorname{cis} \frac{\pi}{4}$$

(ii) If $Z_1 = w \cdot Z_2$ express w in the modulus/argument form.

Find w if $Z_1 = w \cdot Z_2$

$$w = \frac{z_1}{z_2}$$

$$\therefore |w| = \frac{|z_1|}{|z_2|} = \frac{\sqrt{2}}{\sqrt{2}} = 1$$


$$\begin{aligned} \operatorname{Arg}(w) &= \operatorname{Arg} z_1 - \operatorname{Arg} z_2 \\ &= \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4} \end{aligned}$$

$$\therefore w = \operatorname{cis} \frac{\pi}{4}$$

(iii) Show Z_1 , Z_2 and $Z_1 + Z_2$ on an Argand diagram. Hence show that

$$\operatorname{Arg}(Z_1 + Z_2) = \frac{3\pi}{8}$$

Use the diagram to find the exact value of $\tan \frac{3\pi}{8}$



OPSQ is a rhombus, therefore OS bisects $\angle POQ$

$$\text{hence } \angle SOQ = \frac{\pi}{4} \div 2 = \frac{\pi}{8}$$

$$\begin{aligned}\text{Arg}(z_1 + z_2) &= \angle xOQ + \angle QOS \\ &= \frac{\pi}{4} + \frac{\pi}{8} \\ &= \frac{3\pi}{8}\end{aligned}$$

$$\text{In } \triangle OSR, \tan \angle SOR = \frac{SR}{OR}$$

$$\text{Hence, } \boxed{\tan \frac{3\pi}{8} = \sqrt{2} + 1}$$

(B) If Z_1, Z_2 are complex numbers, prove that

4

$$\frac{|Z_1|}{|Z_2|} = \frac{|Z_1|}{|Z_2|}$$

Given a complex number $Z = \frac{c+2i}{c-2i}$ where c is real.

Find $|Z|$ and hence describe the exact locus of Z if c varies from -1 to 1.

Prove

$$\frac{|z_1|}{|z_2|} = \frac{|z_1|}{|z_2|}$$

$$\text{Let } z_1 = |z_1| \text{cis } \alpha$$

$$z_2 = |z_2| \text{cis } \beta$$

$$\begin{aligned}\frac{z_1}{z_2} &= \frac{|z_1| (\cos \alpha + i \sin \alpha)}{|z_2| (\cos \beta + i \sin \beta)} \times \frac{(\cos \beta - i \sin \beta)}{(\cos \beta - i \sin \beta)} \\ &= \frac{|z_1|}{|z_2|} (\cos(\alpha - \beta) + i \sin(\alpha - \beta))\end{aligned}$$

$$\frac{z_1}{z_2} = \frac{|z_1|}{|z_2|} \text{cis } (\alpha - \beta)$$

$$\text{Therefore, modulus of } \frac{z_1}{z_2} = \frac{|z_1|}{|z_2|}$$

$$\text{let } z = \frac{c+2i}{c-2i}$$


$$\text{let } z_1 = c+2i \quad \text{then } |z_1| = \sqrt{c^2+4}$$

$$z_2 = c-2i \quad \text{then } |z_2| = \sqrt{c^2+4}$$

$$|z| = \frac{|z_1|}{|z_2|} = \frac{\sqrt{c^2+4}}{\sqrt{c^2+4}} = 1$$

Therefore, in general, the locus of z is the circle, centre at origin and radius = 1, equation $x^2 + y^2 = 1$

When the value of c varies from -1 to 1 , Locus of z becomes only part of that circle, that is it is an arc from point A to B which contains the corner $(-1,0)$ as shown in the following figure:



(C) If $w = 2\sqrt{3}i - 2$, find $|w|$ and $\arg w$, then indicate on an Argand diagram the complex number w , \bar{w} , iw , $\frac{1}{w}$, $-w$.

4

$$\text{Show that } w^2 = 4\bar{w}$$


Prove that w is a root of the equation $Z^3 - 64 = 0$. Find other roots.

$$w = 2\sqrt{3}i - 2$$

$$|w| = \sqrt{4 + 12} = 4$$

$$\arg(w) = \tan^{-1}\left(\frac{2\sqrt{3}}{-2}\right) = \frac{2\pi}{3}$$

show in Argand Diagram



• show that $w^2 = 4\bar{w}$

$$w^2 = (2\sqrt{3}i - 2)^2 = -12 - 8\sqrt{3}i + 4 = -(8 + 8\sqrt{3}i)$$

$$4\bar{w} = 4(-2 - 2\sqrt{3}i) = -(8 + 8\sqrt{3}i)$$

∴ $w^2 = 4\bar{w}$

• since $w = 4 \operatorname{cis} \frac{2\pi}{3}$

Then $w^3 = 4^3 \operatorname{cis} 2\pi = 64$

Therefore $w^3 - 64 = 0$

Polynomial equation $w^3 - 64 = 0$ has real coefficients and one complex root $w = -2 + 2\sqrt{3}i$, then it also has other complex root which is the conjugate $\bar{w} = -2 - 2\sqrt{3}i$. The last root is real, ie $w = 4$.

Question 3


12

(A) P is the point $(3\cos \theta, 2\sin \theta)$ and Q is the point $(3\sec \theta, 2\tan \theta)$.

Sketch the curves which are the loci, as θ varies, of P and Q, marking on them the range of positions occupied by P and Q respectively as θ varies from $\frac{\pi}{2}$ to π .

(i) Prove that for any value of θ , the line PQ passes through one of the common points of the 2 curves. 2

a) $P(3\cos \theta, 2\sin \theta)$, $Q(3\sec \theta, 2\tan \theta)$, $\frac{\pi}{2} \leq \theta \leq \pi$



Equation of PQ:

$$\begin{aligned}\frac{y - 2\sin\theta}{x - 3\cos\theta} &= \frac{2\tan\theta - 2\sin\theta}{3\sec\theta - 3\cos\theta} \\ &= \frac{2\sin\theta(\sec\theta - 1)}{3\cos\theta(\sec^2\theta - 1)} \\ &= \frac{2\sin\theta}{3(1 + \cos\theta)}\end{aligned}$$

$$\text{Simplify: } \underline{2x\sin\theta - 3y(1 + \cos\theta) - 6\sin\theta = 0}$$

• Show that PQ passes through common point (3,0)

Substitute (3,0) into equation of PQ

$$6\sin\theta - 3 \times 0(1 + \cos\theta) - 6\sin\theta = 0$$

$$6\sin\theta - 6\sin\theta = 0 \quad (\text{True})$$

Therefore PQ passes through (3,0)

(ii) Show that the tangent at P to the first curve meets the tangent at Q to the second curve in a point which lies on the common tangent to the two curves at their other common point. 2

ii) Equation of tangent to Ellipse at P

$$\frac{x \cdot \cos\theta}{3} + \frac{y \sin\theta}{2} = 1 \quad (1)$$

Equation of tangent to Hyperbola at Q

$$\frac{x \sec\theta}{3} - \frac{y \tan\theta}{2} = 1 \quad (2)$$

• Divide equation (1) by $\cos\theta$.

$$\frac{x}{3} + \frac{y \tan\theta}{2} = \sec\theta \quad (3)$$

Note: the sign of the second term $\frac{y \sin\theta}{2}$ has to be changed to $-$ because θ is in 2nd quadrant, $\sin\theta$ and $\tan\theta$ are opposite sign

• point of intersection T: (3) - (2) gives

$$\frac{x}{3} - \frac{x \sec\theta}{3} = \sec\theta - 1$$

$$\frac{x}{3} (1 - \sec\theta) = \sec\theta - 1$$

$$\therefore \boxed{x = -3}$$

Therefore, T lies on the line $x = -3$, which is the common tangent of Ellipse and Hyperbola at the common point $(-3, 0)$

(iii) Prove that the two curves have the same length of the latus rectum. 2

iii) Length of Latus rectum:

• of Ellipse: $\frac{x^2}{9} + \frac{y^2}{4} = 1$

$e = \frac{\sqrt{5}}{3}$, $S(\sqrt{5}, 0)$.

Equation of Latus rectum $x = \sqrt{5}$

Intersection points: $\frac{5}{9} + \frac{y^2}{4} = 1$, $y^2 = \frac{16}{9}$, $y = \pm \frac{4}{3}$

Therefore, Length of Latus rectum of Ellipse

$$L = \frac{8}{3}$$

• of Hyperbola: $\frac{x^2}{9} - \frac{y^2}{4} = 1$

$e = \frac{\sqrt{13}}{3}$, $S(\sqrt{13}, 0)$, $x = \sqrt{13}$

Intersection points $\frac{13}{9} - \frac{y^2}{4} = 1$, $y^2 = \frac{16}{9}$, $y = \pm \frac{4}{3}$

Length of latus rectum $L = 2y = \frac{8}{3}$

Therefore, the 2 curves E & H have the same length of Latus Rectum.

(B) Show that the condition for a straight line $y = mx + c$ to touch the ellipse E of 3

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \quad \text{is} \quad c^2 = b^2 + a^2 m^2$$

Hence show that the locus of the point P(x, y) from which the 2 tangents to the ellipse E

$\frac{x^2}{16} + \frac{y^2}{9} = 1$ are perpendicular together, is a curve with the centre at the origin and a radius of 5.

Condition to be a tangent to an ellipse.

Point of intersection.

$$\frac{x^2}{a^2} + \frac{(mx + c)^2}{b^2} = 1$$

$$b^2x^2 + a^2m^2x^2 + 2a^2mcx + a^2c^2 - a^2b^2 = 0$$
$$(b^2 + a^2m^2)x^2 + (2a^2mc)x + (a^2c^2 - a^2b^2) = 0$$

$$\text{Discriminant : } \Delta = (2a^2mc)^2 - 4(b^2 + a^2m^2)(a^2c^2 - a^2b^2)$$
$$= 4a^4m^2c^2 - 4a^2b^2c^2 + 4a^2b^4 - 4a^4m^2c^2$$
$$+ 4a^4b^2m^2$$

To be a tangent, the line has only one common point with the ellipse or $\Delta = 0$

$$\Delta = 4a^2b^2(b^2 + a^2m^2 - c^2) = 0$$

$$\therefore \boxed{c^2 = b^2 + a^2m^2}$$

• Locus of P(x, y)

Let equation of tangent through P(x, y)
 $y = mx + c$

Using condition above: $c^2 = b^2 + a^2m^2$
 $c^2 = 9 + 16m^2$

$$\therefore y = mx + \sqrt{9 + 16m^2}$$

Solving equation in terms of m

$$(y - mx)^2 = 9 + 16m^2$$

$$y^2 - 2xym + m^2x^2 = 9 + 16m^2$$

Quadratic equation:

$$m^2(x^2 - 16) - 2xym + (y^2 - 9) = 0$$

Since from external point P(x, y), there are 2 tangents with gradient m_1 and m_2 to the ellipse

The tangents are perpendicular, then $m_1 \times m_2 = -1$

• m_1 and m_2 are 2 roots of the above quadratic equation, their product ($m_1 \times m_2$) is equal $\frac{c}{a}$, which is

$$m_1 \times m_2 = \frac{y^2 - 9}{x^2 - 16} = -1$$

Therefore $x^2 + y^2 = 25$

So the locus of P is a circle with radius = 5

Question 4

12

(A) A function is defined with the polar equation as follows:

$$\begin{cases} x = 8 \cos^3 \theta \\ y = 8 \sin^3 \theta \end{cases} \quad -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$$

(i) Find $\frac{dy}{dx}$ in term of θ and show that the graph of this function touches the x and y axis. Sketch the curve.

3

Find $\frac{dy}{dx}$, using chain rule

$$\frac{dy}{dx} = \frac{dy}{d\theta} \times \frac{d\theta}{dx} = \frac{24 \cos \theta \times \sin^2 \theta}{-24 \sin \theta \times \cos^2 \theta}$$

$$\therefore \frac{dy}{dx} = -\frac{\sin \theta}{\cos \theta} = -\tan \theta$$

• When $\theta = 0$, $\frac{dy}{d\theta} = 0$, $x = 8$, $y = 0$

\therefore The tangent at the x intercept $(8, 0)$ is horizontal, OR the curve touches x axis

• When $\theta = \pm \frac{\pi}{2}$, $\frac{dy}{d\theta} = \infty$, $x = 0$, $y = \pm 8$

\therefore The tangents at the y intercepts $(0, \pm 8)$ is vertical, OR the curve touches y axis at 2 points.

(ii) Show that the Cartesian equation of that function is

3

$$x^{\frac{2}{3}} + y^{\frac{2}{3}} = 4$$

Show that the equation of the tangent to the curve at the point $P(x_0, y_0)$ is:

$$y_0^{1/3} x + x_0^{1/3} y = 4x_0^{1/3} \cdot y_0^{1/3}$$

Prove that the segment intercepted on this tangent by the coordinate axis is independent of the position of P on the curve.

change to cartesian form

$$\cos \theta = \frac{x^{1/3}}{2}$$

$$\sin \theta = \frac{y^{1/3}}{2}$$

Then :

$$\cos^2 \theta + \sin^2 \theta = 1$$

$$\frac{x^{2/3}}{4} + \frac{y^{2/3}}{4} = 1$$

OR $\boxed{x^{2/3} + y^{2/3} = 4}$

Equation of tangent at $P(x_0, y_0)$

$$\text{Since } \frac{dy}{dx} = -\frac{\sin \theta}{\cos \theta} = -\frac{y^{1/3}}{x^{1/3}}$$

$$\text{gradient of tangent } m_T = -\frac{y_0^{1/3}}{x_0^{1/3}}$$

Equation of tangent

$$y - y_0 = -\frac{y_0^{1/3}}{x_0^{1/3}} (x - x_0)$$

$$x_0^{1/3} \cdot y - y_0 \cdot x_0^{1/3} = -y_0^{1/3} \cdot x + x_0 \cdot y_0^{1/3}$$

$$\therefore y_0^{1/3} \cdot x + x_0^{1/3} \cdot y = x_0 y_0^{1/3} + y_0 x_0^{1/3}$$

$$= x_0^{1/3} \cdot y_0^{1/3} (x_0^{2/3} + y_0^{2/3})$$

$$= 4 \cdot x_0^{1/3} \cdot y_0^{1/3}$$

Equation of tangent

$$\boxed{y_0^{1/3} \cdot x + x_0^{1/3} \cdot y = 4 x_0^{1/3} \cdot y_0^{1/3}}$$

X and Y intercepts of the tangent.

$$\text{X intercept, } y=0, x = 4x_0^{1/3}$$

$$\text{Y intercept, } x=0, y = 4y_0^{1/3}$$

$$\begin{aligned}\text{Length of XY segment} &= \sqrt{(4x_0^{1/3})^2 + (4y_0^{1/3})^2} \\ &= \sqrt{16(x_0^{2/3} + y_0^{2/3})} \\ &= \sqrt{16 \times 4} = 8\end{aligned}$$

\therefore This length is independent of (x_0, y_0)

(B) Consider the function $f(x) = 2 - \frac{4x}{x^2 + 1}$

(i) Show that the function is always positive for any value of x. 1

show that $f(x)$ is positive definite

$$f'(x) = \frac{2x^2 + 2 - 4x}{x^2 + 1} = \frac{2(x^2 - 2x + 1)}{x^2 + 1}$$

$$f(x) = \frac{2(x-1)^2}{x^2 + 1} \text{ always greater or equal zero.}$$

(ii) Find the asymptote (if any) and the stationary point of that curve. 1

Asymptote : Let $x \rightarrow \infty$

$$\text{Limit } f(x) = 2 - \frac{4}{\infty} = 2$$

Horizontal asymptote $\boxed{y = 2}$

$$\text{Stationary point : } f'(x) = \frac{4x^2 - 4}{(x^2 + 1)^2}$$

$$\text{when } x = \pm 1, f'(x) = 0$$


$$y = 0 \text{ OR } 4$$

Maximum point $(-1, 4)$

Minimum point $(1, 0)$

(iii) Sketch the curve $y = f(x)$.


1



(iv) On a separate diagram, sketch the relating curves:

a) $y = f(|x|)$


1



1


1

b) $y = \frac{1}{f(x)}$



c) $y = \ln f(x)$


$y = \ln f(x)$



Question 5


12

(A) The area bounded by the curve $y = \sin x$, the two lines $y = -x$ and $x = \pi$ is rotated about the line $y = -x$. Find the volume of the solid shape of that formation. 6



The solid shape can be divided by 2 separated volumes:

i) The 1st part produced by rotating area bounded by the curve $y = \sin x$, the perpendicular line about the line $y = -x$ as shown in the following figure



By using the slicing method: The slice is a piece of cylinder, with volume is

$$dV = \pi R^2 dh$$


with R is the perpendicular distance PQ to the

Line $y = -x$ or $x + y = 0$

$$R = PQ = \frac{|x+y|}{\sqrt{2}}$$

Calculate dh by the figure:

$$dh = \sqrt{2} \cdot dx$$



$$\begin{aligned} \therefore dV &= (x+y)^2 \cdot \sqrt{2} \cdot dx \\ &= (x + \sin x)^2 \cdot \sqrt{2} dx \\ &= \sqrt{2} x^2 + 2\sqrt{2} x \sin x + \sqrt{2} \sin^2 x dx \end{aligned}$$

Therefore:

$$V = \lim_{dx \rightarrow 0} \sum dV = \int_0^{\pi} \sqrt{2} x^2 + 2\sqrt{2} x \sin x + \sqrt{2} \sin^2 x dx$$

There are 3 separated integrals.


$$\bullet \int_0^{\pi} \sqrt{2} x^2 dx = \frac{\sqrt{2}}{3} \pi \sqrt{\pi}$$

$$\begin{aligned} \bullet \int_0^{\pi} 2\sqrt{2} x \cdot \sin x dx &= \left[-x \cos x + \sin x \right]_0^{\pi} \\ &= 2\sqrt{2} \pi \end{aligned}$$

$$\bullet \int_0^{\pi} \sqrt{2} \sin^2 x dx = \frac{\sqrt{2}}{2} \left[x - \frac{1}{2} \sin 2x \right]_0^{\pi} = \frac{\sqrt{2} \pi}{2}$$

$$\text{Therefore } V = \frac{\sqrt{2}}{3} \pi \sqrt{\pi} + 2\sqrt{2} \pi + \frac{\sqrt{2}}{2} \pi = 11.34 u^3$$

ii) The 2nd part of that volume is the right-angled cone with the radius and the height equal to $\frac{\pi}{\sqrt{2}}$




The volume of a cone: $V = \frac{1}{3} \pi R^2 h$

$$= \frac{1}{3} \pi \left(\frac{\pi}{\sqrt{2}}\right)^2 \left(\frac{\pi}{\sqrt{2}}\right)$$

$$= \frac{\pi^4}{6\sqrt{2}} = 11.49 \text{ u}^3$$

Total volume of the solid shape = 22.8 unit cube


(B) A cylindrical timber is chopped at two ends by 2 planes which are inclined 60° and 45° respectively. If the two ends are tilt toward each other and the shortest length of 2 ends is 120cm. Find the volume of that timber. Give the radius of timer is 10cm. 6



Volume of a chopped cylindrical timber:

The total volume can be divided by 3 parts, The 2 ends are the pieces of timber which are chopped into half, and the body is the full cylinder.

• One end:



$$V = \frac{1}{2} \pi 10^2 \times 20 = 1000 \pi$$

• Other end: is similar, except the length of that piece is $\frac{20}{\sqrt{3}}$ cm

$$V = \frac{1}{2} \pi 10^2 \times \frac{20}{\sqrt{3}} = \frac{1000 \pi}{\sqrt{3}}$$

- The body is the full cylinder

$$V = \pi \times 10^2 \times 120 = 12000\pi$$

Therefore total volume


$$V = 1000\pi + \frac{1000\pi}{\sqrt{3}} + 12000\pi = \underline{42654 \text{ cm}^3}$$

$$\boxed{V = 42.65 \text{ Litre}}$$

Question 6

12

(A)



Two circles intersect at A and D. P is the point on the major arc of one circle. The other circle has the radius r. PA produced and PD produced meet the other circle at B and C respectively. Let $\angle APD = \alpha$ and $\angle ACD = \beta$.

- (i) Show that $BC = 2r \sin(\alpha + \beta)$.

2

From centre O, draw $OM \perp$ bisects BC

$$\angle BOC = 2\angle BAC \text{ (angle at the centre)}$$

$$\angle BAC = \alpha + \beta \text{ (ext. } \angle \text{ of } \triangle ACP)$$

$$\therefore \angle BOC = 2(\alpha + \beta)$$

$$\therefore \angle BOM = \alpha + \beta \text{ (in isosceles } \triangle BOC)$$

$$\text{In Right angle } \triangle OBM, \sin(\alpha + \beta) = \frac{BM}{OB}$$

$$\therefore BM = r \cdot \sin(\alpha + \beta)$$

$$\text{Then } BC = 2BM = 2r \sin(\alpha + \beta)$$

- (ii) As P moves along the major arc AD on its circle, show that the length of the chord BC is independent of the position of P. 2

Prove BC is independent of the position of P.

Since AD is common chord of both circles, AD is constant, therefore angle α and β

subtend AD on both circles will be constant, no matter of P moves along the arc. Therefore length of BC will not change.

- (iii) If the 2 circles have equal radii, show that 2

$$BC = 2 \cos \alpha \cdot AD$$

If the 2 circles are the same, then

$$\alpha = \beta \text{ (angles subtend equal arcs)}$$

$$\text{Hence } BC = 2r \sin 2\alpha = 2r \cdot 2 \sin \alpha \cdot \cos \alpha$$

Draw ON \perp bisects AD,

$$\angle AOD = 2\angle ACD = 2\beta$$

$$\angle AON = \beta$$

$$\text{Hence in } \triangle AON, \sin \beta = \frac{AN}{OA}$$

$$AN = r \cdot \sin \beta$$

$$AD = 2r \sin \beta$$

$$\text{Since } \alpha = \beta, \text{ then } AD = 2r \sin \alpha.$$

Substitute into BC

$$BC = 2 AD \cos \alpha$$

- (B) P is any point (ct, c/t) on the Hyperbola $xy = c^2$, whose centre is O.

- (i) M and N are perpendicular roots of P to the 2 asymptotes. Prove that PM.PN is constant. 1

- (ii) Find the equation of tangent at P, and show that OP and this tangent are equally inclined to the asymptotes. 2

(iii) If the tangent at P meet the asymptotes at A and B, and Q is the fourth vertex of the rectangle OAQB, find the locus of Q. 1

(iv) Show that $PA=PB$ and hence conclude that the area of ΔOAB is independent of position of P. 2

Question 7

12

(A) If the polynomial $P(x) = x^4 - 4x^3 + 11x^2 - 14x + 10$ has two zeros $(a + ib)$ and $(a - 2ib)$ where a and b are real, then find the values of a and b .

4

Hence find the zeros of $P(x)$ over the complex field \mathbb{C} , and express $P(x)$ as the product of 2 quadratic factors with rational coefficients.

$P(x) = x^4 - 4x^3 + 11x^2 - 14x + 10$ with 2 zeros $(a + ib)$ and $(a - 2ib)$, then find a, b .

Apply the conjugate zeros, $P(x)$ has other 2 zeros which are conjugate to the above. They are $a - ib$ and $a + 2ib$. Let $P(x) = 0$

$$x^4 - 4x^3 + 11x^2 - 14x + 10 = 0$$

The sum of 4 roots = $-\frac{b}{a}$

$$a + ib + a - ib + a + 2ib + a - 2ib = 4$$

$$4a = 4$$

$$a = 1$$

The products of 4 roots = $\frac{c}{a}$

$$(1 + ib)(1 - ib)(1 + 2ib)(1 - 2ib) = 10$$

$$(1 + b^2)(1 + 4b^2) = 10$$

$$4b^4 + 5b^2 - 9 = 0$$

$$(4b^2 + 9)(b^2 - 1) = 0$$

$$b^2 = 1$$

$$b = 1$$

- Therefore, 4 zeros of $P(x)$ are $1 + i, 1 - i, 1 + 2i, 1 - 2i$

- Factorise $P(x)$ in real set.

$$\begin{aligned} P(x) &= (x - 1 + i)(x - 1 - i)(x - 1 + 2i)(x - 1 - 2i) \\ &= ((x - 1)^2 + 1)((x - 1)^2 + 4) \end{aligned}$$

$$P(x) = (x^2 - 2x + 2)(x^2 - 2x + 5)$$

(B) Show that if the polynomial $P(x) = 0$ has a root α of multiplicity m , then

4

$P'(x) = 0$ has a root α of multiplicity $(m - 1)$.

Given that $P(x) = x^4 + x^3 - 3x^2 - 5x - 2 = 0$ has a 3-fold root, find all the roots of $P(x)$.

b) If $P(x) = 0$ has $x = \alpha$ as a multiple root, then

$$P(x) = (x - \alpha)^m \cdot Q(x) \quad \text{with } m \text{ is the multiplicity}$$

$$P'(x) = m(x - \alpha)^{m-1} \cdot Q(x) + (x - \alpha)^m \cdot Q'(x)$$

$$P'(x) = (x - \alpha)^{m-1} [m \cdot Q(x) + (x - \alpha) \cdot Q'(x)]$$

Let $x = \alpha$, $P'(\alpha) = 0$, so α is also the root of $P'(x)$.

$$\text{Let } P(x) = x^4 + x^3 - 3x^2 - 5x - 2 = 0$$

$$P'(x) = 4x^3 + 3x^2 - 6x - 5$$

$$P''(x) = 12x^2 + 6x - 6 = 0$$

$$6(2x - 1)(x + 1) = 0$$

$$\therefore x = -1 \text{ or } \frac{1}{2}$$

Substitute $x = -1$ into $P(x)$ and $P'(x)$ we get

$$P(-1) = P'(-1) = P''(-1) = 0$$

$\therefore x = -1$ is a multiple root multiplicity = 3

Let the 4th root be α , using the sum of 4

$$\text{roots} = -\frac{b}{a}$$

$$-1 \times 3 + \alpha = -1$$

$$\alpha = 2$$

All the roots are, $-1, -1, -1$ and 2

(C) Find the cubic roots of unity and express them in the form $r(\cos \theta + i \sin \theta)$.

Show these roots on an Argand diagram.

If w is one of the complex roots, prove that the other root is w^2 and show that

$$1 + w + w^2 = 0.$$

(i) Prove that if n is a positive integer, then $1 + w^n + w^{2n} = 3$ or 0 depending on whether n is or is not a multiple of 3. 2

c) Find the cube roots of unity $z^3 = 1$.

$$|z^3| = |z|^3 = 1 \quad \therefore |z| = 1$$

$$\text{Let } z = \cos \theta + i \sin \theta$$

$$z^3 = \cos 3\theta + i \sin 3\theta = 1$$

$$\cos 3\theta = 1$$

$$3\theta = 0, 2\pi, 4\pi$$

$$\theta = 0, \frac{2\pi}{3}, \frac{4\pi}{3}$$

The 3 roots are: $z = \cos 0 + i \sin 0 = 1$

$$z = w = \cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3}$$

$$z = w^2 = \cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3}$$

Since $z^3 - 1 = 0$

$$\text{Sum of 3 roots} = -\frac{b}{a} = 0$$

$$\text{Then } 1 + w + w^2 = 0$$

• Show that $\boxed{1 + w^n + w^{2n} = 3 \text{ or } 0}$

• i) if n is a multiple of 3, let

$$n = 3p.$$

$$\begin{aligned} \text{then } 1 + w^n + w^{2n} &= 1 + (w^3)^p + (w^3)^{2p} \\ &= \frac{1 + 1 + 1 = 3}{(\text{since } w^3 = 1)} \end{aligned}$$

• ii) if n is NOT a multiple of 3, let

$$n = 3p + 1 \text{ or } 3p + 2$$

$$\begin{aligned} \text{Then } 1 + w^n + w^{2n} &= 1 + w^{3p+1} + w^{6p+2} \\ &= \underline{1 + w^3 \cdot w + w^{6p} \cdot w^2} \\ &= 1 + w + w^2 \\ &= 0 \end{aligned}$$

(ii) If $x = a + b$, $y = aw + bw^2$ and $z = aw^2 + bw$, show that 2

$$z^2 + y^2 + x^2 = 6ab$$

If $x = a + bw$, $y = aw + bw^2$ and $z = aw^2 + bw^3$.

Show that $z^2 + y^2 + x^2 = 6ab$

$$\begin{aligned} z^2 &= (aw^2 + bw^3)^2 = a^2w^4 + 2abw^5 + b^2w^6 \\ &= \underline{a^2w + b^2w^2 + 2ab} \quad (1) \end{aligned}$$

$$\begin{aligned} y^2 &= (aw + bw^2)^2 = a^2w^2 + 2abw^3 + b^2w^4 \\ &= \underline{a^2w^2 + b^2w + 2ab} \quad (2) \end{aligned}$$

$$x^2 = (a + b)^2 = \underline{a^2 + b^2 + 2ab} \quad (3)$$

$$\begin{aligned} \therefore z^2 + y^2 + x^2 &= a^2(1+w+w^2) + b^2(1+w+w^2) + 6ab \\ &= 6ab \end{aligned}$$

Question 8 - Optional

12


(A) A particle is projected vertically upward with initial speed u . The air resistance is proportional to the speed of the particle.

(a) If $\ddot{x} = -(g + kv)$ with k is the constant, then find the maximum height reached by the particle and the time to do so.

2

Vertical projectile motion:

a) Project upward:



Equation of motion

$$a = \frac{v dv}{dx} = -g - kv$$

$$\int \frac{v dv}{g + kv} = \int -dx$$

$$\frac{1}{k} \int \frac{kv + g}{kv + g} dv - \frac{g}{k} \int \frac{dv}{kv + g} = -x + c$$

$$\therefore x + c = \frac{g}{k^2} \ln(kv + g) - \frac{v}{k}$$

When $x = 0$, $v = u$,

$$c = \frac{g}{k^2} \ln(ku + g) - \frac{u}{k}$$

Maximum height, when $v=0$, $x=H$


$$H = \frac{u}{k} + \frac{g}{k^2} \ln g - \frac{g}{k^2} \ln(ku+g)$$

$$H = \frac{u}{k} - \frac{g}{k^2} \ln\left(\frac{ku+g}{g}\right)$$

(b) Set up the differential equation for the downward motion.

2

b) Downward motion



Equation of motion

$$a = g - kv$$

$$\int \frac{v dv}{g - kv} = \int dx$$

$$-\frac{1}{k} \int \frac{-kv + g}{g - kv} dv + \int \frac{g dv}{g - kv} = x + c$$

$$\therefore x + c = -\frac{v}{k} - \frac{g}{k^2} \ln(g - kv)$$

when $x=0$, $v=0$

$$c = -\frac{g}{k^2} \ln g$$

when $x=H$, $v=V$

$$\frac{u}{k} + \frac{g}{k^2} \ln g - \frac{g}{k^2} \ln(ku+g) - \frac{g}{k^2} \ln g = -\frac{V}{k} - \frac{g}{k^2} \ln(g - kV)$$

Simplify both sides by k^2 :

$$ku + kV = g \ln(ku+g) - g \ln(g - kV)$$

$$\therefore k(u+V) = g \ln\left[\frac{ku+g}{g - kV}\right]$$

(c) Show that the particle returns to its point of projection with speed v given by 2

$$k(u+v) = g \log_e \left[\frac{g+ku}{-g-kv} \right]$$

(B) Show that for $n \geq 1$ 3

$$1 \cdot \ln \frac{2}{1} + 2 \ln \frac{3}{2} + 3 \ln \frac{4}{3} + \dots + n \ln \left(\frac{n+1}{n} \right) = \ln \left(\frac{(n+1)^n}{n!} \right)$$

show that for $n \geq 1$

$$1 \ln \frac{2}{1} + 2 \ln \frac{3}{2} + 3 \ln \frac{4}{3} + \dots + n \ln \left(\frac{n+1}{n} \right) = \ln \frac{(n+1)^n}{n!}$$

$$\text{LHS} = \ln \left(\frac{2}{1} \right)^1 + \ln \left(\frac{3}{2} \right)^2 + \ln \left(\frac{4}{3} \right)^3 + \dots + \ln \left(\frac{n+1}{n} \right)^n$$

$$= \ln \left[\frac{2^1 \cdot 3^2 \cdot 4^3 \cdot 5^4 \cdot \dots \cdot n^{n-1} \cdot (n+1)^n}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot \dots \cdot (n-1)^{n-1} \cdot n^n} \right]$$

simplify the indices of the top and bottom :

$$\text{LHS} = \ln \left[\frac{(n+1)^n}{1 \cdot 2 \cdot 3 \cdot \dots \cdot n} \right] = \ln \left(\frac{(n+1)^n}{n!} \right) = \text{RHS}$$

(C) By using the induction method, prove that 3

$(35)^n + 3 \times 7^n + 3 \times 5^n + 6$ is divisible by 12 for $n \geq 1$

STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1}x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a}e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left(x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2} \right)$$

NOTE : $\ln x = \log_e x, \quad x > 0$