

# Year 12 Mathematics Trial HSC Examination 2014

## **General Instructions**

- Reading time 5 minutes
- Working time 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided on the back page of this question paper
- In Questions 11 16, show relevant mathematical reasoning and/or calculations

## Total marks – 100

Section I

#### 10 marks

- Attempt Questions 1 10
- Allow about 15 minutes for this section.



#### 90 marks

- Attempt Questions 11-16
- Start each question in a new writing booklet
- Write your name on the front cover of each booklet to be handed in
- If you do not attempt a question, submit a blank booklet marked with your name and "N/A" on the front cover
- Allow about 2 hours 45 minutes for this section

# DO NOT REMOVE THIS PAPER FROM THE EXAMINATION ROOM

**BLANK PAGE** 

#### Section I

#### 10 marks Attempt Questions 1–10 Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1–10.

- 1 What is the primitive of  $\frac{2}{x} \cos x$ ?
  - (A)  $-\frac{2}{x^2} + \sin x + C$

$$(B) \quad -\frac{2}{x^2} - \sin x + C$$

- (C)  $2\ln x + \sin x + C$
- (D)  $2\ln x \sin x + C$
- 2 What are the values of x for which |4-3x| < 13?
  - (A) x < -3 and  $x < \frac{17}{3}$
  - (B) x > -3 and  $x > \frac{17}{3}$
  - (C) x > -3 and  $x < \frac{17}{3}$
  - (D)  $x < -3 \text{ and } x > \frac{17}{3}$

3

- What is the simultaneous solution to the equations 2x + y = 7 and x 2y = 1?
- (A) x = 3 and y = 1
- (B) x = -1 and y = 9
- (C) x = 2 and y = 3
- (D) x = 5 and y = 1

**4** Factorise  $2x^2 - 7x - 15$ 

- (A) (2x-3)(x-5)
- (B) (2x+3)(x-5)
- (C) (2x-5)(x-3)
- (D) (2x+5)(x-3)

5 The value of  $\frac{5.79 + 0.55}{\sqrt{4.32 - 3.28}}$  is closest to:

- (A) 4
- (B) 6
- (C) 9
- (D) 10

6 What are the values of p and q given  $(3\sqrt{12} + \sqrt{75})(2 + \sqrt{48}) = p + q\sqrt{3}$ ?

- (A) p = 132 and q = 15
- (B) p = 396 and q = 15
- (C) p = 132 and q = 22
- (D) p = 396 and q = 22
- 7 The line 6x ky = 8 passes through the point (3,2). What is the value of k?
  - (A) -13
  - (B) -5
  - (C) 5
  - (D) 15

8 The semi-circle  $y = \sqrt{4-x^2}$  is rotated about the *x*-axis. Which of the following expressions is correct for the volume of the solid of revolution?

(A) 
$$V = \pi \int_{0}^{2} (4 - x^{2}) dx$$
  
(B)  $V = 2\pi \int_{0}^{2} (4 - x^{2}) dx$   
(C)  $V = \pi \int_{0}^{2} (4 - y^{2}) dy$ 

(D) 
$$V = 2\pi \int_{0}^{2} (4 - y^{2}) dy$$

**9** A circle has the equation  $4x^2 - 4x + 4y^2 + 24y + 21 = 0$ . What is the radius and centre?

(A) Centre 
$$\left(\frac{1}{2}, -3\right)$$
 and radius of 2.

(B) Centre 
$$\left(\frac{1}{2},3\right)$$
 and radius of 2.

(C) Centre 
$$\left(\frac{1}{2}, -3\right)$$
 and radius of 4.

(D) Centre 
$$\left(\frac{1}{2},3\right)$$
 and radius of 4.

**10** An infinite geometric series has a first term of 12 and a limiting sum of 15. What is the common ratio?

(A) 
$$\frac{1}{5}$$
  
(B)  $\frac{1}{4}$   
(C)  $\frac{1}{3}$   
(D)  $\frac{1}{2}$ 

## Section II

#### 90 marks Attempt Questions 11–16 Allow about 2 hours 45 minutes for this section

Answer each question in the appropriate writing booklet. Extra writing booklets are available.


In Questions 11–16, your responses should include relevant mathematical reasoning and/or calculations.

Marks

1

#### Question 11 (15 marks)

(a) The diagram shows the points A(-1,1), B(3,6), C(5,1) and D(-3,-9).



- (iii) Find the perpendicular distance from *A* to the line *BE*. 2
- (iv) Show that *ABED* is a parallelogram. 2
- (v) Find the area of *ABED*.

#### Question 11 continues over the page

- (b) Find the equation of the tangent to the curve y = log<sub>e</sub> x-1 at the point (e,0).
  (c) The equation of a parabola is given by y = x<sup>2</sup> 2x + 5.
  (i) Find the coordinates of its vertex.
  - (ii) State the focal length of the parabola. 1
  - (iii) Find the equation of the normal at the point P(2,5). 2

#### Question 12 (15 marks) Use a SEPARATE writing booklet

#### Marks

1

(a) There are 200 tickets sold in a raffle with only two prizes. These tickets are placed in a bag and two are drawn, one at a time. Once a ticket is drawn it is not placed back in the bag. One boy bought 3 tickets. calculate the probability the boy wins:

(i)	First prize.	1
(ii)	Both prizes.	1
(iii)	The second prize only.	1

- (iv) No prize at all.
- (b) Differentiate with respect to *x*.

(i) 
$$e^{3x} \tan x$$
 2

(ii) 
$$\frac{\sin x}{5-x}$$
 2

(c) Evaluate

(i) 
$$\int \frac{1}{1-2x} dx$$
 2

(ii) 
$$\int_{0}^{\pi} \sec^2 \frac{x}{3} dx$$
 2

(d) The roots of the equation  $2x^2 - x - 15 = 0$  are  $\alpha$  and  $\beta$ . Find the value of:

- (i)  $\alpha + \beta$  1
- (ii)  $\alpha\beta$  1

(iii) 
$$\frac{1}{\alpha} + \frac{1}{\beta}$$
 1


## Question 13 (15 marks) Use a SEPARATE writing booklet

Find the perimeter of the *n*th hexagon.

(i)

(b)

(a) Melanie is using wire to construct a geometrical design which consists of *n* regular hexagons with sides 2cm, 5cm, 8cm and so on going up by the same amount each hexagon. The diagram below shows the first 3 hexagons of her design.



Show that the total length of the wire is  $L = 9n^2 + 3n$ . 2 (ii) (iii) If the total length of the wire is 6 metres, find the number of hexagons that Melanie has constructed. 2 Let  $f(x) = x^3 - 3x^2 - 9x + 22$ (i) Show that f''(x) = 6x - 61 (ii) Find the coordinates of the stationary points on y = f(x) and determine their nature. 2 2 (iii) Find the coordinates of the point(s) of inflexion. Sketch the graph of y = f(x), indicating where the curve meets the (iv) y-axis, stationary points and the point(s) of inflexion. 2

(v) For what values of x is the graph of y = f(x) concave down?


#### Marks

1

1

-9-


(c) Alex and Bella leave from point *O* at the same time. Alex travels at 20km/h along a straight road in the direction 085°. Bella travels at 25km/h along another straight road in the direction 340°.



Find the distance Alex and Bella are apart to the nearest kilometre after two hours.

2


(a)




*ABCD* is a rhombus, *BE* is perpendicular to *AD* and intersects *AC* at *F*. Copy or trace the diagram into your writing booklet.

Question 14 continues over the page				
	(iii)	The number of bacteria doubles every $x$ days. Find $x$ . Answer correct to 1 decimal place.	2	
	(ii)	When $t = 3$ the number of bacteria was estimated at $1.5 \times 10^8$ . Evaluate A. Answer correct to 2 significant figures.	1	
	(i)	Show that the number of bacteria increases at a rate proportional to the number present.	2	
	Where t is measured in days and A is a constant.			
(b)	A scientist grows the number of bacteria according to the equation $N(t) = Ae^{0.15t}$			
	(iv)	Hence, or otherwise, find the size of $\angle FDC$ .	1	
	(iii)	Show that $\angle FBC$ is a right angle.	1	
	(ii)	Prove that the triangles <i>BFC</i> and <i>DFC</i> are congruent.	3	
	(i)	Explain why $\angle BCA = \angle DCA$ .	1	

(c) During a survey the area of an irregular headland was to be found. Measurements of the area were noted on the diagram below.






(d) Solve the equation  $(\cos x + 2)(2\cos x + 1) = 0$  in the domain  $0 \le x \le 2\pi$ .

2



2



The diagram shows the graphs of  $y = e^x - 2$  and  $y = e^{-x}$ .

(i)	Find the area between the curves from $x = 1$ to $x = 2$ . Leave your answer in terms of $e$ .	3
(ii)	Show that the curves intersect when $e^{2x} - 2e^x - 1 = 0$ .	2

(iii) Hence, using the substitution  $u = e^x$ , or otherwise, find the point of intersection of the curves.

(b) Velocity of an object moving along the *x*-axis is given by  $v = 2\sin t + 1$  for  $0 \le t \le 2\pi$ 


Where v is measured in metres per second and t in seconds.

(i)	When is the object at rest?	2
(ii)	Sketch the graph of v as a function of t for $0 \le t \le 2\pi$	2
(iii)	Find the maximum velocity of the object for this period.	1
(iv)	When is the object travelling in the negative direction during this period?	1
(v)	Calculate the total distance travelled by the object in the period $0 \le t \le \pi$	2

# Question 16 (15 marks) Use a SEPARATE writing booklet

) George is saving for a holiday. He opens a savings account with an interest rate of 0.4% per month compounded monthly at the end of each month. George decides to deposit \$450 into the account on the first day of each month. He makes his first deposit on the 1 <sup>st</sup> of January 2012 and his last on the 1 <sup>st</sup> of December 2014. George withdraws the entire amount, plus interest, immediately after his final interest payment on the 31 <sup>st</sup> December 2014.		
(i)	How much in total did George deposit into his savings account over this period?	1
(ii)	How much did George withdraw from his account on the 31 <sup>st</sup> December 2014? Answer correct to the nearest dollar.	3
(iii)	George's holiday is postponed due to family illness. He decides to deposit \$12 000 into a different account with an interest rate of 5% p.a. compounded quarterly for 2 years. How much will George receive at the end of the investment period from this \$12000 investment? Answer correct to the nearest dollar.	2
	inter mon of ea his la amou 31 <sup>st</sup> [ (i) (ii)	<ul> <li>interest rate of 0.4% per month compounded monthly at the end of each month. George decides to deposit \$450 into the account on the first day of each month. He makes his first deposit on the 1<sup>st</sup> of January 2012 and his last on the 1<sup>st</sup> of December 2014. George withdraws the entire amount, plus interest, immediately after his final interest payment on the 31<sup>st</sup> December 2014.</li> <li>(i) How much in total did George deposit into his savings account over this period?</li> <li>(ii) How much did George withdraw from his account on the 31<sup>st</sup> December 2014? Answer correct to the nearest dollar.</li> <li>(iii) George's holiday is postponed due to family illness. He decides to deposit \$12 000 into a different account with an interest rate of 5% p.a. compounded quarterly for 2 years. How much will George receive at the end of the investment period from this \$12000</li> </ul>

# Question 16 continues over the page



(b)

*ABCD* is a rectangle with CD = 3 cm and AD = 2 cm. *F* and *E* lie on the lines *BC* and *BA*, so that, *F*, *D* and *E* are collinear. Let CF = x cm and AE = y cm.

(i)	Show that $\Delta FCD$ and $\Delta DAE$ are similar.	3
(ii)	Show that $xy = 6$	1
(iii)	Show that the area (A) of $\Delta FBE$ is given by $A = 6 + \frac{3}{2}x + \frac{6}{x}$ .	2
(iv)	Find the height and base of $\Delta FBE$ with minimum area. Justify your	

3

#### **End of Examination**

answer.

# **STANDARD INTEGRALS**

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \ x \neq 0, \ \text{if } n < 0$$

$$\int \frac{1}{x} dx = \ln x , \qquad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \qquad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \qquad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \qquad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \qquad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \qquad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \qquad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \qquad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - a^2}} dx = \sin^{-1} \frac{x}{a}, \qquad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left(x + \sqrt{x^2 + a^2}\right)$$

*Note*  $\ln x = \log_e x$ , x > 0

2014 Year 12 Trial HSC Examination

Student Name: .....

## Mathematics

# Section I Multiple-Choice Answer Sheet

1	$A \bigcirc$	B 🔿	С 🔿	D 🔿
2	A 🔿	B 🔿	C 🔿	D 🔿
3	A 🔿	B 🔿	С 🔿	D 🔿
4	A 🔿	B 🔿	С 🔿	D 🔿
5	A 🔿	B 🔿	С 🔿	D 🔿
6	A 🔿	B 🔿	С 🔿	D 🔿
7	A 🔿	B 🔿	C 🔿	D 🔿
8	A 🔿	B 🔿	С 🔿	D 🔿
9	A 🔿	B 🔿	С 🔿	D 🔿
10	A 🔿	B 🔿	С 🔿	D 🔿

# 2014 Year 12 Trial HSC Examination

# Mathematics

# Section I Multiple-Choice Answer Sheet

1	A 🔿	BO	C ()	D 🌀
2	A 🔿	BO	C 🔘	DO
3	Α 💿	BO	C ()	DO
4	A O	B 🔘	C ()	DO
5	A 🔿	B 🎯	C ()	DO
6	A 🔿	BO	С 🎯	DO
7	A 🔿	BO	С 🍥	DO
8	A O	В 🧼	C ()	DO
9	Α 🧼	BO	C ()	DO
10	A 💿	BO	C ()	DO

Year 12 20 Trial HSC 2014  $\int_{-\infty}^{\infty} \frac{1}{2} - \cos x \, dx = 2 \ln x - \sin x + c.$ 2) 4-32 <13 4-3x <13 4-32>-13 52. 773x x < 17, -9<3x 207-3 3) 2x+y=7...0x-2y=1...04x+2y = 14x-2y = 1Sx = 15x=3y=1 $2x^2 - 7x - 15 = (2x + 3)(x - 5)$ 4 (312+75)(2+148)=(613+513)(2+43) = 11/3(2+4/3)= 22/3 + 132 p = 132q = 22,

7) 6(3) - 2k = 8.2k = 10C 8)  $V = \pi \int^2 4 - x^2 dx$ . B 9) 4x2-4x+4y2+24y =-21  $x^2 - x + y^2 + 6y = -21$  $x^2 - x + \frac{1}{4} + y^2 + 6y + 9 = -\frac{2}{4} + \frac{1}{4} + 9$  $(x - \frac{1}{2})^2 + (y + 3)^2 = 4$  $\binom{1}{2}, -3$  R=2 10) a=12. $S_{1}=15$ 15= 12 ---= 12 r= 1-4 21

Question 11 AM B(3,6) A(-1,1) 72. E(1,-4 -3,-9)  $\left(\frac{5-3}{2},\frac{1-9}{2}\right)$ , -4 i) MBE = 6+4 3-1 = 5 y + 4 = 5(x - 1)Y+4= 5x-5 5x - y -9=0 iù  $M_{4E}(-1+1), 1-4)$  $M_{BD}(\frac{3-3}{2}, \frac{6-9}{2})$  $M_{BD}\left(\begin{array}{c}0\\2\end{array}\right)$  $M_{AE}\left(0,-\frac{3}{2}\right)$ ABED is a parallelogram (Diagonals, bisect each other)

(iii) A(-1,1)  $\frac{5(-1) - (1) - 9}{5^2 + 1^2}$ = 1526  $\sqrt{(3-1)^2+(6+4)^2}$ BE= 104 2 = 2,26  $= 2\sqrt{26} \times \frac{15}{\sqrt{26}}$  $=30 u^{2}$  $) \gamma = |n\chi|$ 6  $\gamma' = \frac{1}{\chi}$ Q x = e y' = ? $\gamma = 1$ e. MA == e  $=\frac{1}{e}(x-e)$ Y=x -1 OR x-ey-e=

(i)  $y - 5 = x^2 - 2x$ y-5+1= x2-2x+1  $y - 4 = (x - 1)^2$ V : (1, 4)a=1/4 y = 2x - 2iii ax=2,y' = 2 $m_N = -\frac{1}{2}$  $y - 5 = -\frac{y}{2}(x - 2)$ 2y - 10 = -x + 2x + 2y - 12 = 0or  $y = 6 - \frac{\pi}{2}$ 

Question 12  $P(1st) = \frac{3}{200}$ 3 × 2 200 × 199 ) P(1st +2nd) = = 319900 ìii  $P(2nd) = \frac{197}{200} \times \frac{3}{199}$ = 591 39800  $P(none) = \frac{197}{200} \times \frac{197}{199}$ 9653 9950  $dx(e^{3x}\tan x) = 3e^{3x}\tan x + e^{3x}sec^{2}x$ e3x (3tanx + secx)  $\frac{(sin)(-1)}{(s-x)^2} = \cos((s-x) - sinx.(-1))$  $= \frac{5\cos x - x\cos x + \sin x}{(5 - x)^2}.$ 

 $\int \frac{1}{1-2x} dx = -\frac{1}{2} \ln(1-2x) + C \sqrt{2}$ Ci  $\int_0^T \sec^2 \frac{2\pi}{3} dx = \left[ 3\tan^2 \frac{\pi}{3} \right]_0^T \sqrt{\frac{\pi}{3}}$ = 3tan Iz =33 /  $| \alpha + \beta = \frac{1}{2} \sqrt{2}$ x13=-15 V  $\Delta + \dot{B} = \Delta + \Delta f$ 12/15/2 0 = - 15

Question 13 P= 6(2+(n-1)×3) ai) = 6(2+3n-3)= 18n - 6ii)  $S_n = \frac{n}{2} (18n - 6 + 2x6)$  $= \frac{n}{2}(18n+6)$  $= 9n^2 + 3n$ 111  $600 = 90^{2} + 3n$  $3n^{2} + n - 200 = 0$ (3n + 25)(n - 8) = 0n=-22, 8. n = 8 $b_i$ )  $f'(x) = 3x^2 - 6x - 9$ f''(pc) = 6x - 6i) Stat pts @f'(x)=0  $0 = 3(x^2 - 2x - 3)$ (x-3)(x+1)=0x = 3, -1

@x = -1, y = 27y'' = -6-6= -12 (-1,27) is a max tp. V (ax = 3, y = (-5))y'' = 18 - 6= 12. - (3,-5) is a min t.p. (3,-12) V ii) POI @ y"=Q 0 = 6x - 6Check DC 0 1 2 V" - 0 + (1,11) is a POI 1-1,27) 11 N 22 3,-5

XXI B 25 km/hr 200 850 20 km/hr. A AOB = 20 + 85=  $105^{\circ}$ A0 = 40 kmB0 = 50 km AB2 = 402 +502 - 2×40×50 × Coslos? V = 5135.28 - - -AB = 72 km

Question 14 ai) Diagonals in a rhombus bisect the angles they pass through. i) FC is common. LBCA=LDCA (given (i)) BC = DC (equal sides in a rhombus) . ABFC=ADFC (SAS) iii) LFBC=90° (alt L's on 11 lines) in) LFDC=90° (comes L's in congruent d's  $b_1$   $N(t) = Ae^{0.1St}$ dN = 0.15 Ae0.15+ V = 0.15N. AN XN ii) @1=3, N=1.5×108  $1.5 \times 10^8 = Ae^{0.45}$  $A = \frac{1.5 \times 10^8}{e^{0.45}}$  $= 9.6 \times 10^{7}$ me 96000000

iii) N = 2. $2 = e^{0.15t}$ n2 = 0.15tt = ln2= 4.62. =4,6 days V C A = 10 (52.8 + 19.6 + 62.8) V = 450.7= 451 m<sup>2</sup> V d) (cosx+2)(2cosx+1)=0  $\cos x = -2, -\frac{1}{2}, \cos x \neq -2,$ x=25,45 \$ V

Question 15  $A = \int e^{x} e^{x} - 2 - e^{-x} dx. \quad \checkmark$  $= \left[ e^{x} - 2x + e^{-x} \right]^{2} \checkmark$  $=(e^{2}-4+e^{-2})-(e-2+e^{-1})$  $= e^2 - e - 2 - e^{-1} + e^{-2}$ . ii)  $y = e^{x} - 2$  $y = e^{-x}$  $e^{x} - 2 = e^{-x}$  $e^{\alpha}-2=1$  $e^{2x} - 2e^{x} = 1$  $e^{2\alpha} - 2e^{\alpha} - 1 = 0$  $\tilde{i}$ u² #-2u-1=0  $= 1 \pm \sqrt{2}$  $e^{x} = 1 \pm \sqrt{2}$ .  $\sqrt{2}$  $x = \ln(1+52) = 0.881$  y = 0.414

bi) @v=0, +=?  $0 = 2 \sin t + 1$ Sint = - 1/2  $t = 7\pi 11\pi 5;$ Ĩ 3 111 V=3 おくナく管 iv d= j Zsint+1dt / [t-280st - $=(T - 2\cos T) - (0 - 2\cos 0)$  $= \pi + 2 + 2$  $= \pi + 4$ 

Question 16 ai) r = 0.004A= 450×36 =\$10800\$16200 V ii)  $A_{1} = 450 \times 1.004^{36}$  $A_2 = 450 \times 1.004^{35}$ A36 - 450 × 1.004  $Total = 450 (1.004 + 1.004^2 + ... + 1.004^2)$  $=450 \times 1.004(1.004^{36}-1)$ 0.004 =\$17 456.70 =\$17 457  $i_{1,1}$  = 0.05 = 4 = 0.0125 A = 12000 (1.0025)8 V = 13253.83 =\$13254

 $= \sqrt{(15 + 35 + \frac{3}{15})}$  $(9+3x+1z+9)^{2}=$ 9+ - c+ xs+ - x) == (9+hx+x+hx)== f(2+2)(2+3) M  $H = N (BE) \times (BE)$ 9 = hacラミト 1 a  $\frac{Q}{A} = \frac{Q}{A}$ y (equiangular) JFCDIILL DAE (IZV7=7017 a mins (x-0b)x= 06-081=0347 06 = -081 = 3047 (2457 x=70=17 tor ( PECD= CDAE= do (7,2 ou a struight line) (19

13 Ą 67 <del>}</del>  $\mathcal{X}$ ., 612 0 dA. W 62 3 Ć . . いうの V  $\square$ CM 6.cm  $\leq \rho$ ¢  $\frac{d}{d}$  $\frac{12}{\lambda^3}$ 4 dzA (  $\lambda = 2$ ٢. >0 ĺ m , <u>.</u>