

2010

**TRIAL
HIGHER SCHOOL CERTIFICATE**

GIRRAWEEN HIGH SCHOOL

Mathematics Extension 1

General Instructions:

- Reading Time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen.
- Board - approved calculators may be used
- A table of standard integrals is provided at the back of this paper.
- All necessary working should be shown in every question.

Total marks – 84

- Attempt Questions 1 – 7
- All questions are of equal value

Total marks – 84

Attempt Questions 1 –7

All questions are of equal value.

Question 1 (12 marks). Start on a SEPARATE page.

Marks

- (a) The line $y = mx$ makes an angle of 45° with the line $y = 2x - 3$. Find the possible values of m . 2
- (b) Find the coordinates of the point $P(x, y)$ which divides the interval joining $A(-4, -6)$ and $B(6, -1)$ externally in the ratio 3:2. 2
- (c) Solve for x : $\frac{2x+1}{x-1} \geq 3$ 2
- (d) Differentiate $y = x \tan^{-1} \frac{x}{2}$ 3
- (e) Use the substitution $u = \sqrt{x}$ to evaluate $\int_1^4 \frac{dx}{x + \sqrt{x}}$ 3

Question 2 (12 marks). Start on a SEPARATE page.

(a) Find the coefficient of x^9 in the expansion of $\left(x^2 + \frac{2}{x}\right)^{12}$ 2


(b) Evaluate: $\lim_{x \rightarrow 0} \frac{\sin 6x}{7x}$ 2

(c) If $f(x) = 4 \cos^{-1} \frac{x}{3}$, find

(i) the domain and range of $f(x)$. 2

(ii) Sketch the curve. 2

(d)




In the above figure, AP is a tangent to the circle at A . $PBCQ$ and ADQ

are straight lines. Prove that $\angle PAB = \frac{1}{2}(\angle CFD + \angle CQD)$ 4

Question 3 (12 marks) Start on a SEPARATE page.

- (a) A pole, OT , of length l metres stands on horizontal ground. The pole leans towards the east, making an angle of θ with the vertical. From P , a metres south of O , the elevation of T is α .



- (i) Copy the diagram above onto your booklet. Find expressions, in terms of l and θ for OM and MT . 2
- (ii) Prove that $PM = l \cos \theta \cot \alpha$. 1
- (iii) Prove that $l^2 = \frac{a^2}{\cos^2 \theta \cot^2 \alpha - \sin^2 \theta}$ 3
- (iv) Find the length of the pole, to the nearest metre, if $a = 25$, $\theta = 20^\circ$ and $\alpha = 24^\circ$. 1
- (b) A tea party is arranged for 16 people along two sides of a long table with 8 chairs on each side. Four persons wish to sit on one particular side and two on the other side. In how many ways can they be seated? 2
- (c) In an election 40% of the voters favoured Party A. If an interviewer selected 5 voters at random, what is the probability that
- (i) exactly three of them favoured Party A.
 - (ii) A majority of those selected favoured Party A
 - (iii) At most two favoured Party A. 3

Question 4 (12 marks). Start on a SEPARATE page.

- (a) Prove the following by the Principle of mathematical induction.

$$\log 2 + \log \left(\frac{3}{2}\right) + \log \left(\frac{4}{3}\right) + \dots + \log \left(\frac{n}{n-1}\right) = \log n \text{ for all}$$

integers $n \geq 2$.

3

- (b) $P(2ap, ap^2)$ is a point on the parabola $x^2 = 16y$. The equation of the normal at P is given by $x + py = 4p^3 + 8p$.

- (i) Find the point of intersection R of the normals at P and Q , the end points of focal chord PQ .

2

- (ii) Find the locus of R .

2

- (c) For the function $y = \frac{2x^2 - 2}{x^2 - 9}$

- (i) Write down the equations of horizontal and vertical asymptotes.

2

- (ii) Sketch the curve showing intercepts with axes and asymptotes.

3

Question 5 (12 marks)

- (a) By expanding both sides of the identity
- $(1+x)^5(1+x)^5 = (1+x)^{10}$
- , show

$$\text{that } \sum_{k=0}^5 \binom{5}{k}^2 = \binom{10}{5} \quad 3$$

- (b) (i) Write the expansion of
- $(1+x)^n$
- .
- 1

(ii) By integrating, show that

$$\binom{n}{0} + \frac{1}{2} \binom{n}{1} + \frac{1}{3} \binom{n}{2} + \dots + \frac{1}{n+1} \binom{n}{n} = \frac{2^{n+1} - 1}{n+1} \quad 3$$

- (c) The rate at which an object warms in air is proportional to the difference between its temperature
- T
- and the constant temperature
- A
- of the surrounding air. This rate can be expressed by the differential equation

$$\frac{dT}{dt} = k(T - A),$$

Where t is the time in minutes, T and A are measured in degrees centigrade, and k is a constant.


- (i) Show that $T = A + Ce^{kt}$, where C is a constant is a solution of the differential equation. 2
- (ii) An object warms from 10°C to 15°C in 20 minutes. The air temperature surrounding the object is 25°C . Determine the temperature of the object after a further 30 minutes have passed. Give your answer to the nearest degree. 2
- (iii) Using the equation for T , given in part (i), explain the behaviour of T as t increases to large values. 1

Question 6 (12 marks). Start on a SEPARATE page.

- (a) (i) Given $f(x) = x \sin^{-1}x + \sqrt{1-x^2}$. Find $f'(x)$. 3
- (ii) Hence evaluate $\int_0^{\frac{1}{2}} \sin^{-1}x \, dx$ 2
- (b) (i) show that there exists a root of the equation $\tan x - x = 0$ between $x = 4$ and $x = 4.5$. 1
- (ii) By halving the interval twice find an approximate value of the root Correct to 1 decimal place. 2
- (c) Assume tides at a harbour rise and fall in SHM. At low tide the harbour is 12 m deep, and at high tide 17 m deep. Low tide is at 9-00 am and high tide at 3.00 pm. Assuming a ship needs 14 m to go safely,
- (i) at what time can the ship go into the harbour. 3
- (ii) if the ship take 30 minutes to go out, before what time must it depart the harbour. 1

Question 7 (12 marks). Start on a SEPARATE page.

(a)



A small funnel in the shape of a cone is being emptied of fluid at the rate of $12 \text{ cm}^3/\text{s}$. The height of the funnel is 20 cm and the radius of the top is 4 cm . How fast is the fluid level dropping when the level stands 5 cm above the vertex of the cone?

3

(b) Given that $x^3 + x^2 - 10 = 0$ has a root between 1 and 2. By taking 2 as the initial value find an approximation to the root using Newton's method, correct to one decimal place.

2


(c) A particle is projected from a point P on horizontal ground, with initial speed $V \text{ m/s}$ at an angle of elevation α to the horizontal. Its equations of motion are $x = 0$ and $y = -g$. The horizontal and vertical component of velocity and displacement of the particle at any time t are given by

$$\frac{dx}{dt} = V \cos \alpha \quad \text{and} \quad \frac{dy}{dt} = V \sin \alpha - gt$$

$$x = Vt \cos \alpha \quad \text{and} \quad y = Vt \sin \alpha - \frac{1}{2}gt^2 \quad (\text{do not prove these})$$

(i) Determine the time of flight of the particle.

2



- (ii) The particle reaches a point Q , as shown, where the direction of the flight makes an angle β with the horizontal. Find an expression for $\tan \beta$. 1

- (iii) Hence show that the time taken to travel from P to Q is

$$\frac{V \sin(\alpha - \beta)}{g \cos \beta} \text{ seconds.} \quad 2$$

- (iv) Consider the case where $\beta = \frac{\alpha}{2}$. If the time taken to travel from P to Q is one third of the total time of flight, find the value of α . 2

End of paper

Trial HSC Extension 1, 2010 - Solutions

Question 1 (12 marks)

(a) $m_1 = m$ $m_2 = 2$

$$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$$

$$1 = \left| \frac{2 - m}{1 + 2m} \right| \quad (2)$$

$$\frac{2 - m}{1 + 2m} = 1 \quad \text{OR} \quad \frac{2 - m}{1 + 2m} = -1$$

$$2 - m = 1 + 2m$$

$$2 - m = -1 - 2m$$


$$3m = 1$$

$$-m = 3$$

$$\underline{\underline{m = \frac{1}{3}}}$$

$$\underline{\underline{m = -3}}$$

(b)



$$x = \frac{(-3 \times 6) + (2 \times -4)}{-3 + 2} = 26$$

$$y = \frac{(-3 \times -1) + (2 \times -6)}{-3 + 2} = 9$$

$$\underline{\underline{P(26, 9)}} \quad (2)$$

(c) $\frac{2x+1}{x-1} \geq 3$


$$(x-1)(2x+1) \geq 3(x-1)^2, \quad x \neq 1$$

$$3(x-1)^2 - (x-1)(2x+1) \leq 0$$

$$(x-1) [3(x-1) - (2x+1)] \leq 0$$

$$(x-1)(3x-3-2x-1) \leq 0$$

$$(x-1)(x-4) \leq 0$$



$$\underline{\underline{1 < x \leq 4}}$$

(d) $y = x \tan^{-1} \frac{x}{2}$

$$y' = x \times \frac{1}{1 + \frac{x^2}{4}} \times \frac{1}{2} + \tan^{-1} \frac{x}{2} \times 1$$

$$= x \times \frac{1}{\frac{4+x^2}{4}} \times \frac{1}{2} + \tan^{-1} \frac{x}{2}$$

$$= x \times \frac{4}{4+x^2} \times \frac{1}{2} + \tan^{-1} \frac{x}{2}$$

$$= \underline{\underline{\frac{2x}{4+x^2} + \tan^{-1} \frac{x}{2}}} \quad (3)$$

$$(c) \int_1^4 \frac{dx}{x + \sqrt{x}}, \quad u = \sqrt{x}$$

$$\frac{du}{dx} = \frac{1}{2\sqrt{x}}; \quad \frac{dx}{du} = 2\sqrt{x}$$

$$dx = 2\sqrt{x} du = 2u du$$

When $x=1$, $u = \sqrt{1} = 1$

When $x=4$, $u = \sqrt{4} = 2$

$$\int_1^2 \frac{2u du}{u^2 + u} = \int_1^2 \frac{2u du}{u(u+1)}$$

$$= \int_1^2 \frac{2 du}{u+1} = 2 \int_1^2 \frac{du}{u+1}$$

$$= 2 \left[\log(u+1) \right]_1^2 \quad (3)$$

$$= 2 \left[\log 3 - \log 2 \right]$$

$$= \underline{\underline{2 \log \frac{3}{2}}}$$

Question 2 (12 marks)

$$(a) T_{r+1} = 12 C_r (x^2)^{12-r} \left(\frac{2}{x}\right)^r$$

$$= 12 C_r x^{24-2r} \frac{2^r}{x^r}$$

$$= 12 C_r x^{24-3r} 2^r$$

$$24 - 3r = 9$$

$$3r = 15$$

$$r = 5$$

$$T_6 = 12 C_5 x^{24-15} 2^5$$

$$= 12 C_5 x^9 2^5 \quad (2)$$

$$\text{Coefficient of } x^9 = \underline{\underline{12 C_5 x^5}}$$

$$(b) \lim_{x \rightarrow 0} \frac{\sin 6x}{7x}$$

$$= \lim_{x \rightarrow 0} \frac{\sin 6x}{6x} \times \frac{6x}{7x} \quad (2)$$

$$= \underline{\underline{\frac{6}{7}}} \quad \left(\because \lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1 \right)$$

$$(c) f(x) = 4 \cos^{-1} \frac{x}{3}$$

$$(i) D: -1 \leq \frac{x}{3} \leq 1$$


$$-3 \leq x \leq 3 \quad (2)$$

$$R: 0 \leq y \leq 4\pi$$

$$x = -3, \quad y = 4 \cos^{-1}(-1) = 4\pi$$


$$x = 0, \quad y = 4 \cos^{-1} 0 = 4 \times \frac{\pi}{2} = 2\pi$$

$$x = 3, \quad y = 4 \cos^{-1} 1 = 0$$



Question 3 (12 marks)

(d)



$\angle PAB = \angle ACB$ (angle between tangent and chord is equal to the angle in the alternate segment)

$\angle ACB = \angle ADB$ (angles at the circumference standing on the same chord)

$\angle BCA = \angle CQF + \angle CFQ$
(exterior angle of $\triangle FQC$)

$\angle ADF = \angle DFQ + \angle DQF$
(exterior angle of $\triangle FQD$)

$\angle BCA + \angle ADF$

$= \angle CQF + \angle CFQ + \angle DFQ + \angle DQF$

$= \angle CQF + \angle DQF + \angle CFQ + \angle DFQ$


$= \angle CQD + \angle CFD$

But $\angle BCA + \angle ADF = 2\alpha$

$2\alpha = \angle CQD + \angle CFD$ (4)

$\alpha = \frac{1}{2}(\angle CQD + \angle CFD)$

(e)



(i) $\sin \theta = \frac{OM}{l}$

$OM = l \sin \theta$ (2)

$\cos \theta = \frac{MT}{l}$; $MT = l \cos \theta$

(ii) $\cot \alpha = \frac{PM}{MT}$; $PM = MT \cot \alpha$ (1)
 $= l \cos \theta \cot \alpha$

(iii) $PM^2 - OM^2 = a^2$

$l^2 \cos^2 \theta \cot^2 \alpha - l^2 \sin^2 \theta = a^2$

$l^2 (\cos^2 \theta \cot^2 \alpha - \sin^2 \theta) = a^2$

$l^2 = \frac{a^2}{\cos^2 \theta \cot^2 \alpha - \sin^2 \theta}$ (3)

(iv) $l^2 = \frac{25^2}{\cos^2 20^\circ \cot^2 24^\circ - \sin^2 20^\circ}$

$l = 12$ (1)

(b) Total number of ways in which 16 people can be seated = $8P_4 \times 8P_2 \times 10!$ (2)

$$(i) P(x=3) = {}^5C_3 (0.4)^3 (0.6)^2 = 0.2304$$

$$(ii) P(x=3) + P(x=4) + P(x=5) = 0.2304 + {}^5C_4 (0.4)^4 (0.6)^1 + {}^5C_5 (0.4)^5 (0.6)^0 = 0.31744$$
 (3)

$$(iii) P(x=0) + P(x=1) + P(x=2) = 1 - 0.31744 = \underline{\underline{0.68256}}$$

Question 4 (12 marks)

(a) when $n=2$,

$$\text{LHS} = \log \frac{2}{2-1} = \log 2$$

$$\text{RHS} = \log 2$$

$$\text{LHS} = \text{RHS}$$

\therefore the result is true for $n=2$

Assume the result is true for $n=k$

$$\text{i.e. } \log 2 + \log \left(\frac{3}{2}\right) + \dots + \log \left(\frac{k}{k-1}\right) = \log k \text{ --- (1)}$$

To prove that the result is true for $n=k+1$

$$\text{i.e. } \log 2 + \log \left(\frac{3}{2}\right) + \log \left(\frac{4}{3}\right) + \dots + \log \left(\frac{k}{k-1}\right) + \log \left(\frac{k+1}{k}\right) = \log k+1$$

Now

$$\log 2 + \log \frac{3}{2} + \dots + \log \left(\frac{k}{k-1}\right) + \log \left(\frac{k+1}{k}\right) = \log k + \log \left(\frac{k+1}{k}\right) \text{ by assumption (1)}$$

$$= \log \left(k \times \frac{k+1}{k}\right)$$

$$= \log(k+1) \text{ (3)}$$

\therefore the result is true for $n=k+1$

Hence by the principle of mathematical induction, the result is true for $n \geq 2$

(b) (i) Normal at P

$$x + py = 4p^3 + 8p \text{ --- (1)}$$

Normal at Q

$$x + qy = 4q^3 + 8q \text{ --- (2)}$$

① - ② gives

$$y(p-q) = 4(p^3 - q^3) + 8(p-q)$$

$$y(p-q) = 4(p-q)(p^2 + pq + q^2) + 8(p-q)$$

$$y = 4(p^2 + pq + q^2) + 8$$

$$= 4(p^2 - 1 + q^2) + 8$$

$$= 4p^2 - 4 + 4q^2 + 8$$

$$= 4p^2 + 4q^2 + 4$$

Substitute in ①

$$2x + p(4p^2 + 4q^2 + 4) = 4p^3 + 8p$$

$$2x + 4p^3 + 4pq^2 + 4p = 4p^3 + 8p$$

$$2x + 4pq^2 = 4p$$

$$2x + 4pq \times q = 4p$$

$$2x - 4q = 4p$$

$$2x = 4p + 4q$$

$$= 4(p+q)$$

$$2x = 4(p+q) \text{ --- ③}$$

$$y = 4p^2 + 4q^2 + 4 \text{ --- ④}$$

$$\text{From ③ } p+q = \frac{2x}{4}$$

$$\text{From ④ } \frac{y}{4} = p^2 + q^2 + 1$$

$$p^2 + q^2 = \frac{y}{4} - 1$$

$$p^2 + q^2 = (p+q)^2 - 2pq$$

$$= (p+q)^2 - 2x - 1$$

$$= (p+q)^2 + 2$$

$$\frac{y}{4} - 1 = \left(\frac{2x}{4}\right)^2 + 2$$

$$\frac{y}{4} - 1 = \frac{x^2}{16} + 2$$

$$4y - 16 = x^2 + 32 \text{ --- ③}$$

$$x^2 = 4y - 48$$

$$\underline{\underline{x^2 = 4(y-12)}}$$

$$\textcircled{c} y = \frac{2x^2 - 2}{x^2 - 9} = \frac{2(x+1)(x-1)}{(x+3)(x-3)}$$

Vertical asymptotes

$$x = -3 \text{ and } x = 3$$

$$\lim_{x \rightarrow \infty} \frac{2x^2 - 2}{x^2 - 9}$$

$$= \lim_{x \rightarrow \infty} \frac{2 - \frac{2}{x^2}}{1 - \frac{9}{x^2}} \text{ --- ③}$$

$$= 2 \text{ (since } \lim_{x \rightarrow \infty} \frac{1}{x^2} = 0)$$

Horizontal asymptote is

$$y = 2$$


x intercepts

$$y=0 \Rightarrow 2(x+1)(x-1)=0$$

$$x=-1 \text{ or } x=1$$

y intercept

$$x=0 \Rightarrow y = \frac{-2}{-9} = \frac{2}{9} = 0.2$$

Question 5 (12 marks)

$$(a) (1+x)^5 (1+x)^5 = (1+x)^{10}$$

$$\left[{}^5C_0 + {}^5C_1 x + {}^5C_2 x^2 + {}^5C_3 x^3 + {}^5C_4 x^4 + {}^5C_5 x^5 \right] \left[{}^5C_0 + {}^5C_1 x + {}^5C_2 x^2 + {}^5C_3 x^3 + {}^5C_4 x^4 + {}^5C_5 x^5 \right]$$

$$= {}^{10}C_0 + {}^{10}C_1 x + {}^{10}C_2 x^2 + {}^{10}C_3 x^3 + {}^{10}C_4 x^4 + {}^{10}C_5 x^5 + \dots + {}^{10}C_{10} x^{10}$$

Equating coefficients of x^5 on both sides, page 7

$${}^5C_0 \times {}^5C_5 + {}^5C_1 \times {}^5C_4 + {}^5C_2 \times {}^5C_3 + {}^5C_3 \times {}^5C_2 + {}^5C_4 \times {}^5C_1 \\ + {}^5C_5 \times {}^5C_0 = {}^{10}C_5$$

$${}^5C_0 \times {}^5C_0 + {}^5C_1 \times {}^5C_1 + {}^5C_2 \times {}^5C_2 + {}^5C_3 \times {}^5C_3 + {}^5C_4 \times {}^5C_4 \\ + {}^5C_5 \times {}^5C_5 = {}^{10}C_5 \quad (\text{since } nC_r = nC_{n-r})$$

$$({}^5C_0)^2 + ({}^5C_1)^2 + ({}^5C_2)^2 + ({}^5C_3)^2 + ({}^5C_4)^2 + ({}^5C_5)^2 = {}^{10}C_5$$

$$\text{i.e. } \sum_{k=0}^5 ({}^5C_k)^2 = {}^{10}C_5 \quad (3)$$

$$(b) (i) (1+x)^n = nC_0 + nC_1x + nC_2x^2 + \dots + nC_r x^r + \dots + nC_n x^n \quad (1)$$

$$(ii) \int_0^1 (1+x)^n dx = \int_0^1 (nC_0 + nC_1x + nC_2x^2 + \dots + nC_n x^n) dx$$

$$\left[\frac{(1+x)^{n+1}}{n+1} \right]_0^1 = nC_0 x + nC_1 \frac{x^2}{2} + nC_2 \frac{x^3}{3} + \dots + nC_n \frac{x^{n+1}}{n+1} \Big|_0^1$$

$$\frac{2^{n+1}}{n+1} - \frac{1}{n+1} = nC_0 + \frac{nC_1}{2} + \frac{nC_2}{3} + \dots + \frac{nC_n}{n+1} \quad (3)$$

$$\underline{\underline{nC_0 + \frac{nC_1}{2} + \frac{nC_2}{3} + \dots + \frac{nC_n}{n+1} = \frac{2^{n+1} - 1}{n+1}}}$$

$$(1) (i) T = A + Ce^{kt}$$

$$\frac{dT}{dt} = Ce^{kt} \times k$$

$$= k \times Ce^{kt} \quad (2)$$

$$= k(T-A) \quad (\text{since } e^{kt} = T-A)$$

$\therefore T = A + Ce^{kt}$ is a solution of the equation.

$$\frac{dT}{dt} = k(T-A)$$

$$(ii) T = 25 + Ce^{kt}$$

$$\text{when } t=0, T = 10^\circ\text{C}$$

$$10 = 25 + C$$

$$C = 10 - 25 = -15$$

$$\therefore T = 25 - 15e^{kt}$$

$$\text{when } t=20, T = 15^\circ$$

$$15 = 25 - 15e^{20k}$$

$$15e^{20k} = 10$$

$$e^{20k} = \frac{10}{15}$$

$$20k = \log\left(\frac{10}{15}\right)$$

$$k = \frac{1}{20} \log\left(\frac{10}{15}\right)$$

$$\text{when } t=50, T = ?$$

$$T = 25 - 15e^{50k}$$

$$= 25 - 15e^{50 \times \frac{1}{20} \log\left(\frac{10}{15}\right)} \quad (2)$$

$$= 20^\circ\text{C} \quad (\text{to the nearest degree})$$

$$(iii) k = \frac{1}{20} \log\left(\frac{10}{15}\right) = -0.02$$

$$T = 25 - 15e^{-0.02t}$$

Since $k < 0$, as $t \rightarrow \infty$ $e^{kt} \rightarrow 0$

\therefore as t increases indefinitely the object's temperature approaches air temperature. (1)

Question 6 (12 marks)

$$(a) (i) f(x) = x \sin^{-1}x + \sqrt{1-x^2}$$

$$f'(x) = x \times \frac{1}{\sqrt{1-x^2}} + \sin^{-1}x \times 1 + \frac{1}{2\sqrt{1-x^2}} \times -2x$$

$$= \frac{x}{\sqrt{1-x^2}} + \sin^{-1}x - \frac{x}{\sqrt{1-x^2}} = \sin^{-1}x$$

$$\frac{d}{dx} (x \sin^{-1}x + \sqrt{1-x^2}) = \sin^{-1}x \quad (3)$$

$$\int_0^{\frac{1}{2}} \sin^{-1}x dx = \left[x \sin^{-1}x + \sqrt{1-x^2} \right]_0^{\frac{1}{2}}$$

$$= \left(\frac{1}{2} \sin^{-1}\left(\frac{1}{2}\right) + \sqrt{1-\frac{1}{4}} \right) - (0 + \sqrt{1})$$

$$= \frac{1}{2} \times \frac{\pi}{6} + \frac{\sqrt{3}}{2} - 1 = \frac{\pi}{12} - 1 + \frac{\sqrt{3}}{2}$$

$$= 0.128$$


(2)

(b)(i) $\tan x - x = 0$

$\tan 4 - 4 = -2.8$

$\tan 4.5 - 4.5 = 0.14$

Since $f(4) < 0$ and $f(4.5) > 0$ there is a root of $f(x) = 0$ between $x = 4$ and




$f(4.25) = \tan 4.25 - 4.25 = -2.24$

$f(4.375) = \tan 4.375 - 4.375 = -1.52$

Approximate value of the root

$= \frac{4.375 + 4.5}{2} = 4.4$

(c)(i)



$T = 2 \times (3:00 \text{ pm} - 9:00 \text{ am})$
 $= 2 \times 6 \text{ h}$
 $= 12 \text{ h}$

$T = \frac{2\pi}{n} \quad n = \frac{2\pi}{T} = \frac{2\pi}{12} = \frac{\pi}{6}$

Let 9:00 am denote $t = 0$

$x = a \cos(nt + \pi)$
 $= 2.5 \cos\left(\frac{\pi}{6}t + \pi\right)$

when the harbour is 14m deep

$x = -0.5$

$-0.5 = 2.5 \cos\left(\frac{\pi}{6}t + \pi\right)$

$\cos\left(\frac{\pi}{6}t + \pi\right) = -\frac{1}{5}$

$\frac{\pi}{6}t + \pi = \pi - \cos^{-1}\left(\frac{1}{5}\right), \pi + \cos^{-1}\left(\frac{1}{5}\right)$

$\frac{\pi t}{6} = -1.3694, 1.3694$

$\frac{\pi t}{6} = 1.3694$ (t can't be negative)

$t = 1.3694 \times \frac{6}{\pi}$

$= 2.6154$

$= 2 \text{ hr } 37 \text{ minutes.}$ (3)


Time taken from A to B

$= 3 \text{ hr} - 2 \text{ hr } 37 \text{ min}$

$= 23 \text{ min}$

The ship can go into the harbour


at 11:37 am



The ship must depart before

5:53 pm. (1)

Question 7 (12 marks)



$$V = \frac{1}{3} \pi r^2 h$$

$$\frac{dV}{dt} = 12 \text{ cm}^3/\text{s}$$

By similar triangles

$$\frac{r}{4} = \frac{h}{20}$$

$$r = \frac{4h}{20} = \frac{h}{5}$$

$$V = \frac{1}{3} \pi \times \frac{h^2}{25} \times h$$

$$= \frac{\pi h^3}{75}$$

(3)

$$\frac{dV}{dt} = \frac{dV}{dh} \times \frac{dh}{dt}$$

$$\frac{dV}{dh} = \frac{\pi \times 3h^2}{75} = \frac{\pi h^2}{25}$$

$$12 = \frac{\pi h^2}{25} \times \frac{dh}{dt}$$

$$\frac{dh}{dt} = \frac{12 \times 25}{\pi h^2}$$

When $h = 5$,

$$\frac{dh}{dt} = \frac{12 \times 25}{\pi \times 25} = 3.82$$

The fluid level is dropping at the rate of 3.82 cm/s.

$$(b) f(x) = x^3 + x^2 - 10$$

$$f'(x) = 3x^2 + 2x$$

$$a = 2$$

$$a_1 = 2 - \frac{f(2)}{f'(2)}$$

$$= 2 - \left[\frac{8 + 4 - 10}{3 \times 4 + 4} \right] = 2 - \frac{1}{8} = 1.875$$

$$= 1.9$$

$$a_2 = 1.875 - \frac{f(1.875)}{f'(1.875)}$$

$$= 1.875 - \left[\frac{(1.875)^3 + (1.875)^2 - 10}{3 \times (1.875)^2 + 2 \times 1.875} \right]$$

$$= 1.867 = 1.9$$

(2)

$$a_1 = a_2 = 1.9$$

\therefore the root of $f(x) = 0$ correct to one decimal place is 1.9

(c) (i) When the particle strikes the ground $y = 0$

$$Vt \sin \alpha - \frac{1}{2} g t^2 = 0$$

$$t(V \sin \alpha - \frac{1}{2} g t) = 0$$

$$t = 0 \quad \text{or} \quad V \sin \alpha = \frac{1}{2} g t$$


$$t=0 \text{ or } 2V\sin\alpha = gt$$

$$t=0 \text{ or } t = \frac{2V\sin\alpha}{g}$$

Now $t=0$ refers to the instant of projection (2) and hence $t = \frac{2V\sin\alpha}{g}$

is the required time, the time of flight.

(ii)



(1)

$$\tan\beta = \frac{dy/dt}{dx/dt}$$

$$= \frac{V\sin\alpha - gt}{V\cos\alpha}$$

$$\text{iii) } \frac{\sin\beta}{\cos\beta} = \frac{V\sin\alpha - gt}{V\cos\alpha}$$

$$V\sin\beta\cos\alpha = V\sin\alpha\cos\beta - gt\cos\beta$$

$$gt\cos\beta = V\sin\alpha\cos\beta - V\sin\beta\cos\alpha$$

$$gt\cos\beta = V(\sin\alpha\cos\beta - \cos\alpha\sin\beta)$$

$$gt\cos\beta = V\sin(\alpha - \beta)$$

(2)

$$t = \frac{V\sin(\alpha - \beta)}{g\cos\beta}$$

(iv) when $\beta = \frac{\alpha}{2}$ we have

$$t = \frac{V\sin(\alpha - \frac{\alpha}{2})}{g\cos\frac{\alpha}{2}} = \frac{V\sin\frac{\alpha}{2}}{g\cos\frac{\alpha}{2}}$$

$$= \frac{V}{g} \tan\frac{\alpha}{2}$$

$$\text{Given that } \frac{V}{g} \tan\frac{\alpha}{2} = \frac{1}{3} \frac{2V\sin\alpha}{g}$$

$$\tan\frac{\alpha}{2} = \frac{1}{3} \times 2\sin\alpha$$

$$3\tan\frac{\alpha}{2} = 2\sin\alpha$$

$$= \frac{2 \times 2\tan\frac{\alpha}{2}}{1 + \tan^2\frac{\alpha}{2}}$$

$$3 = \frac{4}{1 + \tan^2\frac{\alpha}{2}}$$

$$4 = 3 + 3\tan^2\frac{\alpha}{2}$$

(2)

$$3\tan^2\frac{\alpha}{2} = 1$$

$$\tan^2\frac{\alpha}{2} = \frac{1}{3}$$

$$\tan\frac{\alpha}{2} = \frac{1}{\sqrt{3}} \left(\frac{\alpha}{2} \text{ is acute} \right)$$

$$\frac{\alpha}{2} = \frac{\pi}{6}$$

$$\alpha = \frac{\pi}{3}$$

=====