


**QUESTION 1. (Start on a new sheet of paper)****MARKS**

- a) If  $y = x \tan^{-1} x$ , find  $\frac{dy}{dx}$ . 2
- b) If  $f(x) = \sin^{-1}(1 - 2x)$ , show that  $f'(x) = \frac{-1}{\sqrt{x-x^2}}$ . 3
- c)  $P(x)$  is an odd polynomial of degree 3. It has  $(x+4)$  as a factor and, when it is divided by  $(x-3)$ , the remainder is 21. Find  $P(x)$ . 3
- d) By making the substitution  $u = x - 2$ , evaluate  $\int_4^5 \frac{x(x-4)}{(x-2)} dx$  4

**QUESTION 2. (Start on a new sheet of paper)**


- a) The points  $A, B, C$  and  $D$  lie on the circumference of a circle centred at  $O$ .  $CD$  is a diameter of the circle and  $AB$  is parallel to  $CD$ .  $\angle ACD = x^\circ$ . Find an expression for  $\angle ACB$  in terms of  $x$ . 3



- b) Use the method of mathematical induction to show that the expression  $9^n - 8n - 1$  is divisible by 64 for all integers  $n \geq 2$ . 5
- c) i) Given that  ${}^n C_r = \frac{n!}{r!(n-r)!}$ , show that  $\frac{r \times^n C_r}{{}^n C_{r-1}} = n - r + 1$  1
- ii) Hence show that : 3
- $$\frac{{}^n C_1}{{}^n C_0} + \frac{2 \times^n C_2}{{}^n C_1} + \frac{3 \times^n C_3}{{}^n C_2} + \dots + \frac{n \times^n C_n}{{}^n C_{n-1}} = \frac{n}{2}(n+1).$$

**QUESTION 3. (Start on a new sheet of paper)****MARKS**

- a) Evaluate the definite integral  $\int_0^{\frac{1}{\sqrt{2}}} \frac{x}{\sqrt{1-x^4}} dx$  by using the substitution  $u = x^2$ . **4**
- b) The point  $P(2ap, ap^2)$  is on the parabola  $x^2 = 4ay$  and a straight line  $OQ$  is drawn through the vertex parallel to the tangent at  $P$ . This line meets the parabola again at  $Q$  and the tangent to the parabola at  $Q$  meets the tangent at  $P$  in  $R$ , as shown in the diagram.



You are given that the tangent at  $P$  has equation  $y = px - ap^2$ .

- i) Write down the equation of the line  $OQ$ . **1**
- ii) Find the coordinates of  $Q$  in terms of  $a$  and  $p$ . **2**
- iii) Show that the equation of the tangent at  $Q$  is  $y = 2px - 4ap^2$ . **1**
- iv) Find the coordinates of  $R$ . **2**
- v) Show that, as  $P$  varies on the parabola,  $R$  moves on another parabola whose equation is  $x^2 = \frac{9}{2}ay$ . **2**

**QUESTION 4. (Start on a new sheet of paper)****MARKS**

- a) Consider the function  $f(x) = \frac{e^x}{(1+e^x)}$ .
- i) Find  $f'(x)$  and deduce that  $f(x)$  is increasing for all  $x$ . **2**
  - ii) State the range of  $f(x)$ . **1**
  - iii) Find the inverse function  $f^{-1}(x)$  **2**
  - iv) Draw  $y = f(x)$  and  $y = f^{-1}(x)$  on the same diagram. **2**
- b) A particle moves in a straight line on the  $x$  axis. At time  $t$  its velocity is  $v$  and its acceleration is  $a$ .
- i) If  $a = 4x - 4$  and initially  $x = 6$  and  $v^2 = 64$ , show that  $v^2 = 4x^2 - 8x - 32$ . **2**
  - ii) Use this expression for  $v^2$  to find the possible values of  $x$ . **1**
  - iii) Describe the motion of the particle if  $v = -8$  initially. **2**

**QUESTION 5. (Start on a new sheet of paper)**


- a) A particle moves in a straight line with displacement in centimetres from the point  $x = 0$  at time  $t$  seconds given by  $x = \sin 3t + 2 \cos 3t$  for  $t \geq 0$ .
- i) Express  $x$  in the form  $R \sin(3t + \alpha)$  where  $R > 0$  and  $0 \leq \alpha \leq \frac{\pi}{2}$ . **2**
  - ii) Show that the motion is simple harmonic. **1**
  - iii) Write down the period of motion. **1**
  - iv) Find at what time, to the nearest tenth of a second, the particle first reaches  $x = 1$ . **2**
- b) Fred deposited \$20,000 at the beginning of January into an account which paid interest at the rate of 0.5% per month compounded monthly. He withdrew \$50 each month from the account each month, immediately after the interest was paid.
- i) How much money was in the account immediately after the first withdrawal? **1**
  - ii) Show that, after making the  $n^{\text{th}}$  withdrawal, his account balance is given by the expression  $\$(10,000 \times 1.005^n + 10,000)$  **3**
  - iii) Find the number of months it will take for his account balance to be \$50,000 **2**

**QUESTION 6. (Start on a new sheet of paper)**

**MARKS**

a) Find the term independent of  $x$  in the expansion of  $\left(\frac{2x^2}{3} - \frac{3}{2x}\right)^9$ . **3**

b) An aeroplane,  $A$ , flying at a height of 2000m observes a stationary blimp,  $B$ , at a height of 3200m drop an object. As the object is dropped, the plane fires a projectile towards it at a speed of 240m/s and at an angle  $\theta$  to the horizontal. The horizontal distance between the plane and the blimp is 3600m at the time that the projectile is fired.



The origin of coordinates,  $O$ , is taken to be the point on the ground below  $A$ .

The particle's coordinates at time  $t$  (secs) are given by :  $x = 240t \cos \theta$ ,  
 $y = 2000 + 240t \sin \theta - \frac{gt^2}{2}$

The coordinates of the dropped object at time  $t$  are :  $x = 3600$  ,  
 $y = 3200 - \frac{gt^2}{2}$

(You may use  $g=10\text{m/s}^2$ )

i) What is the angle  $\theta$  at which the projectile must be fired to intercept the object, and how long does it take to reach it? **3**

ii) At what height does the projectile intercept the object? **1**

c) A man notices two towers, one due North and one in a direction  $N\theta E$  (i.e. at an angle  $\theta$  east of north). The angle of elevation  $\beta$  of both towers is the same but the height of one tower is twice the height of the other. Show that

$$\cos \theta = \frac{5 \cot^2 \beta - \cot^2 \alpha}{4 \cot^2 \beta} \quad \mathbf{5}$$

where  $\alpha$  is the angle of elevation of the top of the taller tower from the top of the shorter.


**QUESTION 7. (Start on a new sheet of paper)****MARKS**

a) A group of four contestants for a quiz game are to be selected at random from a class of eight girls and five boys.

i) What is the probability that the team comprises three girls and one boy? **2**

ii) Find the probability that there are more girls chosen than boys. **2**

b) A water trough takes the shape of a hollow semi-circular prism with length  $l$  and radius  $r$ . It is placed on horizontal ground and filled with water. The surface of the water is at a distance  $h$  below the top of the trough, as shown in the diagram.



i) Show that the area  $A$  of the flat surface area of water is given by

$$A = 2l\sqrt{r^2 - h^2} \quad \mathbf{2}$$

ii) Show that the volume  $V$  of water in the trough is given by

$$V = l \left( r^2 \cos^{-1} \left( \frac{h}{r} \right) - h\sqrt{r^2 - h^2} \right) \quad \mathbf{2}$$

iii) If the water level is falling, show that  $\frac{dV}{dt} = -2l\sqrt{r^2 - h^2} \frac{dh}{dt} = -A \frac{dh}{dt}$ . **3**

iv) On a sunny day, the rate of evaporation at any time (and hence  $-\frac{dV}{dt}$ ) is proportional to  $A$ . Show that the water level falls at a constant rate. **1**

**END OF THE PAPER**

1) a)  $y = x \tan^{-1} x$   
 $\frac{dy}{dx} = \tan^{-1} x + \frac{x}{1+x^2}$  (Product rule)

b)  $f(x) = \sin^{-1}(1-2x)$   
 Let  $u = 1-2x$ ,  $\frac{du}{dx} = -2$   
 $\therefore f'(x) = \frac{1}{\sqrt{1-(1-2x)^2}} \cdot x - 2$   
 $= \frac{-2}{\sqrt{1-(1+4x^2-4x)}} = \frac{-2}{\sqrt{4x-4x^2}} = \frac{-1}{\sqrt{x-x^2}}$

c)  $P(x) = Ax(x+4)(x-4)$   
 $P(3) = 3A \times 7 \times -1 = 21 \therefore A = 1$   
 $\therefore P(x) = -x(x+4)(x-4)$   
 $(= 16x - x^3)$

1) Let  $I = \int_4^5 \frac{x(x-4)}{(x-2)} dx$   
 Let  $u = x-2$   $x=5 \Rightarrow u=3$   
 $x=4 \Rightarrow u=2$   
 $\frac{du}{dx} = 1 \therefore "dx = du"$   
 $\therefore I = \int_2^3 \frac{(u+2)(u-2)}{u} du$   
 $= \int_2^3 \frac{u^2-4}{u} du$   
 $= \int_2^3 u - \frac{4}{u} du$   
 $= \left[ \frac{u^2}{2} - 4 \ln u \right]_2^3$   
 $= \left( \frac{9}{2} - 4 \ln 3 \right) - \left( 2 - 4 \ln 2 \right)$   
 $= \frac{5}{2} - 4 \ln \left( \frac{3}{2} \right)$

2) a) ABCD is a cyclic quadrilateral  
 $\angle CAD = 90^\circ$  (Angle at circumference in semi circle)  
 $\angle BAC = x^\circ$  (Alternate angles are equal  $AB \parallel CD$ )  
 $\therefore \angle BAP = x^\circ + 90^\circ$   
 $\therefore \angle BCD = 180^\circ - (x^\circ + 90^\circ)$  (Opposite angles in cyclic quad. supp.)  
 $= 90^\circ - x^\circ$   
 $\therefore \angle BCA = \angle BCD - \angle ACD$   
 $= 90^\circ - 2x^\circ$

b) When  $n=2$ ,  
 $T_2 = 9^2 - 16 - 1 = 64$   
 which is divisible by 64  
 So induction starts.  
 Assume that  
 $9^k - 8k - 1 = 64A$  for  $k \geq 2$   
 and  $A \in \mathbb{Z}$ .

Then  
 $9^{k+1} - 8(k+1) - 1 = 9 \cdot 9^k - 8k - 9$   
 $= 9(9^k - 1) - 8k$   
 $= 9(9^k - 8k - 1) + 64k$   
 $= 9 \cdot 64A + 64k$   
 $= 64(9A + k)$   
 Thus if true for  $n=k$ , also true for  $n=k+1$ .  
 $\therefore$  By principle of M.I.,  
 $9^n - 8n - 1$  is divisible by 64 for  $n \geq 2$

c) i)  ${}^r C_r = \frac{r!}{(n-r)! r! n!} \cdot (n-r)!$   
 ${}^n C_{r-1} = \frac{n!}{(n-r)! (r-1)! n!}$   
 $= \frac{r(r-1)! (n-r+1)!}{r! (n-r)!}$   
 $= \frac{n-r+1}{r}$

2 c) ii) Using result from 1)  
 $n + (n-1) + (n-2) + \dots + 2 + 1$   
 This an AP with  $a=n, d=-1$   
 $Sum = \frac{n}{2} (2n + (n-1)(-1))$   
 $= \frac{n(n+1)}{2}$

3) a) Let  $I = \int_0^{\sqrt{1/2}} \frac{x dx}{\sqrt{1-x^4}}$   
 Let  $u = x^2$   $x = \sqrt{1/2} \Rightarrow u = 1/2$   
 $"du = 2x dx"$   $x=0 \Rightarrow u=0$   
 $\therefore I = \frac{1}{2} \int_0^{1/2} \frac{du}{\sqrt{1-u^2}}$   
 $= \left[ \frac{1}{2} \sin^{-1} u \right]_0^{1/2}$   
 $= \frac{1}{2} \cdot \frac{\pi}{6} = \frac{\pi}{12}$

b) i)  $y = px$   
 ii) Crosses parabola where  
 $x^2 = 4apx$   
 $x = 4ap$  ( $x \neq 0$ )  
 $y = 4ap^2$   
 $Q$  is  $(4ap, 4ap^2)$


iii)  $Q$  is point with parameter " $2p$ "  
 $\therefore$  Tangent is  $y = (2p)x - a(2p)^2$   
 $y = 2px - 4ap^2$

iv) Two tangents cross at  $R$   
 $px - ap^2 = 2px - 4ap^2$   
 $\Rightarrow x = 3ap$  ( $p \neq 0$ )  
 $y = 2ap^2$   
 $R$  is  $(3ap, 2ap^2)$

v) Eliminate  $p$  from  $x = 3ap$   
 $y = 2ap^2$   
 $R = \frac{x}{3a}, y = 2a \left( \frac{x}{3a} \right)^2$   
 $x^2 = \frac{9ay}{2}$

4) i)  $f(x) = 1 - \frac{1}{1+e^x} = 1 - (1+e^x)^{-1}$   
 $f'(x) = \frac{e^x}{(1+e^x)^2}$   
 $e^x > 0$   $(1+e^x)^2 > 0 \therefore f'(x) > 0$   
 $\therefore f(x)$  increasing for all  $x$

ii) Range  $\{y: 0 < y < 1\}$   
 iii) Let  $y = 1 - \frac{1}{1+e^x}$   
 $\frac{1}{1+e^x} = 1-y$   
 $1+e^x = \frac{1}{1-y}$   
 $e^x = \frac{1}{1-y} - 1 = \frac{y}{1-y}$   
 $f'(x) = \ln \left( \frac{x}{1-x} \right)$



b) i)  $\frac{d}{dx} \left( \frac{v^2}{2} \right) = 4x - 4$   
 $\frac{dv^2}{dx} = 2v^2 - 4x + 4$   
 $\therefore \frac{v^2}{2} = 2x^2 - 4x + k$   
 When  $x=6, v^2=64$   
 $\therefore 32 = 72 - 24 + k \Rightarrow k = -16$   
 $\therefore v^2 = 4x^2 - 8x - 32$   
 ii)  $v^2 = 4(x^2 - 2x - 8)$   
 $= 4(x-4)(x+2)$   
 $v^2 \geq 0 \therefore -2 \leq x \leq 4$

iii) Particle moving to left from  $x=6$   
 Stops at  $x=4$  and immediately moves off to right, accelerating  
 $\therefore$  velocity  $> 0$

Equating coefficients:

$$\begin{cases} R \sin \alpha = 2 \\ R \cos \alpha = 1 \end{cases} \quad R = \sqrt{5} \quad \tan \alpha = 2$$

$$x = \sqrt{5} \sin(3t + \tan^{-1}(2))$$

$$i) \quad \dot{x} = 3\sqrt{5} \cos(3t + \tan^{-1}(2))$$

$$\ddot{x} = -9\sqrt{5} \sin(3t + \tan^{-1}(2))$$

$$\ddot{x} = -9x$$

which is of the form  $\ddot{x} = -\omega^2 x$  which defines S.H.M.

$$iii) \quad \text{Period} = \frac{2\pi}{\omega} = \frac{2\pi}{3} \text{ secs.}$$

iv) When  $x = 1$ ,

$$\sin(3t + \tan^{-1}(2)) = \frac{1}{\sqrt{5}}$$

$$3t + \tan^{-1}(2) = n\pi + (-1)^n \sin^{-1}\left(\frac{1}{\sqrt{5}}\right)$$

$n=1$  gives 1st +ve sol:

$$3t = \pi - \sin^{-1}\left(\frac{1}{\sqrt{5}}\right) - \tan^{-1}(2)$$

$$t = \frac{1}{3} \left( \pi - \sin^{-1}\left(\frac{1}{\sqrt{5}}\right) - \tan^{-1}(2) \right)$$

$$= 0.5 \text{ secs (to 10)}$$

b) i) After 1 withdrawal

$$\$ (20000 \times 1.005 - 50) = \$20050$$

ii) Let  $P = \$20000$ ,  $D = \$50$ ,  $r = 0.005$

$$\text{After 1 month } P(1+r) - D$$

$$\dots 2 \dots P(1+r)^2 - D(1+r) - D$$

$$\dots n \dots P(1+r)^n - D(1+r)^{n-1} - D(1+r)^{n-2} - \dots - D$$

$$= \frac{P(1+r)^n - D((1+r)^n - 1)}{r}$$

$$= 20000(1.005)^n - 10000(1.005)^n + 10000$$

$$= 10000(1.005)^n + 10000$$

$$10000 \times (1.005)^n \geq 40000$$

$$(1.005)^n \geq 4$$

$$n \log(1.005) \geq \log 4$$

$$n \geq \frac{\log 4}{\log 1.005}$$

$$\therefore n = 278 \text{ (months)}$$

$$6) a) r^{\text{th}} \text{ term is } {}^9C_r \left(\frac{2x}{3}\right)^r \left(\frac{-3}{2x}\right)^{9-r}$$

$$= {}^9C_r \left(\frac{2}{3}\right)^r \left(\frac{-3}{2}\right)^{9-r} x^{3r-9}$$

$$\text{Required term has } 3r-9=0 \quad r=3$$

$$\therefore \text{Coefficient is } {}^9C_3 (-1)^6 \left(\frac{3}{2}\right)^3$$

$$= {}^9C_3 \left(\frac{3}{2}\right)^3 = \frac{567}{2} = 283\frac{1}{2}$$

b) i) Intercept when  $x$  and  $y$  equal simultaneously

$$240t \cos \theta = 3600$$

$$t \cos \theta = 15$$

$$t = 15 \text{ sec}$$

$$2000 + 240t \sin \theta - \frac{gt^2}{2} = 3200 - \frac{gt^2}{2}$$

$$240t \sin \theta = 1200$$

$$t \sin \theta = 5$$

Solving  $\tan \theta = \frac{1}{3}$

$$\theta = \tan^{-1}\left(\frac{1}{3}\right) = 18^\circ 26'$$


At time  $t = 15 \text{ sec}$

$$= 5\sqrt{10} \text{ sec. } (= 15.81 \text{ s})$$

ii) Putting  $t = 5\sqrt{10}$  and  $g = 10$  into last equation

$$\text{height} = 3200 - \frac{10 \cdot 250}{2}$$

$$= 1950 \text{ m}$$



WPQRST defined in diagram  
 $\Delta PSQ \parallel \Delta PTR$  (Equiangular)

$$\frac{PR}{PQ} = \frac{RT}{QS} = 2$$

$$\text{But } PQ = h \cot \beta \quad \therefore PR = 2h \cot \beta$$

$$QR = SW = h \cot \alpha$$

Apply cosine rule to  $\Delta PQR$

$$\cos \theta = \frac{PQ^2 + PR^2 - QR^2}{2 \cdot PQ \cdot PR}$$

$$= \frac{h^2 \cot^2 \beta + 4h^2 \cot^2 \beta - h^2 \cot^2 \alpha}{2 \cdot h \cot \beta \cdot 2h \cot \beta}$$

$$\cos \theta = \frac{5 \cot^2 \beta - \cot^2 \alpha}{4 \cot^2 \beta}$$

7) a) i) Ways of choosing team of 4 =  ${}^{13}C_4 (= 715)$

Ways of choosing 3 boys and 1 girl =  ${}^8C_3 {}^5C_1 = 280$


$$\therefore \text{Probability} = \frac{{}^8C_3 {}^5C_1}{{}^{13}C_4} (= 0.39)$$

ii) Ways of choosing 4 girls =  ${}^8C_4 = 70$

Prob all girls =  $\frac{{}^8C_4}{{}^{13}C_4}$

Prob 3 or 4 girls

$$= \frac{{}^8C_4 + {}^8C_3 {}^5C_1}{{}^{13}C_4} (= 0.49)$$



i) Consider cross section of trough.  
 $OA = r \quad \therefore AB = \sqrt{r^2 - h^2}$  (Pythag)

$$\therefore \text{Surface area} = l \times 2AB = 2l\sqrt{r^2 - h^2}$$

ii) Let  $\angle AOB = \theta (= \cos^{-1}(\frac{h}{r}))$

Area of shaded segment is

$$\frac{r^2}{2} (2\theta - \sin 2\theta)$$

$$= \frac{r^2}{2} (\theta - \sin \theta \cos \theta)$$

$$= \frac{r^2}{2} \left( \cos^{-1}\left(\frac{h}{r}\right) - \frac{\sqrt{r^2 - h^2}}{r} \cdot \frac{h}{r} \right)$$

$$\therefore \text{Vol} = l \left( r^2 \cos^{-1}\left(\frac{h}{r}\right) - h\sqrt{r^2 - h^2} \right)$$

$$iii) \quad \frac{dV}{dt} = l \left( \frac{-\frac{h}{r}}{\sqrt{1 - \frac{h^2}{r^2}}} - \sqrt{r^2 - h^2} \right) \frac{dh}{dt}$$

$$+ \frac{h^2}{\sqrt{r^2 - h^2}} \frac{dh}{dt}$$

$$= \left( -\frac{r^2 - (r^2 - h^2) + h^2}{\sqrt{r^2 - h^2}} \right) l \frac{dh}{dt}$$

$$= -\frac{2l(r^2 - h^2) \frac{dh}{dt}}{\sqrt{r^2 - h^2}}$$

$$= -2l\sqrt{r^2 - h^2} \frac{dh}{dt} = -A \frac{dh}{dt}$$

$$iv) \quad -\frac{dV}{dt} < A \quad \therefore \frac{dV}{dt} = kA$$

where  $k$  is constant of proportionality

$$\therefore kA = A \frac{dh}{dt} \Rightarrow \frac{dh}{dt} = k$$

$\therefore h$  increasing at constant rate (ie. water falls at constant rate)