



SYDNEY BOYS HIGH SCHOOL
MOORE PARK, SURRY HILLS

2000

**TRIAL HIGHER SCHOOL
CERTIFICATE EXAMINATION**

Mathematics Extension 1


Examiner: B. Dowdell

- (a) State the domain and range of $4\sin^{-1} 3x$ 2
- (b) Solve for x : $(x - 2)^2 \leq 4$ 2
- (c) Differentiate: 4
- (i) $x \cos^{-1} 2x$
- (ii) $\frac{1}{4 + x^2}$
- (d) Find x correct to 3 decimal places if $x^{\frac{3}{4}} = 10$ 2
- (e) The point $P(11, 7)$ divides AB externally in the ratio 3:1. If B is $(6, 5)$, find the coordinates of A . 2

Question 2: START A NEW BOOKLET

Marks

(a)



2

The angle between the lines $y = mx$ and $y = \frac{1}{2}x$ is 135° . Find the exact value of m .

(b) Using $u = \sqrt{x}$ evaluate $\int_1^4 \frac{dx}{x + \sqrt{x}}$

2

(c) Write down the exact value of $\cos^{-1}(\cos \frac{4\pi}{3})$

2

(d) Find a primitive of

4

(i) $\frac{2}{\sqrt{1-4x^2}}$

(ii) $\frac{x}{4+x^2}$

(e) Find the values of a for which $f(x) = e^{-ax}(x-a)$ is stationary at $x = \frac{5}{2}$.


2

Question 3: START A NEW BOOKLET

Marks

(a)

3



O is the centre of the circle, XA is a tangent.

$$XY = 3 \text{ and } XA = 5$$

Calculate the size of $\angle AXY$ correct to the nearest minute.

- (b) (i) Sketch the graphs of $y = e^x$ and $y = \cos x$ on the same diagram for $0 \leq x \leq \frac{\pi}{2}$, clearly showing any points of intersection.

4

Shade the area enclosed by the two curves and the line $x = \frac{\pi}{2}$.

- (ii) Calculate the volume of the solid formed when this area is rotated about the x axis.

- (c) (i) Prove that $\frac{d^2x}{dt^2} = \frac{d}{dx} \left(\frac{1}{2}v^2 \right)$.

5

- (ii) A particle moves in a straight line with velocity given by $v^2 = 36 - 4x^2$ where x is measured in metres and is the displacement from a fixed point O and t is the time measured in seconds.

(α) Show that the motion is simple harmonic

(β) Find the period and amplitude of the motion.

Question 4: START A NEW BOOKLET

- (a) When $P(x) = ax^3 + bx + c$ is divided by $x - 1$ the remainder is -4 . 3
When $P(x)$ is divided by $x^2 - 4$, the remainder is $-4x + 3$.
Find a, b and c .
- (b) Prove by induction that 4
$$1 + (1 + 2) + (1 + 2 + 3) + \dots + (1 + 2 + 3 + \dots + n) = \frac{n}{6}(n + 1)(n + 2)$$
for all positive integers n .
- (c) (i) Show that the point $A(6p, 3p^2)$ lies on the parabola $x^2 = 12y$. 5
(ii) The chord joining $A(6p, 3p^2)$ and $B(6q, 3q^2)$, when produced, passes through $C(8, 0)$. Show that $4(p + q) = 3pq$ and hence find the locus of M , the midpoint of AB .

Question 5: START A NEW BOOKLET

Marks

(a) Prove that $2 \tan^{-1} \theta = \tan^{-1} \left(\frac{2\theta}{1-\theta^2} \right)$ provided that $|\theta| < 1$.


2

(b) A balloon is being filled with helium at a constant rate of $30 \text{ cm}^3/\text{s}$. Find the rate at which the surface area is increasing when its diameter is 40 cm.

4

(c)

6



A projectile is fired from a point P , a metres above O with an initial velocity $V \text{ ms}^{-1}$ at an angle of elevation of θ . It is subject to a constant downward acceleration of $g \text{ ms}^{-2}$.

- (i) Find expressions for the horizontal (x) and vertical (y) displacements from P after t seconds.
- (ii) Show that the time taken to reach Q , a metres from O in a horizontal direction is given by $\frac{2V(\sin \theta + \cos \theta)}{g}$ seconds.
- (iii) Show that $a = \frac{V^2(\sin 2\theta + \cos 2\theta + 1)}{g}$ metres.

Question 6: START A NEW BOOKLET

-) Eight people attend a meeting. They are provided with two circular tables, one seating 3 people, the other 5 people. 4

- (i) How many seating arrangements are possible?
(ii) If the seating is done randomly, what is the probability that a particular couple are on different tables?

-) If $f(x) = u(x) - \ln(u(x) + 1)$ 4

(i) Show that $f'(x) = \frac{u(x) \cdot u'(x)}{1 + u(x)}$.

- (ii) Hence or otherwise evaluate

$$\int_0^{\frac{\pi}{2}} \frac{\cos x \cdot \sin x}{1 + \sin x} dx$$

-) A function $L(x)$ is defined by 4

$$L(x) = Pe^{\frac{x}{3}} + Qe^{-\frac{2x}{3}} \text{ where } P \text{ and } Q \text{ are constants.}$$

It is given that $L(0) = 30$ and $L'(0) = -14$.


- (i) Find the values of P and Q .
(ii) Find $L'(3)$ and explain why $L(x)$ must have a minimum for some value of x between 0 and 3.

Question 7: START A NEW BOOKLET

Mark

(a)

3




AB is any chord of a circle. AB is produced to P , and PT is a tangent. The bisector of $\angle APT$ meets TB at M and TA at N .

- (i) Copy the diagram into your answer booklet.
- (ii) Prove that $\triangle TMN$ is isosceles.

(b)

9

AOB is a sector of a circle, such that, when the radius is x cm, $\angle AOB = (\pi - x)$ radians and x varies from 0 to π .



- (i) Find the maximum value of the perimeter of sector AOB . Comment on the minimum value of the perimeter of the sector.
- (ii) If the area of **triangle** AOB is given by $t(x)$
 - (α) Show that $t(x) = \frac{x^2 \sin x}{2}$.
 - (β) Show that when $t(x)$ is a maximum, $2 \tan x = -x$.
 - (γ) By sketching $y = \tan x$ and a suitable line, show that a solution to the equation in (β) is close to $x = \frac{3\pi}{4}$.
 - (δ) Taking $\frac{3\pi}{4}$ as a first approximation, use Newton's method once to obtain a better approximation (leave your answer in terms of π).

END OF PAPER



SYDNEY BOYS HIGH SCHOOL
MOORE PARK, SURRY HILLS

2000

**TRIAL HIGHER SCHOOL
CERTIFICATE EXAMINATION**

Mathematics Extension 1

Sample Solutions

(Q1) (a) $y = 4 \sin^{-1} 3x$

D: $-1 \leq 3x \leq 1$ R: $-\frac{\pi}{2} \leq \frac{y}{4} \leq \frac{\pi}{2}$
 $-\frac{1}{3} \leq x \leq \frac{1}{3}$ $-2\pi \leq y \leq 2\pi$

(b) $(x-2)^2 \leq 4$

$\therefore -2 \leq x-2 \leq 2$

$\therefore 0 \leq x \leq 4$

(c) (i) $\frac{d(\cos^{-1} 2x)}{dx} = \cos^{-1} 2x - \frac{x \times 2}{\sqrt{1-4x^2}}$

$= \cos^{-1} x - \frac{2x}{\sqrt{1-4x^2}}$

(ii) $\frac{d\left(\frac{1}{4+x^2}\right)}{dx} = \frac{d(4+x^2)^{-1}}{dx}$

$= -(4+x^2)^{-2} \times 2x$

$= \frac{-2x}{(4+x^2)^2}$

(d) $x^{3/4} = 10$
 $\therefore x = 10^{4/3}$
 ≈ 21.544

e) $P\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}\right)$

$A(x_1, y_1)$ $B(6, 5)$ $P(11, 7)$ $\begin{matrix} m & n \\ 3 & -1 \end{matrix}$
 $x_2 \ y_2$

$\therefore 11 = \frac{3 \times 6 - x_1}{2}, \quad 7 = \frac{3 \times 5 - y_1}{2}$

$18 - x_1 = 22, \quad 15 - y_1 = 14$

$x_1 = -4, \quad y_1 = 1$

$A(-4, 1)$

(2) (a) the acute angle is 45°

$$\tan 45 = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| \quad m_2 = \frac{1}{2}$$

$$\therefore \tan 45^\circ = \left| \frac{m_1 - \frac{1}{2}}{1 + m_1 \cdot \frac{1}{2}} \right|$$

$$\therefore \left| \frac{m_1 - \frac{1}{2}}{1 + \frac{m_1}{2}} \right| = 1 \Rightarrow \left| \frac{m_1 - \frac{1}{2}}{2 + m_1} \right| = 2$$

$$\therefore \frac{m_1 - \frac{1}{2}}{2 + m_1} = 2, \quad \frac{m_1 - \frac{1}{2}}{2 + m_1} = -2$$

$$m_1 - \frac{1}{2} = 4 + 2m_1 \quad m_1 - \frac{1}{2} = -4 - 2m_1$$

$$m_1 = 4\frac{1}{2}, \quad m_1 = -3\frac{1}{2}$$

$$\therefore \boxed{m = -3\frac{1}{2}, 4\frac{1}{2}}$$

(b) $u = \sqrt{x} \Rightarrow x = u^2$

$$\therefore dx = 2u du$$

$$\int_1^4 \frac{dx}{x + \sqrt{x}}$$

$$x=1 \Rightarrow u=1$$

$$x=4 \Rightarrow u=2$$

$$= \int_1^2 \frac{2u du}{u^2 + u}$$

$$= \int_1^2 \frac{2}{u+2}$$

$$= \ln|u+2| \Big|_1^2$$

$$= \ln 4 - \ln 3$$

$$= \ln\left(\frac{4}{3}\right)$$

(c) $\cos^{-1}\left(\cos \frac{4\pi}{3}\right)$

$$= \cos^{-1}\left(-\frac{1}{2}\right)$$

$$= \frac{2\pi}{3}$$

(d) (i) $\int \frac{2 dx}{\sqrt{1-4x^2}} = \sin^{-1}(2x) + C$

(ii) $\int \frac{x}{4+x^2} dx$

$$= \frac{1}{2} \int \frac{2x}{4+x^2} dx$$

$$= \frac{1}{2} \ln(x^2+4) + C$$

(2) (e)

$$f(x) = e^{-ax}(x-a)$$

$$f'(x) = e^{-ax} + (x-a) \times -ae^{-ax}$$
$$= e^{-ax}(1 - a(x-a))$$

$$f'(\frac{5}{2}) = 0$$

$$e^{-ax} \neq 0 \quad \therefore 1 - a(\frac{5}{2} - a) = 0$$


$$\therefore 2 - 5a + 2a^2 = 0$$

$$\therefore 2a^2 - 5a + 2 = 0$$

$$(2a-1)(a-2) = 0$$

$$a = \frac{1}{2}, 2$$

(3) (a)



$$XZ \cdot XY = XA^2$$

$$\therefore 25 = 3 \times XZ$$

$$XZ = \frac{25}{3} = 8\frac{1}{3}$$

$$\therefore YZ = 5\frac{1}{3}$$


$$\therefore OA = 8/3$$

$$\text{Let } \alpha = \angle AXY$$

$$\tan \alpha = \frac{8/3}{5} = 8/15$$

$$\alpha = 28^\circ 04'$$

(i)
(b)



$$\text{Area} = \int_0^{\pi/2} (e^x - \cos x) dx$$

$$= [e^x - \sin x]_0^{\pi/2}$$

$$= (e^{\pi/2} - \sin \frac{\pi}{2}) - (e^0 - \sin 0)$$

$$= e^{\pi/2} - 1 - 1$$

$$= e^{\pi/2} - 2$$

3 (b) (ii)

$$V = \pi \int_0^{\pi/2} (e^{2x} - \cos^2 x) dx$$

$$= \pi \int_0^{\pi/2} (e^{2x} - \frac{1}{2} - \frac{1}{2} \cos 2x) dx$$

$$= \pi \left[\frac{1}{2} e^{2x} - \frac{1}{2} x - \frac{1}{4} \sin 2x \right]_0^{\pi/2}$$

$$= \pi \left[\left(\frac{1}{2} e^{\pi} - \frac{\pi}{4} \right) - \left(\frac{1}{2} \right) \right]$$

$$= \frac{\pi}{2} \left(e^{\pi} - \frac{\pi}{2} - 1 \right)$$

$$\left[\cos^2 x = \frac{1}{2} (1 + \cos 2x) \right]$$

(c) (i) R.H.S = $d \left(\frac{1}{2} v^2 \right)$
 $\frac{dv}{dx}$

$$= d \left(\frac{1}{2} v^2 \right) \times \frac{dv}{dx}$$

$$= v \frac{dv}{dx} = \frac{dx}{dt} \cdot \frac{dv}{dx}$$

$$= \frac{dv}{dt}$$

$$= a$$

$$= \ddot{x}$$

$$= \text{L.H.S}$$

(ii) $v^2 = 36 - 4x^2$

$$\frac{1}{2} v^2 = 18 - x^2 \Rightarrow a = d \left(\frac{1}{2} v^2 \right) / dx$$

(\alpha) $a = -2x$

This is one of the defining equations for SHM, centred at $x=0$

(\beta) $n^2 = 2 \Rightarrow n = \sqrt{2}$

$$T = \frac{2\pi}{n} = \frac{2\pi}{\sqrt{2}} = \sqrt{2}\pi$$

$$v=0 \Rightarrow x^2 = 18$$

$$x = \pm 3\sqrt{2}$$

$$\therefore \text{Amplitude} = 3\sqrt{2} \text{ m}$$

$$(4) \quad (a) \quad P(x) = ax^3 + bx + c$$

$$P(1) = -4 \quad \Rightarrow \quad a + b + c = -4 \quad -\textcircled{1}$$

$$P(x) = (x^2 - 4)Q(x) + (-4x + 3)$$

$$\therefore P(2) = -5 \quad \Rightarrow \quad 8a + 2b + c = -5 \quad -\textcircled{2}$$

$$P(-2) = 11 \quad \Rightarrow \quad -8a - 2b + c = 11 \quad -\textcircled{3}$$

$$\textcircled{2} + \textcircled{3}: \quad 2c = 6$$

$$c = 3$$

$$\textcircled{1} \Rightarrow a + b = -7 \quad -\textcircled{4}$$

$$\textcircled{2} \Rightarrow 8a + 2b = -8 \quad -\textcircled{5}$$

$$4a + b = -4 \quad -\textcircled{6}$$

$$\textcircled{6} - \textcircled{4} \quad 3a = 3$$

$$a = 1$$

$$\text{sub into } \textcircled{4} \quad b = -8$$

$$\therefore a = 1, b = -8, c = 3$$

$$(b) \quad 1 + (1+2) + (1+2+3) + \dots + (1+2+3+\dots+n) = \frac{n}{6}(n+1)(n+2), \quad n > 0$$

Using the sum of an arithmetic series

$$e. \quad 1 + (1+2) + \dots + \frac{n(n+1)}{2} = \frac{n}{6}(n+1)(n+2), \quad n > 0 \quad -\textcircled{*}$$

$$\text{Test } n=1: \quad \text{LHS} = 1$$

$$\text{RHS} = \frac{1}{6}(2)(3) = 1$$

\therefore true for $n=1$

Assume $\textcircled{*}$ is true for some integer $n=k$.

$$e. \quad 1 + (1+2) + \dots + \frac{k(k+1)}{2} = \frac{k}{6}(k+1)(k+2)$$

We need to prove $\textcircled{*}$ is true for the integer $n=k+1$

$$i.e. \quad 1 + (1+2) + \dots + \frac{(k+1)(k+2)}{2} = \frac{(k+1)(k+2)(k+3)}{6}$$

4(b)

$$\text{LHS} = 1 + (1+2) + \dots + \frac{k(k+1)}{2} + \frac{(k+1)(k+2)}{2}$$

$$= \frac{k(k+1)(k+2)}{6} + \frac{(k+1)(k+2)}{2}$$

$$= (k+1)(k+2) \left[\frac{k}{6} + \frac{1}{2} \right]$$

$$= (k+1)(k+2) \frac{(k+3)}{6}$$

$$= \frac{1}{6} (k+1)(k+2)(k+3)$$

$$= \text{RHS}$$

\therefore Since the statement is true for $n=k+1$ when the statement is true for $n=k$. By the principle of mathematical induction


$$1 + (1+2) + \dots + (1+2+\dots+n) = \frac{n(n+1)(n+2)}{2}, \quad n > 0$$

(c) (i) $A(6p, 3p^2)$

$$\text{LHS} = x^2 = 36p^2$$

$$\text{RHS} = 12y = 12(3p^2) = 36p^2$$

$\therefore A$ lies on $x^2 = 12y$



$$x_1, y_1 \quad x_2, y_2$$

$$A(6p, 3p^2) \quad B(6q, 3q^2)$$

$$m_{AB} = \frac{3q^2 - 3p^2}{6q - 6p} = \frac{3(q-p)(q+p)}{6(q-p)}$$

$$= \frac{q+p}{2}$$

$$\therefore y - 3q^2 = \frac{q+p}{2}(x - 6q)$$

$$2y - 6q^2 = (q+p)x - 6q(q+p)$$

$$\therefore 2y = (q+p)x - 6qp \quad \text{--- (1)}$$

4 (c) (ii) $C(8,0)$ lies on (1)

$$\text{i.e. } 0 = (q+p)8 - 6qp$$

$$\therefore 6qp = 8(q+p) \Rightarrow 3pq = 4(p+q) \quad - (*)$$

$$\text{Midpoint AB } \left(\frac{6p+6q}{2}, \frac{3p^2+3q^2}{2} \right)$$

$$x = 3(p+q)$$

$$y = \frac{3}{2}(p^2+q^2)$$

$$= \frac{3}{2}[(p+q)^2 - 2pq]$$

$$= \frac{3}{2}(p+q)^2 - 3pq$$

$$= \frac{3}{2}(p+q)^2 - 4(p+q) \quad \text{from } (*)$$

$$= \frac{3}{2}\left[\frac{x}{3}\right]^2 - 4\left[\frac{x}{3}\right]$$

$$= \frac{x^2}{6} - \frac{4x}{3}$$

$$\therefore \text{Locus of M is } y = \frac{x^2}{6} - \frac{4x}{3}$$

Question 5(a)

$$2 \tan^{-1} \theta = \tan^{-1} \left(\frac{2\theta}{1-\theta^2} \right)$$

$|\theta| < 1$

$$\begin{aligned} \tan(2 \tan^{-1} \theta) &= \frac{2 \tan(\tan^{-1} \theta)}{1 - \tan^2(\tan^{-1} \theta)} \\ &= \frac{2\theta}{1-\theta^2} \end{aligned}$$

$$\therefore 2 \tan^{-1} \theta = \tan^{-1} \left(\frac{2\theta}{1-\theta^2} \right)$$

Now if $|\theta| > 1$

$2 \tan^{-1} \theta > \frac{\pi}{2}$ if $\theta > 1$
and $2 \tan^{-1} \theta < -\frac{\pi}{2}$ if $\theta < -1$

But $-\frac{\pi}{2} < \tan^{-1} x < \frac{\pi}{2}$

So R.H.S. has

$$-\frac{\pi}{2} < \tan^{-1} \left(\frac{2\theta}{1-\theta^2} \right) < \frac{\pi}{2}$$

So no valid solution

$$c) \frac{dv}{dt} = 30 \quad (v = \frac{4}{3} \pi r^3)$$

$$\frac{dv}{dt} = \frac{dv}{dr} \times \frac{dr}{dt} \quad \text{--- (1)}$$

$$\therefore 30 = 4\pi r^2 \times \frac{dr}{dt}$$

$$\Rightarrow \frac{dr}{dt} = \frac{15}{2\pi r^2} \quad \text{--- (2)}$$

$$s = 4\pi r^2$$


$$\frac{ds}{dt} = 8\pi r \frac{dr}{dt} \quad \text{--- (3)}$$

Subst (2) into (3)

$$= \cancel{8\pi r} \times \frac{15}{\cancel{2\pi r^2}}$$

$$\frac{ds}{dt} = \frac{60}{r}$$

$$\text{When } r = 20, \frac{ds}{dt} = 3$$



$$\ddot{x} = 0, \quad \dot{x} = v \cos \theta$$

$$x = (v \cos \theta) t \quad \text{--- (1)}$$

$$\ddot{y} = -g, \quad \dot{y} = (v \sin \theta) - gt$$

$$y = (v \sin \theta) t - \frac{gt^2}{2} + a \quad \text{--- (2)}$$

When $x = a, y = 0$

When $x = a$

$$t = \frac{a}{v \cos \theta} \quad \text{--- (3)}$$

and

When $t = \frac{a}{v \cos \theta}, y = 0$

Subst. (3) into (2)

We have

$$0 = v \sin \theta \left(\frac{a}{v \cos \theta} \right) - \frac{g}{2} \left(\frac{a}{v \cos \theta} \right)^2 + a$$

divide each term by a
and rearrange.

$$0 = \tan \theta - \frac{gt}{2v \cos \theta} + 1$$

$$\frac{gt}{2v \cos \theta} = \frac{\sin \theta + \cos \theta}{\cos \theta}$$

$$\therefore t = \frac{2v(\sin \theta + \cos \theta)}{g} \quad \text{--- (4)}$$

$a = (v \cos \theta) t$ — (5)
 In bst (4) into (5) we
 have


$$a = \frac{(v \cos \theta)(2v)(\sin \theta + \cos \theta)}{g}$$

$$= \frac{v^2(2\sin \theta \cos \theta + 2\cos^2 \theta)}{g}$$

$$= \frac{v^2(2\sin \theta \cos \theta + (2\cos^2 \theta - 1) + 1)}{g}$$

$$= \frac{v^2(\sin 2\theta + \cos 2\theta + 1)}{g}$$

Q nest in (6)
 (a)



(a) The 1st person has
 8 choices, the 2nd person
 (i) has 7 choices ...
 $\therefore \frac{8!}{5! 3!}$

(ii)



$$P(E) = \frac{\binom{2 \times 6!}{5! 3!}}{8!}$$

$$= \frac{2}{8 \times 7} = \frac{1}{28}$$

(b)

(i) $f(x) = u(x) - \ln[u(x)+1]$

$$f'(x) = u'(x) - \frac{u'(x)}{u(x)+1}$$

$$= u'(x) \left[1 - \frac{1}{u(x)+1} \right]$$

$$= u'(x) \left[\frac{u(x)+1-1}{u(x)+1} \right]$$

(ii)

$$\int_0^{\pi/2} \frac{\sin x \cos x}{1+\sin x} dx$$

$$= \left[\sin x - \ln(\sin x + 1) \right]_0^{\pi/2}$$

$$= (1 - \ln 2) - (0)$$

$$= 1 - \ln 2$$

(c)

$$L(0) = 30$$

$$\therefore 30 = p + q$$

$$L'(0) = -14$$

Now, $L'(x)$

$$= \frac{p}{3} e^{\frac{x}{3}} - \frac{2q}{3} e^{-\frac{2x}{3}}$$

$$\therefore -14 = \frac{p}{3} - \frac{2q}{3}$$

$$\therefore p - 2q = -42 \quad \text{--- (1)}$$

$$p + q = 30 \quad \text{--- (2)}$$

$$\Rightarrow p = 6, 6 + q = 30$$


$$\therefore q = 24$$

$$\therefore L'(0) = -14 < 0$$

$$\text{and } L'(3) = 2e^{-1} - 6e^{-2} > 0$$

$\therefore L(x_1)$ must be
 minimum for $0 < x_1 < 3$.

Question (7)



Let $\angle PAT = \beta$.

$\therefore \angle PTB = \beta$

(alternate segment theorem.)

Now


$\angle TNP = \alpha + \beta$

(ext. $\angle =$ sum of int. opp. \angle 's).

Similarly In $\triangle TPM$,

$\angle TMN = \alpha + \beta$.

$\therefore \triangle TMN$ is isosceles.



(b).

(i)

$$P = 2x + x(\pi - x)$$

$$\therefore P = (\pi + 2)x - x^2$$

$$\frac{dP}{dx} = \pi + 2 - 2x$$

$$\frac{dP}{dx} = 0, \quad 2x = \pi + 2$$

$$\therefore x = \frac{\pi + 2}{2}$$

$$\frac{d^2P}{dx^2} = -2 < 0$$

$\therefore P$ is max when $x = \frac{\pi + 2}{2}$

$$P_{\max} = \pi + 2 + \frac{\pi + 2}{2} \left(\frac{\pi}{2} - 1 \right)$$

$$= \pi + 2 + \left(\frac{\pi}{2} + 1 \right) \left(\frac{\pi}{2} - 1 \right)$$

$$= \pi + 2 + \frac{\pi^2}{4} - 1$$

$$= \frac{\pi^2}{4} + \pi + 1$$

$$= \frac{\pi^2 + 4\pi + 4}{4}$$

$$t(x) = \frac{x^2}{2} \sin(\pi - x)$$

$$\sin(\pi - x) = \sin x$$

$$\therefore t(x) = \frac{x^2 \sin x}{2}$$


$$\frac{dt(x)}{dx} = x \sin x + \frac{x^2}{2} \cos x$$

$$\frac{dt(x)}{dx} = 0, \quad x \left(\sin x + \frac{x \cos x}{2} \right) = 0$$

$$\therefore \sin x = -\frac{x \cos x}{2}$$

$$\Rightarrow \tan x = -\frac{x}{2}$$

$$\therefore 2 \tan x = -x$$



$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$= \frac{3\pi}{4} - \frac{-2 + \frac{3\pi}{4}}{1 + \frac{3\pi}{4}}$$

$$=$$