



Sydney Girls High School

2007

TRIAL HIGHER SCHOOL  
CERTIFICATE EXAMINATION

# Mathematics

## Extension 1

This is a trial paper ONLY. It does not necessarily reflect the format or the contents of the 2007 HSC Examination Paper in this subject.

### General Instructions

- Reading Time - 5 mins
- Working time - 2 hours
- Attempt ALL questions
- ALL questions are of equal value
- All necessary working should be shown in every question. Marks may be deducted for careless or badly arranged work.
- Standard integrals are supplied
- Board-approved calculators may be used.
- Diagrams are not to scale
- Each question attempted should be started on a new sheet. Write on one side of the paper only.

--	--	--	--	--	--	--	--	--	--

Candidate Number

**Question 1 (12 marks)**

(a) Find  $\int \cos^2(2x) dx$  2

(b) Using the substitution  $u = e^x$  find  $\int \frac{e^x}{1+e^{2x}} dx$  3

c) Evaluate  $\lim_{x \rightarrow 0} \frac{\sin \frac{x}{2}}{4x}$  2

(d) The point  $M(-3, 8)$  divides the interval  $AB$  externally in the ratio  $k:1$   
If  $A = (6, -4)$  and  $B = (0, 4)$ , Find the value of  $k$ . 3

(e) Prove the identity 2


$$\frac{2 \tan A}{1 + \tan^2 A} = \sin 2A$$

**Question 2 (12 marks)**

- (a) Consider the function  $f(x) = 3 \sin^{-1}\left(\frac{x}{2}\right)$  4
- (i) Evaluate  $f(2)$
- (ii) Draw the graph of  $y = f(x)$
- (iii) State the Domain and Range of  $y = f(x)$
- (b) One root of the polynomial equation  $x^3 + 6x^2 - x - 30 = 0$  is equal to the sum of the other two roots. Find all three roots. 3
- (c) Use Newton's Method to find a second approximation to the positive root of the equation  $x = 2 \sin x$  taking  $x = 1.7$  as the first approximation. Give answer in radians correct to 1 decimal place. 3
- (d) Solve the inequality  $\frac{2}{x-1} < 1$  2

## Question 3 (12 marks)

- (a) 3
- (i) Find the point of intersection of the line  $y = x$  with the curve  $y = x^3$  in the first quadrant.
- (ii) Then find the size of the acute angle between the line and the curve at this point to the nearest degree.



- (i) Copy this diagram onto your answer page.
- (ii) Prove that  $\triangle PRT$  and  $\triangle QPT$  are similar.
- (iii) Hence prove that  $PT^2 = QT \times RT$
- (c) Let  $T$  be the temperature inside a room at time  $t$  hours and let  $A$  be the constant outside air temperature. Newton's Law of Cooling states that the rate of change of the temperature  $T$  is proportional to  $(T - A)$ .
- (i) Show that  $T = A + Ce^{kt}$  where  $C$  and  $k$  are constants satisfies Newton's Law of Cooling. 1
- $$\frac{dT}{dt} = k(T - A)$$
- (ii) The outside air temperature is  $5^{\circ}\text{C}$  when a system failure causes the inside room temperature to drop from  $20^{\circ}\text{C}$  to  $17^{\circ}\text{C}$  in half an hour. After how many hours is the inside room temperature equal to  $10^{\circ}\text{C}$ ? Give answer correct to 1 decimal place. 3

## Question 4 (12 marks)

- (a) The acceleration of a particle moving in a straight line is given by  $\ddot{x} = 2x - 3$  where  $x$  is the displacement, in metres, from the origin  $O$  and  $t$  is the time in seconds. Initially the particle is at rest at  $x = 4$ . 4
- (i) If the velocity of the particle is  $V \text{ ms}^{-1}$  show that  $V^2 = 2(x^2 - 3x - 4)$
- (ii) Show that the particle does **not** pass through the origin.
- (iii) Find the position of the particle when  $V = 10 \text{ ms}^{-1}$
- (b)
- (i) Find the inverse function  $f^{-1}(x)$  in terms of  $x$  for  $f(x) = 2x - x^2$  over the restricted domain  $x \geq 1$ . Write the Domain and Range of the inverse function. 4
- (ii) Find the point common to both  $f(x)$  and  $f^{-1}(x)$  in this domain.
- (c) From the top of a mountain 200 metres above ground an observer sights two landmarks A and B. Point A has a bearing of  $300^\circ \text{T}$  at an angle of depression of  $10^\circ$ . Point B has a bearing of  $040^\circ \text{T}$  at an angle of depression of  $15^\circ$ . Calculate the distance from A to B given that both points are at ground level. (to the nearest metre). 4

**Question 5 (12 marks)**

(a) 3


- (i) Express  $\sqrt{3} \sin \Theta - \cos \Theta$  in the form  $A \sin (\Theta - \alpha)$  where  $\alpha$  is in radians and  $A > 0$
- (ii) Hence, or otherwise find all angles  $\Theta$ , where  $0 \leq \Theta \leq 2\pi$  for which  $\sqrt{3} \sin \Theta - \cos \Theta = 1$

(b) Consider the parabola  $x^2 = 4ay$  where  $a > 0$ . 5

The tangents at  $P(2ap, ap^2)$  and  $Q(2aq, aq^2)$  intersect at the point  $T$ .  
Let  $S(0, a)$  be the focus of the parabola.

- (i) Find the coordinates of  $T$ . (You may assume the equation of the tangent at  $P$  is  $px - y - ap^2 = 0$ )
- (ii) Show that  $SP = ap^2 + a$
- (iii) Now  $P$  and  $Q$  move along the parabola in such a way that  $SP + SQ = 4a$   
Find the locus of  $T$  under this condition.

(c) 4



$QR$  is a tangent touching the circle at  $T$

- (i) Copy this diagram onto your answer page.
- (ii) Prove that  $LMNP$  is a cyclic quadrilateral

**Question 6 (12 marks)**

(a) Prove by mathematical induction that  $n^3 + 2n$  is divisible by 3 for all positive integers  $n$ . 4

(b) 4

(i) Find the exact area bounded by the curve  $y = \frac{x-1}{\sqrt{x+1}}$ , the  $x$  axis and the lines  $x = 3$  and  $x = 8$ .  
Use the substitution  $u^2 = x + 1$

(ii) Now find the volume of the solid of revolution formed by rotating this area about the  $x$  axis. Give answer correct to 1 decimal place.

(c) 4



Car P is North of an intersection and travelling towards O

Car Q is moving away from the intersection eastwards at 60 km / hour

The distance between the two cars at any given time is 10 km.

Find the rate in km per hour at which car P is moving when car Q is 8 km away from the intersection.

**Question 7 (12 marks)**

- (a) A particle's displacement is given by  $x = 2 \cos(t + \frac{\pi}{4})$  metres at time  $t$  seconds 5
- (i) Show that acceleration is proportional to the displacement and hence describe its motion.
  - (ii) Find the initial position
  - (iii) Find the period of the motion
  - (iv) Find the maximum displacement
  - (v) Find the particle's position after  $\frac{\pi}{2}$  secs.
- (b) A sky rocket is fired vertically into the air. At a height of 28 metres it explodes and is projected at an angle of  $60^\circ$  to the horizontal with a velocity of  $30 \text{ ms}^{-1}$ . Take  $g = 10 \text{ ms}^{-2}$  7
- (i) How long from the time of the explosion will it take to fall back to the ground?
  - (ii) How far from its launching site will it land?
  - (iii) At what velocity will it strike the ground? To nearest whole number.
  - (iv) What acute angle will it make with the ground on impact? To nearest degree.

<b>End of Exam</b>
--------------------



## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE :  $\ln x = \log_e x, \quad x > 0$

$$\frac{81}{a} \int \cos^2(2x) dx$$

$$\cos 4x = 2 \cos^2(2x) - 1$$

$$\therefore \cos^2(2x) = \frac{1}{2} + \frac{\cos 4x}{2}$$

$$\int \cos^2(2x) dx = \int \frac{1}{2} + \frac{1}{2} \cos 4x dx$$

$$= \frac{1}{2}x + \frac{1}{8} \sin 4x + C \quad (2)$$

$$\frac{c}{\lim_{x \rightarrow 0} \frac{\sin \frac{x}{2}}{4x}}$$

$$= \frac{1}{8} \lim_{x \rightarrow 0} \frac{\sin \frac{x}{2}}{\frac{x}{2}}$$

$$= \frac{1}{8} \quad (2)$$

$$\frac{d}{x} = \frac{k_1 x_2 + k_2 x_1}{k_1 + k_2}$$

$$-3 = \frac{k \times 0 + 1 \times 6}{k + 1}$$

$$-3k - 3 = 6$$

$$-3k = 9$$

$$\therefore k = -3$$

$$y = \frac{k_1 y_2 + k_2 y_1}{k_1 + k_2}$$

$$8 = \frac{k \times 4 + 1 \times (-4)}{k + 1}$$

$$8k + 8 = 4k - 4$$

$$4k = -12$$

$$\therefore k = -3 \quad (3)$$

$$\frac{b}{\int \frac{e^x}{1+e^{2x}} dx}$$

$$u = e^x$$

$$\frac{du}{dx} = e^x$$

$$\therefore du = e^x dx$$

$$\int \frac{e^x dx}{1+e^{2x}} = \int \frac{du}{1+u^2}$$

$$= \tan^{-1} u + C \quad (3)$$

$$= \tan^{-1}(e^x) + C$$

$\frac{e}{\text{Prove}}$

$$\frac{2 \tan A}{1 + \tan^2 A} = \sin 2A$$

$$\text{L.H.S} = \frac{2 \tan A}{1 + \tan^2 A}$$

$$= \frac{2 \tan A}{\sec^2 A}$$

$$= \frac{2 \sin A}{\cos A} \cdot \cos^2 A$$

$$= 2 \sin A \cos A$$


$$= \sin 2A$$

$$= \text{R.H.S.} \quad (2)$$

Q2

(a)  $f(x) = 3 \sin^{-1}\left(\frac{x}{2}\right)$

$$\begin{aligned} \text{i } f(2) &= 3 \sin^{-1}\left(\frac{2}{2}\right) \\ &= 3 \times \frac{\pi}{2} \\ &= \frac{3\pi}{2} \end{aligned} \quad (1)$$



iii Dom  $-2 \leq x \leq 2$  (1)

Range  $-\frac{3\pi}{2} \leq y \leq \frac{3\pi}{2}$  (1)

(c)  $x = 2 \sin x$

$\therefore x - 2 \sin x = 0$

Let  $f(x) = x - 2 \sin x$

$f'(x) = 1 - 2 \cos x$

Let  $x_1 = 1.7$

$$\begin{aligned} x_2 &= x_1 - \frac{f(x_1)}{f'(x_1)} \\ &= 1.7 - \frac{[1.7 - 2 \sin 1.7]}{[1 - 2 \cos 1.7]} \\ &= 1.9 \end{aligned} \quad (3)$$

(d)  $\frac{2}{x-1} < 1$


Let  $x-1 = 0$

$\therefore x = 1$  (not c.v)

Let  $\frac{2}{x-1} = 1$

$2 = x-1$

$3 = x$  (2nd c.v)



Test  $x = 0$ ,  $\frac{2}{-1} < 1$  True

Test  $x = 2$ ,  $\frac{2}{1} \nless 1$

Test  $x = 4$ ,  $\frac{2}{3} < 1$  True

$\therefore$  Ans  $x < 1, x > 3$

(2)

(b)  $x^3 + 6x^2 - x - 30 = 0$

Roots =  $\alpha, \beta, \gamma$

$\alpha = \beta + \gamma$  (given)

Sum of roots

$= \alpha + \beta + \gamma = -6$

$\alpha + \alpha = -6$

$\therefore \alpha = -3$

Product in pairs

$\alpha\beta + \alpha\gamma + \beta\gamma = -1$

$-3\beta - 3\gamma + \beta\gamma = -1$

$-3(\beta + \gamma) + \beta\gamma = -1$

$-3(\alpha) + \beta\gamma = -1$

$9 + \beta(-3-\beta) = -1$

$9 - 3\beta - \beta^2 = -1$

$0 = \beta^2 + 3\beta - 10$

$(\beta + 5)(\beta - 2) = 0$

$\therefore \beta = -5$  or  $2$

$\alpha = \beta + \gamma$

$-3 = -5 + \gamma$

$\therefore \gamma = 2$

or  $-3 = 2 + \gamma$

$\gamma = -5$

(3)

$\therefore$  Roots are  $-3, -5, 2$

Q3

i  $y = x$  ,  $y = x^3$

$x^3 = x$

$x^3 - x = 0$

$x(x-1)(x+1) = 0$

$\therefore x = 0, x = 1, x = -1$

$\therefore$  In 1st Quad Intersection pt = (1,1) (1)

ii  $y = x^3$  ,  $\frac{dy}{dx} = 3x^2$


$x = 1$  ,  $\frac{dy}{dx} = 3 = m_1$

$y = x$  ,  $\frac{dy}{dx} = 1 = m_2$

$\tan \theta = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$

$= \frac{3 - 1}{1 + 3 \times 1} = \frac{2}{4} = \frac{1}{2}$

$\theta = 27^\circ$  (2)



ii Ans. Prove  $\Delta PRT \parallel \Delta QPT$

Proof. In  $\Delta PRT$  and  $\Delta QPT$

$\angle T$  is common

$\angle TPR = \angle Q$  (Angle in alt Seg.)

$\therefore \angle PRT = \angle QPT$  ( $\angle$  sum of  $\Delta$ )

$\therefore \Delta PRT \parallel \Delta QPT$  (equiangular) (2)

iii  $\frac{PT}{QT} = \frac{RT}{PT}$  (eq ratios sim  $\Delta$ s)

$\therefore PT^2 = QT \times RT$  (2)

Finally put  $T = 10$

$10 = 5 + 15e^{kt}$

$\log_e \left( \frac{5}{15} \right) = kt$

$\therefore t = 2.46 \approx 2.5$  hours. (3)

i  $\frac{dT}{dt} = k(T-A)$

Proposed solution is

$T = A + Ce^{kt}$

L.H.S =  $\frac{dT}{dt} = 0 + Ck e^{kt}$

R.H.S =  $k(T-A)$

$= k(Ce^{kt})$

$\therefore$  L.H.S = R.H.S (1)

ii  $T = A + Ce^{kt}$

$T = 5 + Ce^{kt}$

$t = 0, T = 20$

$20 = 5 + Ce^{k \times 0}$

$\therefore C = 15$

$T = 5 + 15e^{kt}$

$17 = 5 + 15e^{0.5k}$

$\log_e \left( \frac{12}{15} \right) = 0.5k$

$k = -0.446287$

Q4

$$a \text{ i } \ddot{x} = \frac{d}{dx} \left( \frac{1}{2} v^2 \right)$$

$$\frac{d}{dx} \left( \frac{1}{2} v^2 \right) = 2x - 3$$

$$\frac{1}{2} v^2 = \int 2x - 3 \, dx$$

$$= x^2 - 3x + c$$

$$t=0, x=4, v=0$$

$$0 = 16 - 12 + c$$

$$c = -4$$

$$\therefore \frac{1}{2} v^2 = x^2 - 3x - 4 \quad (2)$$

$$\therefore v^2 = 2(x^2 - 3x - 4)$$

ii At origin,  $x=0$

$$\therefore v^2 = -8 \text{ NO soln.}$$

$\therefore$  Particle does NOT pass through origin. (1)

iii  $v = 10$

$$100 = 2(x^2 - 3x - 4)$$

$$0 = x^2 - 3x - 54$$

$$(x-9)(x+6) = 0$$

$$x = 9 \text{ or } -6 \quad (1)$$

Since particle starts at

$x=4$  and can't reach

$x=-6$  (other side of origin)

$\therefore$   $x=9$  m when  $v=10$

$$b \text{ i } f: y = 2x - x^2$$

$$\left. \begin{array}{l} \text{Dom } x \geq 1 \\ \text{Range } y \leq 1 \end{array} \right\} \text{Restrict}$$

$$f^{-1}: x = 2y - y^2$$

$$y^2 - 2y = -x$$

$$y^2 - 2y + 1 = 1 - x$$

$$(y-1)^2 = 1-x$$

$$y-1 = \pm \sqrt{1-x}$$

$$y = 1 \pm \sqrt{1-x}$$

$$\therefore \text{Inv. } f^{-1} \text{ is } y = 1 + \sqrt{1-x}$$

$$f^{-1} \left. \begin{array}{l} \text{Dom } x \leq 1 \\ \text{Range } y \geq 1 \end{array} \right\} (3)$$

ii Common point solve

$$y = x \text{ with}$$

$$y = 2x - x^2$$


$$x = 2x - x^2$$

$$x^2 - x = 0$$

$$x(x-1) = 0$$




$$\therefore (1, 1) = \text{Common Pt.}$$

(1)



04 (c)

4




$$\tan 10^\circ = \frac{200}{x}$$

$$\tan 15^\circ = \frac{200}{y}$$

$$x = \frac{200}{\tan 10^\circ} = 1134.26$$

$$y = \frac{200}{\tan 15^\circ} = 746.41$$



Using Cosine Rule

$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$c^2 = 746.41^2 + 1134.26^2 - 2 \times 746.41 \times 1134.26 \times \cos 100^\circ$$

$$c = 1462.09$$

Ans AB = 1462 m

Q5

$$\begin{aligned} i \quad & \sqrt{3} \sin \theta - \cos \theta \\ & = A \sin(\theta - \alpha) \\ & = A \sin \theta \cos \alpha - A \cos \theta \sin \alpha \end{aligned}$$


$$\begin{aligned} A \cos \alpha &= \sqrt{3}, \quad A \sin \alpha = 1 \\ A^2 &= 4 \quad \therefore A = 2 \end{aligned}$$

$$\frac{\sin \alpha}{\cos \alpha} = \tan \alpha = \frac{1}{\sqrt{3}} \quad \begin{array}{c} 2 \\ \nearrow \\ 1 \end{array} \quad \left(\frac{1}{\sqrt{3}}\right)$$

$\therefore \alpha = 30^\circ = \pi/6$


$$\therefore A \sin(\theta - \alpha) = 2 \sin(\theta - \pi/6)$$

$$\begin{aligned} ii \quad & 2 \sin(\theta - \pi/6) = 1 \\ & \sin(\theta - \pi/6) = 1/2 \\ & \theta - \frac{\pi}{6} = \pi/6 \quad \text{or} \quad 5\pi/6 \\ & \therefore \theta = \pi/3 \quad \text{or} \quad \pi \end{aligned} \quad \left(\frac{1}{2}\right)$$



$$\begin{aligned} i \quad & px - y - ap^2 = 0 \quad (1) \\ & qx - y - aq^2 = 0 \quad (2) \\ & (p - q)x = a(p^2 - q^2) \quad (1) - (2) \\ & \therefore x = a(p + q) \\ & ap(p + q) - y - ap^2 = 0 \quad (2) \\ & \therefore y = apq \end{aligned}$$

$$\therefore T = [a(p + q), apq]$$



Aim. Prove LMNP is a cyclic quadrilateral.

Proof. Let  $\angle NTR = \alpha$

$$\angle NTR = \angle PNT = \alpha \quad (\text{alt } \angle\text{'s } PN \parallel AR)$$

$$\text{Also } \angle NTR = \angle TLM = \alpha \quad (\text{angle in alt seg.})$$

$$\angle PNM = 180 - \alpha \quad (\text{adj supp } \angle\text{'s})$$

$\therefore$  LMNP is cyclic quad since  $\angle L + \angle PNM = 180^\circ$   
(opp  $\angle\text{'s}$  supp.)

$$\begin{aligned} \underline{b} \quad ii \quad & SP^2 = (2ap - 0)^2 + (ap^2 - a)^2 \\ & = 4a^2p^2 + a^2p^4 - 2a^2p^2 + a^2 \\ & = a^2p^4 + 2a^2p^2 + a^2 \\ & = a^2(p^2 + 1)^2 \\ \therefore SP &= ap^2 + a \end{aligned}$$

(iii) (over)

Q5

Condition of locus is

b iii  $SP + SQ = 4a$

$$ap^2 + a + aq^2 + a = 4a$$

$$a(p^2 + q^2) = 2a$$

$$\therefore p^2 + q^2 = 2$$

$$x = a(p + q)$$

$$y = apq$$

$$(p + q)^2 = p^2 + 2pq + q^2$$

$$\frac{x^2}{a^2} = 2 + \frac{2y}{a}$$

$$x^2 = 2a^2 + 2ay$$

$$\therefore x^2 = 2a(y + a) \text{ is locus of } T$$

(2)



Q6  
(a) Prove  $n^3 + 2n$  is divisible by 3 for all positive integers  $n$ .

Step 1 Prove true for  $n=1$

$$1^3 + 2 \times 1 = 3 \text{ which is divisible by 3} \quad \therefore \text{True for } n=1$$

Step 2 Assume true for  $n=k$  (= integer)

$$k^3 + 2k = 3m \quad (m = \text{integer})$$


Step 3 Prove true for  $n=k+1$

$$\begin{aligned} & (k+1)^3 + 2(k+1) \\ &= k^3 + 3k^2 + 3k + 1 + 2k + 2 \\ &= (k^3 + 2k) + (3k^2 + 3k + 3) \\ &= 3m + 3(k^2 + k + 1) \end{aligned}$$

which is divisible by 3 since  $k^2 + k + 1 = \text{integer}$ .

$\therefore$  True for  $n=k+1$

Step 4 Since true for  $n=1$  and having assumed true for  $n=k$  and subsequently proven true for  $n=k+1$ , then result is true by Math. Induction for all positive integers  $n$ . (4)



$$\text{Area} = \int_3^8 \frac{x-1}{\sqrt{x+1}} dx$$

$$u^2 = x+1$$

$$u^2 - 1 = x$$

$$\frac{dx}{du} = 2u$$

$$\therefore dx = 2u du$$

Change Limits

$$x=3, u=2$$

$$x=8, u=3$$

$$\text{Area} = \int_2^3 \frac{u^2-2}{u} \cdot 2u du$$

$$= 2 \int_2^3 (u-2) du$$

$$= 2 \left[ \frac{u^2}{2} - 2u \right]_2^3$$

$$= 2 \left[ \left( \frac{27}{2} - 6 \right) - \left( \frac{8}{2} - 4 \right) \right]$$

$$= 8\frac{2}{3} \text{ units}^2 \quad (2)$$

b ii  $\text{Vol} = \pi \int_3^8 y^2 dx$

$$= \pi \int_3^8 \frac{(x-1)^2}{x+1} dx$$

$$u^2 = x+1$$

$$x-1 = u^2 - 2$$

$$(x-1)^2 = (u^2 - 2)^2$$

$$= u^4 - 4u^2 + 4$$

$$\text{Vol} = \pi \int_2^3 \frac{(u^4 - 4u^2 + 4) \cdot 2u du}{u^2}$$

$$= 2\pi \int_2^3 (u^3 - 4u + \frac{4}{u}) du$$

$$= 2\pi \left[ \frac{u^4}{4} - 2u^2 + 4 \log_e u \right]_2^3$$

$$= 2\pi \left[ \frac{81}{4} - 18 + 4 \log_e 3 - 4 + 8 - 4 \log_e 2 \right]$$


$$= 49.46$$

$$\therefore \text{units}^3$$

$$49.5$$

(2)

Q6



$$x^2 + y^2 = 100$$

$$y = \sqrt{100 - x^2}$$
$$= (100 - x^2)^{\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{1}{2} (-2x) (100 - x^2)^{-\frac{1}{2}}$$
$$= \frac{-x}{\sqrt{100 - x^2}}$$

$$\frac{dx}{dt} = +60$$

since moving  
Left to Right.

$$\frac{dy}{dt} = \frac{dy}{dx} \times \frac{dx}{dt}$$

$$= \frac{-x}{\sqrt{100 - x^2}} \times +60$$

Put  $x = +8$  (since left of 0)

$$\frac{dy}{dt} = \frac{8 \times -60}{\sqrt{100 - 64}}$$

$$= \frac{8 \times -60}{\sqrt{36}}$$

$$= -80 \text{ km/h}$$

$\therefore$  Car P is travelling at 80 km/h when  
Car Q is 8 km from the intersection.

4

Q7 (a)  $x = 2 \cos(t + \frac{\pi}{4})$

i  $\dot{x} = -2 \sin(t + \frac{\pi}{4})$

$\ddot{x} = -2 \cos(t + \frac{\pi}{4})$

$\therefore \ddot{x} = -x$

Thus acceleration is proportional to the displacement (x)

$\ddot{x} = -\omega^2 x$

$\therefore$  Motion is S.H.M. (1)

$\checkmark$   $t = \frac{\pi}{2}, x = 2 \cos(\frac{3\pi}{4}) = 2 \times \frac{-1}{\sqrt{2}} = -\sqrt{2} \text{ m}$  (1)

ii let  $t=0$

$x = 2 \cos(\frac{\pi}{4})$


$= 2 \times \frac{1}{\sqrt{2}}$

$= \sqrt{2} = \text{initial position.}$  (1)

iii  $\omega^2 = 1 \therefore \omega = 1$  (1)

$T = \frac{2\pi}{\omega} = \frac{2\pi}{1} = 2\pi \text{ secs.}$

iv Max displacement = a = 2 metres (1)



Horiz. Motion

$\dot{x} = 15$

$x = 15t$

Vertical Motion

$\ddot{y} = -g$

$\dot{y} = -gt + c$

$15\sqrt{3} = 0 + c$

$\dot{y} = -gt + 15\sqrt{3}$

$y = -\frac{gt^2}{2} + 15\sqrt{3} \cdot t + 28$

Data  $t=0, x=0, \dot{x} = 30 \times \frac{1}{2} = 15$

$t=0, y=28, \dot{y} = 30 \times \frac{\sqrt{3}}{2} = 15\sqrt{3}$

$g = 10$

i Put  $y=0$

$0 = -5t^2 + 15\sqrt{3}t + 28$

$5t^2 - 15\sqrt{3}t - 28 = 0$

$t = \frac{15\sqrt{3} \pm \sqrt{675 + 560}}{10}$

$= 6.1$

$= 6 \text{ secs.}$  (2)

ii  $x = 15 \times 6$

$= 90 \text{ m}$

(1)

(ii)  $\dot{y} = -10 \times 6 + 15\sqrt{3}$

$= -34$


$\dot{x} = 15$

$v^2 = 15^2 + (-34)^2$

$= 1381$

$v = 37 \text{ m s}^{-1}$

(2)



$\tan \alpha = \frac{15}{34}$

$\alpha = 23^\circ 48'$

$\therefore \theta = 90^\circ - 23^\circ 48' = 66^\circ$

(2)