

# Sydney Girls High School

2015

TRIAL HIGHER SCHOOL CERTIFICATE  
EXAMINATION

# Mathematics Extension 1

## General Instructions

- Reading Time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen  
Black pen is preferred
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In Questions 11 – 14, show relevant mathematical reasoning and/or calculations

Total marks – 70

**Section I** Pages 3 – 5

**10 Marks**

- Attempt Questions 1 – 10
- Answer on the Multiple Choice answer sheet provided
- Allow about 15 minutes for this section

**Section II** Pages 6 – 9

**60 Marks**

- Attempt Questions 11 – 14
- Answer on the blank paper provided
- Begin a new page for each question
- Allow about 1 hour and 45 minutes for this section

Name: .....

Teacher: .....

**THIS IS A TRIAL PAPER ONLY**

It does not necessarily reflect the format or the content of the 2015 HSC Examination Paper in this subject.

## Section I

10 marks

Attempt Questions 1–10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1-10.


---

1. A committee of six is to be formed from seven women and nine men. Find the number of committees possible if exactly two members of the committee are to be men.

- (A) 1260
- (B) 2646
- (C) 36036
- (D) 60480

2. In the diagram below  $\angle BAC = 34^\circ$  and  $\angle ADE = 85^\circ$ . What is the size of angle  $\angle ACB$ ?

- (A)  $51^\circ$
- (B)  $56^\circ$
- (C)  $61^\circ$
- (D)  $60^\circ$



3. Use the substitution  $u = e^x$  to determine which of the following is an expression for  $\int \frac{e^x}{1+e^{2x}} dx$ .

- (A)  $\tan^{-1}(e^x) + C$
- (B)  $\tan^{-1}(e^{2x}) + C$
- (C)  $\frac{-1}{2(1+e^x)^2} + C$
- (D)  $\frac{-e^x}{(1+e^x)^2} + C$

4. The radius of a spherical balloon is increasing at the rate of 2 cm/s. The rate at which the volume of the balloon is increasing when the radius is 10 cm is :
- (A)  $200\pi \text{ cm}^3/\text{s}$
- (B)  $400\pi \text{ cm}^3/\text{s}$
- (C)  $800\pi \text{ cm}^3/\text{s}$
- (D)  $100\pi \text{ cm}^3/\text{s}$
5. A stone is thrown vertically upwards with a speed of 21 m/s. How long is the stone in the air before it reaches its maximum height? (Assume acceleration due to gravity is  $10 \text{ m/s}^2$ .)
- (A) 4.2 s
- (B) 0.48 s
- (C) 0.95 s
- (D) 2.1 s
6. The polynomial equation  $f(x) = x^3 + x - 1$  has a root near  $x = 0.5$ . Using this as the initial approximation, determine another approximation (correct to four decimal places) to the root using one application of Newton's method.
- (A)  $x = 0.7141$
- (B)  $x = 0.7142$
- (C)  $x = 0.7143$
- (D)  $x = 0.7144$
7. If  $y = \sin^{-1}\left(\frac{a}{x}\right)$ , then  $\frac{dy}{dx} =$
- (A)  $\frac{-a}{\sqrt{x^2 - a^2}}$
- (B)  $\frac{a}{\sqrt{x^2 - a^2}}$
- (C)  $\frac{-a}{x\sqrt{x^2 - a^2}}$
- (D)  $\frac{a}{x\sqrt{x^2 - a^2}}$

8. What is the value of  $\sum_{k=1}^{20} {}^{20}C_k$  ?

- (A) 1 048 574
- (B) 1 048 575
- (C) 1 048 576
- (D) 1 048 577

9. Given that  $a$ ,  $b$  and  $c$  are the roots of the equation  $x^3 - 3x^2 + mx + 24 = 0$ , and that  $-a$  and  $-b$  are the roots of the equation  $x^2 + nx - 6 = 0$ , then the value of  $n$  is :

- (A) 1
- (B) -1
- (C) 7
- (D) -7

10. The sum  $1^4 + 2^4 + 3^4 + 4^4 + \dots + n^4$  is given by the expression  $\frac{6n^5 + an^4 + bn^3 - n}{30}$ .

The value of  $a - b$  is :

- (A) -25
- (B) 25
- (C) -5
- (D) 5

## Section II

Total 60 marks

Attempt Questions 11–14


Allow about 1 hour and 45 minutes for this section

Answer all questions, starting each question on a new page.

In Questions 11–14, your responses should include relevant mathematical reasoning and/or calculations.

---

- Question 11** (15 marks) **Marks**
- (a) Solve  $\frac{2}{x-5} \leq 3$ . 3
- (b) Sketch the function  $f(x) = \pi \cos^{-1}\left(\frac{x}{2}\right)$ , clearly indicating the domain and range of the function. 3
- (c) The velocity of a particle when  $x$  m from the origin is given by  $v^2 = x^2 e^{3x} + 4$ . 2  
Find the acceleration of the particle when  $x = 1$ .
- (d) Find the general solution to the equation  $\sqrt{3} \tan \theta + 1 = 0$ . 2
- (e) Find the value of  $\lim_{x \rightarrow 0} \frac{5x + \sin 3x}{2x}$ . 2
- (f) In the diagram,  $AOB$  is the diameter of the circle with centre at  $O$ .  $TC$  is a tangent to the circle at the point  $C$  such that  $AC$  bisects  $\angle TAB$ . Copy the diagram onto your writing paper. 3  
Prove that  $AT$  is perpendicular to  $TC$ .



**End of Question 11**

**Question 12** (15 marks)**Marks**

- (a) By considering the derivative of  $\ln(\tan x)$ , find  $\int \operatorname{cosec} 2x \, dx$ . 3
- (b) In the expansion of  $\left(x + \frac{2}{x^2}\right)^{10}$ , find the coefficient of  $x$ . 3
- (c)  $P(4p, 2p^2)$  and  $Q(4q, 2q^2)$  are two variable points on the parabola  $x^2 = 8y$ . The tangents at  $P$  and  $Q$  intersect at the point  $T$ .
- (i) Derive the equation of the tangent at  $P$ . 1
- (ii) Hence, show that the point  $T$  has the coordinates  $(2(p+q), 2pq)$ . 2
- (iii) Given that  $p^2 + q^2 = 10$ , determine the cartesian equation of the locus of  $T$ . 2
- (d) Find  $\int (\sin^2 x + 2\cos^2 x + 3\tan^2 x) \, dx$ . 2
- (e) Prove that  $\tan^{-1}(x+1) + \cot^{-1} x = \tan^{-1}(-x^2 - x - 1)$  for  $x > 0$ . 2

**End of Question 12**

**Question 13** (15 marks)

**Marks**

- (a) The growth rate per month of the number  $N$  of birds on a property during a drought is  $-20\%$  of the excess of the bird population over 1000.
- (i) Express the information in the form of a differential equation and show that  $N = 1000 + Ae^{-0.2t}$  (where  $t$  is the time in months) is a solution to this differential equation. 2
- (ii) Given that initially there are 8000 birds on the property, find the amount of time that will elapse before the population is reduced to half. 2
- (b) Ansett Airlines offer two options on all flights for their meal service – chicken or beef (vegetarians choose not to fly with Ansett). If 60% of the time Ansett passengers select the chicken dish, what is the probability that out of 7 randomly selected passengers at least 2 will select chicken for their meal? 2
- (c) An iPhone is thrown from the top of a building, 6 metres high, with an initial velocity of 8 m/s at an angle of  $30^\circ$  to the horizon.
- (i) Using  $10 \text{ m/s}^2$  for acceleration due to gravity, derive the horizontal and vertical equations of motion for the iPhone. 2
- (ii) Determine the greatest height of the iPhone above ground level. 2
- (iii) Find the velocity and direction of the iPhone's path after 1 second. 2
- (d) Prove the following statement is true by mathematical induction for all integers  $n \geq 1$ . 3

$$\sum_{r=1}^n r(r!) = (n+1)! - 1$$

**End of Question 13**

**Question 14** (15 marks)**Marks**

(a) A particle moves in a straight line with simple harmonic motion. At time  $t$  seconds, its displacement  $x$  metres from a fixed point  $O$ , is given by  $x = 2 + 5 \sin\left(3t + \frac{\pi}{4}\right)$ .

(i) Show that  $\ddot{x} = -9(x - 2)$ .

1

(ii) Determine the maximum speed of the particle and its displacement at this time.

2

(b) How many different arrangements of the word MAMMOTH can be made if only five letters are used?

2

(c) Use the substitution  $u^2 = x + 1$  to find the volume of the solid formed by rotating the area bounded by the curve  $y = \frac{x-1}{\sqrt{x+1}}$ , the  $x$  axis and the lines  $x = 3$  and  $x = 8$  about the  $x$  axis. Express your answer in exact form.

4

(d) Use the expansion of  $(1+x)^n$  to prove that

3

$$\frac{n+(-1)^n}{n+1} = \frac{1}{2} {}^n C_1 - \frac{1}{3} {}^n C_2 + \dots + \frac{(-1)^n}{n} {}^n C_{n-1}.$$

(e) Given that  $f(x) = Ax^3 + Bx^2 + Cx + D$  is a function with a double zero at  $x = 1$ , and with a minimum value of  $-4$  when  $x = -1$ , find the values of  $A$ ,  $B$ ,  $C$  and  $D$ .

3

**End of paper**



2015 Trial MSC - Extension 1

SECTION I

1.  ${}^9C_2 \times {}^7C_4 = 1260$  (A)

2.  $\angle ABC = 85$  (ext.  $\angle$  of cyclic quad)  
 $\angle ACB + 34 + 85 = 180$  ( $\angle$  sum of  $\triangle ABC$ )  
 $\therefore \angle ACB = 61^\circ$  (C)

3. let  $u = e^x$   $du = e^x dx$   
 $\int \frac{e^x}{1+(e^x)^2} dx = \int \frac{du}{1+u^2} = \tan^{-1}u + C = \tan^{-1}(e^x) + C$  (A)

4.  $\frac{dV}{dt} = \frac{dV}{dr} \times \frac{dr}{dt}$   $V = \frac{4}{3}\pi r^3$   $\frac{dV}{dr} = 4\pi r^2$   
 $= 4\pi (10)^2 \times 2$   
 $= 800\pi \text{ cm}^3/\text{s}$  (C)

5. when  $t=0$ ,  $y = 21 \text{ m/s}$   $t=?$  when  $y=0$   
 $\ddot{y} = -10$   
 $\dot{y} = -10t + c$   $\therefore c = 21$   
 $0 = -10t + 21$   $10t = 21$   $\therefore t = 2.1 \text{ s}$  (D)

6.  $f(x) = x^3 + x - 1$   $f'(x) = 3x^2 + 1$   
 $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = 0.5 - \frac{(0.5^3 + 0.5 - 1)}{3(0.5)^2 + 1}$   
 $\doteq 0.7143$  (C)

7.  $\frac{dy}{dx} = \frac{1}{\sqrt{1-\frac{a^2}{x^2}}} \times \frac{-a}{x^2} = \frac{-a}{x\sqrt{x^2-a^2}}$  (C)

8. Consider  $(1+x)^n = {}^nC_0 + {}^nC_1 x + {}^nC_2 x^2 + \dots + {}^nC_n x^n$   
 let  $x=1$   
 $(1+1)^n = {}^nC_0 + {}^nC_1 + {}^nC_2 + \dots + {}^nC_n$   
 let  $n=20$   
 $\therefore \sum_{k=1}^{20} {}^{20}C_k + {}^{20}C_0 = (2)^{20}$   
 $\therefore \sum_{k=1}^{20} {}^{20}C_k = 2^{20} - 1 = 1048575$  (B)

9.  $a+b+c = 3$   $abc = -24$   
 $-a-b = -n$   $ab = -6$   
 $a+b = n$   $\therefore c = 4$   
 $\therefore n = 3 - c = 3 - 4$   
 $\therefore n = -1$  (B)

10.  $1^4 = 6 + a + b - 1$   $a+b = 30 - 5 = 25$  (1)  
 $1^4 + 2^4 = \frac{6(2)^5 + a(2)^4 + b(2)^3 - 2}{30}$   
 $190 + 16a + 8b = 510$   
 $2a + b = \frac{320}{8} = 40$  (2)  
 $a = 15, b = 10$   
 $a - b = 5$  (D)

Question 11, (15 Marks)

a)  $\frac{2}{x-5} \leq 3 \quad (x \neq 5)$

$$\frac{2(x-5)^2}{(x-5)} \leq 3(x-5)^2$$


$$2(x-5) \leq 3(x-5)^2$$

$$3(x-5)^2 - 2(x-5) \geq 0$$

$$(x-5)[3(x-5) - 2] \geq 0$$

$$(x-5)(3x-15-2) \geq 0$$

$$(x-5)(3x-17) \geq 0$$




$$x < 5 \text{ or } x \geq \frac{17}{3}$$

(x ≠ 5)      ✓✓✓

\*Many students lost a mark as  $x \neq 5$ .\*

b)  $f(x) = \pi \cos^{-1}\left(\frac{x}{2}\right)$

Domain:  $-1 \leq \frac{x}{2} \leq 1$       Range:  $0 \leq \frac{y}{\pi} \leq \pi$   
 $-2 \leq x \leq 2$        $0 \leq y \leq \pi$



c)  $v^2 = x^2 e^{3x} + 4$

$$a = \frac{d}{dx} \left( \frac{1}{2} v^2 \right)$$

$$= \frac{d}{dx} \frac{1}{2} (x^2 e^{3x} + 4)$$

$$= \frac{1}{2} [x^2 \cdot 3e^{3x} + e^{3x} \cdot 2x]$$

$$= \frac{1}{2} [3x^2 e^{3x} + 2x e^{3x}]$$

$$= \frac{1}{2} e^{3x} (3x^2 + 2x) \quad \checkmark$$

when  $x=1$


$$a = \frac{1}{2} \cdot e^3 \cdot (5)$$

$$a = \frac{5}{2} e^3 \text{ m/s}^2 \quad \checkmark$$

$$\left[ a \doteq 50.21 \text{ m/s}^2 \text{ (2 dec. pl.)} \right]$$

d)  $\sqrt{3} \tan \theta = -1$

$$\tan \theta = \frac{-1}{\sqrt{3}}$$



$$\theta = \pi n + \tan^{-1}\left(\frac{1}{\sqrt{3}}\right)$$

$$\theta = n\pi - \frac{\pi}{6} \quad (n \text{ is an integer})$$

OR  $\theta = n\pi + \frac{5\pi}{6}$



$$e) \lim_{x \rightarrow 0} \frac{5x + \sin 3x}{2x}$$

$$= \lim_{x \rightarrow 0} \frac{5x}{2x} + \lim_{x \rightarrow 0} \frac{\sin 3x}{2x}$$


$$= \frac{5}{2} + \frac{3}{2} \lim_{x \rightarrow 0} \frac{\sin 3x}{3x} \quad (2)$$

$$= \frac{8}{2}$$

$$= 4$$

\* Some confusion for students with this limit.

f)



\* Many more ways to prove that  $AT \perp TC$

\* The proof had to be logical from start to finish.

Join B to C

$\angle ACB = 90^\circ$  ( $\angle$  in a semi circle)

Let  $\angle TAC = \angle CAB = \theta$  ( $AC$  bisects  $\angle TAB$ )

In  $\triangle ACB$ :  $\angle ABC = 90 - \theta$  ( $\angle$  sum of  $\triangle ACB$ )

$\angle TCA = \angle ABC = 90 - \theta$  ( $\angle$  in alternate segment)

In  $\triangle ATC$ :  $\angle ATC + \angle TAC + \angle TCA = 180^\circ$  ( $\angle$  sum  $\triangle ATC$ )

$$\angle ATC + \theta + 90 - \theta = 180$$

$$\angle ATC + 90 = 180$$

$$\angle ATC = 90^\circ$$

$\therefore AT \perp TC$

(3)

Q12 a)  $y = \ln(\tan x)$   
 $y' = \frac{\sec^2 x}{\tan x} = \frac{\frac{1}{\cos^2 x}}{\frac{\sin x}{\cos x}} = \frac{1}{\cos x \sin x}$  ✓

Now  $\frac{1}{\cos x \sin x} = \frac{2}{2 \sin x \cos x} = \frac{2}{\sin 2x} = 2 \operatorname{cosec} 2x$  ✓

$\frac{1}{2} \int 2 \operatorname{cosec} 2x dx = \frac{\ln |\tan x|}{2} + C$  ✓  
 [Some students could not find  $\frac{d[\ln(\tan x)]}{dx}$ , hence couldn't get the final Ans.]

b)  $(x + \frac{2}{x^2})^{10}$

$T_{k+1} = {}^{10}C_k \cdot a^{10-k} \cdot b^k$   
 $= {}^{10}C_k \cdot x^{10-k} \cdot (2x^{-2})^k$  ✓  
 $= {}^{10}C_k \cdot x^{10-k} \cdot 2^k \cdot x^{-2k}$   
 $= {}^{10}C_k \cdot 2^k \cdot x^{10-3k}$

Equating the coefficients of  $x$

$\therefore 10 - 3k = 1$   
 $3k = 9$  ✓  
 $k = 3$

The coefficient of  $x$  is

${}^{10}C_3 \cdot 2^3 = 960$  ✓

[Most students did well in this question.]

Q12 c)  $P(4p, 2p^2)$   $Q(4q, 2q^2)$

i)  $x^2 = 8y$   
 $y = \frac{x^2}{8} \therefore y' = \frac{x}{4}$

At  $P(4p, 2p^2) \therefore y' = \frac{4p}{4} = p$

Equation of the tangent at P.

$y - 2p^2 = p(x - 4p)$

$y = px - 2p^2$  ✓ ①

Similarly the equation of the tangent at Q

$y = qx - 2q^2$  ✓ ②

ii) solve ① and ②

$px - 2p^2 = qx - 2q^2$

$(p - q)x = 2(p + q)(p - q)$  ✓

$x = 2(p + q)$  Sub into ①

$y = p[2(p + q)] - 2p^2$

$y = 2p^2 + 2pq - 2p^2$

$y = 2pq$

Thus  $T(2(p + q), 2pq)$  ✓

[Almost every one did well in parts (c/i) and (c/ii)]

Q12

c/iii)  $x = 2(p+q)$  Given  $p^2+q^2=10$   
 $y = 2pq$

$$x^2 = [2(p+q)]^2 \quad \checkmark$$

$$x^2 = 4(p^2+q^2+2pq)$$

$$x^2 = 4(10+y)$$

$$x^2 = 4(y+10) : \text{Locus of T} \quad \checkmark$$

d)  $\int (\sin^2 x + 2\cos^2 x + 3\tan^2 x) dx$

$$I = \int [\sin^2 x + \cos^2 x + \cos^2 x + 3(\sec^2 x - 1)] dx \quad \checkmark$$

$$I = \int [3\sec^2 x + \frac{1}{2}(1 + \cos 2x) - 2] dx$$

$$I = \int (3\sec^2 x + \frac{1}{2}\cos 2x - \frac{3}{2}) dx$$

$$= 3\tan x + \frac{1}{4}\sin 2x - \frac{3}{2}x + C \quad \checkmark$$

[A number of students forgot Trigonometric identities, double-angle formula, hence could not integrate this question correctly.]

Q12

e) Let  $A = \tan^{-1}(x+1)$ , Let  $B = \cot^{-1}x$   
 $\tan A = x+1$ ,  $\cot B = x$   
 $\tan B = \frac{1}{x}$

$$\tan^{-1}[-(x^2+x+1)] = -\tan^{-1}(x^2+x+1) \quad \checkmark$$

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \cdot \tan B} \quad \checkmark$$

$$= \frac{x+1 + \frac{1}{x}}{1 - (x+1) \cdot \frac{1}{x}}$$

$$= \frac{\frac{x^2+x+1}{x}}{\frac{x-x-1}{x}} = -(x^2+x+1)$$

Thus  $A+B = \tan^{-1}[-(x^2+x+1)] \quad \checkmark$

OR  $\tan^{-1}(x+1) + \cot^{-1}x = \tan^{-1}(-x^2-x-1)$

[Many students could not simplify  $\tan^{-1}(x+1)$ ,  $\cot^{-1}x$  and  $\tan[-(x^2+x+1)]$  correctly]

Q13 Ext 1 2015

a) i)  $\frac{dN}{dt} = k(N-1000)$

$= -0.2(N-1000)$

$N = 1000 + A e^{-0.2t}$

$\frac{dN}{dt} = -0.2 A e^{-0.2t}$

$A e^{-0.2t} = N - 1000$

$= -0.2(N-1000)$

\* The setting out for this question was not very good. For a show question you need to show all the steps.

ii) at  $t=0$ ,  $N=8000$

$8000 = 1000 + A e^0$

$\therefore A = 7000$

$N = 1000 + 7000 e^{-0.2t}$

$4000 = 1000 + 7000 e^{-0.2t}$

$\frac{3}{7} = e^{-0.2t}$

$\ln\left(\frac{3}{7}\right) = -0.2t$

$t = 4.23$  months

\* This question was done very well.

b) P(at least 2 chicken) =

$1 - {}^7C_1(0.6)(0.4)^6 - {}^7C_0(0.6)^7(0.4)^0$

$= 1 - \frac{1344}{78125} - \frac{128}{78125}$

$= 0.98$

\* Many students couldn't do this question.

The particle is travelling at  $9.2$  m/s down wards at  $\theta = 41^\circ$  to the horizontal.

\* many students didn't know how to find the direction

c)

i)

$\ddot{x} = 0$

$\dot{x} = c$

$x = vt \cos \theta$

$x = vt \cos \theta + c$

at  $t=0$ ,  $x=0$

$\therefore c=0$

$x = vt \cos \theta$

$= 8t \cos 30^\circ$

$= 8t \times \frac{\sqrt{3}}{2}$

$x = 4\sqrt{3}t^2$

\* Many students didn't show all the steps of deriving these equations

$\ddot{y} = -10$

$\dot{y} = -10t + c$

$c = v \sin \theta$

$\dot{y} = -10t + v \sin \theta$

$y = -5t^2 + vt \sin \theta + c$

at  $t=0$ ,  $c=6$

$\therefore y = 6$

$y = -5t^2 + vt \sin \theta + 6$

$= -5t^2 + 8t \times \frac{1}{2} + 6$

$y = -5t^2 + 4t + 6$

$y = -5t^2 + 4t + 6$

ii)  $\dot{y} = 0$


$-10t + 4 = 0$

$t = \frac{2}{5}$

\* This question was done well

$y = -5\left(\frac{2}{5}\right)^2 + 8\left(\frac{2}{5}\right) + 6$

$y = 6.8$



iii)  $\dot{x} = 4\sqrt{3}$

$\dot{y} = -10\left(\frac{1}{2}\right) + 8\left(\frac{1}{2}\right)$

$= -6$

$v^2 = (4\sqrt{3})^2 + 36$

$v = 9.2$  m/s

$\tan \theta = \frac{6}{4\sqrt{3}} \therefore \theta = 41^\circ$

d)

$1 + 2(2!) + 3(3!) + \dots + n(n!) = (n+1)! - 1$

when  $n=1$

LHS = 1

RHS =  $2! - 1$

$= 1$

$\therefore$  true for  $n=1$

when

$n=k$

$1 + 2(2!) + \dots + k(k!) = (k+1)! - 1$

prove true for  $n=k+1$

$1 + 2(2!) + \dots + k(k!) + (k+1)(k+1)! = (k+2)! - 1$

LHS =  $(k+1)! - 1 + (k+1)(k+1)!$

$= (k+1)! [1 + k + 1] - 1$

$= (k+1)! (k+2) - 1$

$= (k+2)! - 1$

$=$  RHS

$\therefore$  true for  $n \geq 1$

By Mathematical induction

\* This question was done well, except some students didn't show the last step properly  $\therefore$  lost mark.



### Question 14 (15 Marks)

a)  $x = 2 + 5 \sin\left(3t + \frac{\pi}{4}\right) \Rightarrow x - 2 = 5 \sin\left(3t + \frac{\pi}{4}\right)$

i)  $\dot{x} = 0 + 5 \cos\left(3t + \frac{\pi}{4}\right) \times 3$   
 $= 15 \cos\left(3t + \frac{\pi}{4}\right)$

$$\begin{aligned}\ddot{x} &= -15 \sin\left(3t + \frac{\pi}{4}\right) \times 3 \\ &= -45 \sin\left(3t + \frac{\pi}{4}\right) \quad \checkmark \\ &= -9 \left[ 5 \sin\left(3t + \frac{\pi}{4}\right) \right]\end{aligned}$$

$\therefore \ddot{x} = -9(x - 2)$ , as required.

ii) Maximum speed occurs at the centre of motion i.e. at  $x = 2$

When  $x = 2$ :  $5 \sin\left(3t + \frac{\pi}{4}\right) = 0$   
 $3t + \frac{\pi}{4} = \pi \quad (t > 0)$

$$3t = \frac{3\pi}{4}$$

$$t = \frac{\pi}{4} \text{ sec.}$$

At  $x = 2$ :  $\dot{x} = 15 \cos\left(\frac{3\pi}{4} + \frac{\pi}{4}\right)$   
 $= 15 \cos \pi$   
 $= -15 \text{ m/sec.}$

$\therefore |\dot{x}|_{\max} = 15 \text{ m/sec at } x = 2$

b). different arrangements of the word MAMMOTH using only 5 letters.

Case 1: (1M) M \_ \_ \_ \_ =  $5!$  ways.

Case 2: (2M's) M M \_ \_ \_ =  $\frac{5!}{2!} \times 4C_3$

Case 3: (3M's) M M M \_ \_ =  $\frac{5!}{3!} \times 4C_2$

Total arrangements

$$= 5! + \frac{5!}{2!} \times 4C_3 + \frac{5!}{3!} \times 4C_2$$

$$= 120 + 60 \times 4 + 20 \times 6$$

$$= 120 + 240 + 120$$

$$= \underline{\underline{480}}$$

★ One mark awarded for considering the different cases of repetition.  
One mark for the correct answer.

$$c) V = \pi \int_3^8 y^2 dx$$

$$= \pi \int_3^8 \frac{(x-1)^2}{(x+1)} dx$$

$$\begin{aligned} u^2 &= x+1 \\ 2udu &= dx \end{aligned}$$

$$= \pi \int_2^3 \frac{(u^2-2)^2}{u^2} \cdot 2u du$$

$$\begin{cases} u^2-1 = x \\ u^2-2 = x-1 \end{cases}$$

$$= 2\pi \int_2^3 \left( \frac{u^4}{u} - \frac{4u^2}{u} + \frac{4}{u} \right) du$$

when  $x=8, u^2=9$   
 $u=3$

$$= 2\pi \int_2^3 \left( u^3 - 4u + \frac{4}{u} \right) du$$

$x=3, u^2=4$   
 $u=2$

$$= 2\pi \left[ \frac{u^4}{4} - 2u^2 + 4\ln u \right]_2^3$$

$$= 2\pi \left[ \left( \frac{81}{4} - 18 + 4\ln 3 \right) - \left( 4 - 8 + 4\ln 2 \right) \right]$$

$$= 2\pi \left[ \frac{25}{4} + 4\ln 3 - 4\ln 2 \right]$$

$$= \frac{25\pi}{2} + 8\pi(\ln 3 - \ln 2) \text{ units}^3$$

\* Most students substituted incorrectly with the new variable, back into the integral.

d)

$$(1+x)^n = {}^nC_0 + {}^nC_1 x + {}^nC_2 x^2 + \dots + {}^nC_{n-1} x^{n-1} + {}^nC_n x^n$$

Integrate both sides w.r.t.  $x$ :

$$\frac{(1+x)^{n+1}}{n+1} + C = {}^nC_0 x + \frac{{}^nC_1 x^2}{2} + \frac{{}^nC_2 x^3}{3} + \dots + \frac{{}^nC_{n-1} x^n}{n} + \frac{{}^nC_n x^{n+1}}{n+1}$$

Let  $x=0$ :

$$\frac{1}{n+1} + C = 0 \quad \therefore C = -\frac{1}{n+1}$$

$$\therefore \frac{(1+x)^{n+1} - 1}{(n+1)} = {}^nC_0 x + \frac{{}^nC_1 x^2}{2} + \frac{{}^nC_2 x^3}{3} + \dots + \frac{{}^nC_{n-1} x^n}{n} + \frac{{}^nC_n x^{n+1}}{n+1}$$

Let  $x=-1$ : noting that  ${}^nC_0 = {}^nC_n = 1$

$$\frac{-1}{n+1} = -1 + \frac{1}{2} {}^nC_1 - \frac{1}{3} {}^nC_2 + \dots + \frac{(-1)^n}{n} {}^nC_{n-1} + \frac{(-1)^{n+1}}{n+1}$$

$$1 - \frac{1}{n+1} - \frac{(-1)(-1)^n}{n+1} = \frac{1}{2} {}^nC_1 - \frac{1}{3} {}^nC_2 + \dots + \frac{(-1)^n}{n} {}^nC_{n-1}$$

$$\frac{n+1-1}{n+1} + \frac{(-1)^n}{n+1} = \frac{1}{2} {}^nC_1 - \frac{1}{3} {}^nC_2 + \dots + \frac{(-1)^n}{n} {}^nC_{n-1}$$

$$\therefore \frac{n+(-1)^n}{n+1} = \frac{1}{2} {}^nC_1 - \frac{1}{3} {}^nC_2 + \dots + \frac{(-1)^n}{n} {}^nC_{n-1}$$

\* Most students forgot the constant of integration in the first step.



$$e) f(x) = Ax^3 + Bx^2 + Cx + D$$

$$f'(x) = 3Ax^2 + 2Bx + C$$

$$f(1) = 0 : A + B + C + D = 0 \dots \textcircled{1}$$

$$f'(1) = 0 : 3A + 2B + C = 0 \dots \textcircled{2}$$

$$f'(-1) = 0 : 3A - 2B + C = 0 \dots \textcircled{3}$$

$$f(-1) = -4 : -A + B - C + D = -4 \dots \textcircled{4}$$

$$\textcircled{1} + \textcircled{4} : 2B + 2D = -4 \dots \textcircled{5}$$

$$\textcircled{2} - \textcircled{3} : 4B = 0$$

$$\underline{\underline{B = 0}}$$

$$\text{From } \textcircled{5} : 2D = -4$$

$$\underline{\underline{D = -2}}$$

$$\text{From } \textcircled{1} : A + C - 2 = 0 \dots \textcircled{6}$$

$$\text{From } \textcircled{2} : 3A + C = 0 \dots \textcircled{7}$$

$$\textcircled{7} - \textcircled{6} : 2A + 2 = 0$$

$$2A = -2$$

$$\underline{\underline{A = -1}}$$

$$\text{From } \textcircled{6} : -1 + C - 2 = 0$$

$$\underline{\underline{C = 3}}$$

$$\therefore A = -1, B = 0, C = 3, D = -2.$$