



SYDNEY GRAMMAR SCHOOL
MATHEMATICS DEPARTMENT
TRIAL EXAMINATIONS 2004

FORM VI

MATHEMATICS EXTENSION 1

Examination date

Tuesday 10th August 2004

Time allowed

2 hours (plus 5 minutes reading time)

Instructions

- All seven questions may be attempted.
- All seven questions are of equal value.
- All necessary working must be shown.
- Marks may not be awarded for careless or badly arranged work.
- Approved calculators and templates may be used.
- A list of standard integrals is provided at the end of the examination paper.

Collection

- Write your candidate number clearly on each booklet.
- Hand in the seven questions in a single well-ordered pile.
- Hand in a booklet for each question, even if it has not been attempted.
- If you use a second booklet for a question, place it inside the first.
- Keep the printed examination paper and bring it to your next Mathematics lesson.

Checklist

- SGS booklets: 7 per boy. A total of 1000 booklets should be sufficient.
- Candidature: 121 boys.

Examiner


MLS

QUESTION ONE (12 marks) Use a separate writing booklet. **Marks**

- (a) Solve the inequation $\frac{4}{5-x} \leq 1$. **3**
- (b) For what value of p is the expression $4x^3 - x + p$ divisible by $x + 3$? **2**
- (c) Expand $(a + \frac{1}{2})^5$, expressing each term in its simplest form. **2**
- (d) Given the points $A(1,4)$ and $B(5,2)$, find the co-ordinates of the point that divides the interval AB externally in the ratio 1 : 3. **2**
- (e) Find $\int x(1-x^2)^5 dx$, using the substitution $u = 1-x^2$, or otherwise. **3**

QUESTION TWO (12 marks) Use a separate writing booklet. **Marks**

- (a) Consider the parabola $x = 4t, y = 2t^2$.
 - (i) Find the gradient of the parabola at the point where $t = 4$. **1**
 - (ii) Find the equation of the tangent to the parabola at $t = 4$. **2**



In the diagram above, two circles touch one another externally at the point A . A straight line through A meets one of the circles at T and the other at S . The tangents at T and S meet the common tangent at A at X and Y respectively.


Let $\theta = \angle XTA$.

- (i) Explain why $\angle XAT$ is θ . **1**
- (ii) Prove that $TX \parallel YS$. **2**
- (c) (i) Write down the first three terms in the expansion of $(1 + mx)^n$. **1**
- (ii) If $(1 + mx)^n \equiv 1 - 4x + 7x^2 - \dots$, find the values of m and n . **3**
- (d) Evaluate $\lim_{x \rightarrow 0} \frac{5x \cos 2x}{\sin x}$, showing your reasoning. **2**

QUESTION THREE (12 marks) Use a separate writing booklet.

Marks

(a)



The diagram above shows a container in the shape of a right circular cone. The semi-vertical angle $\theta = \tan^{-1} \frac{1}{2}$.

Water is poured in at the constant rate of 10 cm^3 per minute.

Let the height of the water at time t seconds be h cm, let the radius of the water surface be r cm, and let the volume of water be $V \text{ cm}^3$.

(i) Show that $r = \frac{1}{2}h$. 1

(ii) Show that $V = \frac{1}{12}\pi h^3$. 1

(iii) Find the exact rate at which h is increasing when the height of the water in the cone is 50 cm. 2

(b) Show that there is no term independent of x in the expansion of $\left(2x^2 - \frac{1}{4x}\right)^{11}$. 3

(c) Evaluate $\int_{-1}^0 x\sqrt{1+x} dx$, using the substitution $u = 1+x$. 4

(d) Find $\int \sin x \cos^3 x dx$. 1

QUESTION FOUR (12 marks) Use a separate writing booklet.

Marks

(a) If $y = \frac{1}{200}te^{-t}$, show that $\frac{dy}{dt} = \frac{1}{200}(1-t)e^{-t}$. 1

(b) Fred has recently consumed three standard alcoholic drinks. Immediately after he has finished his last drink, his blood alcohol level is measured over a four-hour period.
Let his blood alcohol level at any time t be A , where t is the time in hours after his last drink.


It is found that the rate of change $\frac{dA}{dt}$ of his blood alcohol content is given by

$$\frac{dA}{dt} = \frac{1}{200}(1-t)e^{-t}, \text{ where } 0 \leq t \leq 4.$$

(i) Show that his blood alcohol content increases during the first hour and decreases after the first hour. 2

(ii) Initially his blood alcohol content was 0.0005. Find A as a function of t . You will need to use part (a). 2

(iii) Determine his maximum alcohol content during the four-hour period. Give your answer correct to four decimal places. 1



The graph of the curve $y = \frac{\pi}{2} - 2 \tan^{-1} x$ is drawn above. It cuts the y -axis at $(0, \frac{\pi}{2})$.

(i) Write down the domain of the inverse function of $y = \frac{\pi}{2} - 2 \tan^{-1} x$. 1

(ii) Find the equation of the inverse function of $y = \frac{\pi}{2} - 2 \tan^{-1} x$. 1

(iii) Find the volume generated when the shaded region is rotated about the y -axis. 4

QUESTION FIVE (12 marks) Use a separate writing booklet.

Marks

(a) Evaluate $\int_0^4 \frac{1}{3 + \sqrt{x}} dx$, using the substitution $x = (u - 3)^2$. 3

(b) (i) Write down the expansion of $(1 + x)^n$ in ascending powers of x . Then differentiate both sides of your identity. 1

(ii) Make an appropriate substitution for x to show that 1

$$\binom{n}{1} + 2 \binom{n}{2} + 3 \binom{n}{3} + 4 \binom{n}{4} + \dots + n \binom{n}{n} = n(2^{n-1}).$$

(iii) Hence find an expression for 1

$$2 \binom{n}{1} + 3 \binom{n}{2} + 4 \binom{n}{3} + 5 \binom{n}{4} + \dots + (n + 1) \binom{n}{n}.$$

(c) Find values for R and α if $\sqrt{3} \sin \theta - \cos \theta = R \cos(\theta + \alpha)$, where R and α are positive constants and $0 < \alpha < 2\pi$. 2


(d) Use the method of mathematical induction to prove that 4

$$\frac{1}{2!} + \frac{2}{3!} + \dots + \frac{n}{(n+1)!} = \frac{(n+1)! - 1}{(n+1)!}, \text{ for all positive integers } n.$$

QUESTION SIX (12 marks) Use a separate writing booklet.

Marks

(a)



The sketch above shows the curve $y = 2 \sin x$ and the line $x - 3y = 0$. The graphs meet at the point A in the first quadrant.

(i) Write down an equation whose solution gives the x -coordinate of A . 1

(ii) An approximate value for the x -coordinate of A is $x = 3$. Apply Newton's method once to find a closer approximation for this value. Give your answer correct to one decimal place. 2

(b) Newton's law of cooling states that a body cools according to the equation

$$\frac{dT}{dt} = -k(T - S),$$

where T is the temperature of the body at time t , S is the temperature of the surroundings and k is a constant.


(i) Show that $T = S + Ae^{-kt}$ satisfies the equation, where A is a constant. 1

(ii) A metal rod has an initial temperature of 470°C and cools to 250°C in 10 minutes. The surrounding temperature is 30°C .

(α) Find the value of A and show that $k = \frac{1}{10} \log_e 2$. 2

(β) Find how much longer it will take the rod to cool to 70°C , giving your answer correct to the nearest minute. 2

(c)



In the diagram above, the straight line ACD is a tangent at A to the circle with centre O . The interval AOB is a diameter of the circle. The intervals BC and BD meet the circle at E and F respectively.

Let $\angle BAF = \beta$.

Copy or trace this diagram into your answer booklet.

(i) Explain why $\angle ABF = \frac{\pi}{2} - \beta$. 1

(ii) Prove that the quadrilateral $CDFE$ is cyclic. 3

QUESTION SEVEN (12 marks) Use a separate writing booklet.

Marks

- (a) Car A and car B are travelling along a straight level road at constant speeds V_A and V_B respectively. Car A is behind car B , but is travelling faster.

When car A is exactly D metres behind car B , car A applies its brakes, producing a constant deceleration of $k \text{ m/s}^2$.

- (i) Using calculus, find the speed of car A after it has travelled a distance x metres under braking. 2

- (ii) Prove that the cars will collide if $V_A - V_B > \sqrt{2kD}$. 4

- (b) A particle is moving in simple harmonic motion of period T about a centre O . Its displacement at any time t is given by $x = a \sin nt$, where a is the amplitude.

- (i) Draw a neat sketch of one period of this displacement–time equation, showing all intercepts. 1

- (ii) Show that $\dot{x} = \frac{2\pi a}{T} \cos \frac{2\pi t}{T}$. 1

- (iii) The point P lies D units on the positive side of O . Let V be the velocity of the particle when it first passes through P . 4

Show that the time between the first two occasions when the particle passes through P is $\frac{T}{\pi} \tan^{-1} \frac{VT}{2\pi D}$.

END OF EXAMINATION

Question 1 2004 Trial Ex I


(a) $\frac{4}{5-x} \leq 1$

$4(5-x) \leq (5-x)^2$ ✓

$4(5-x) - (5-x)^2 \leq 0$

$(5-x)(4 - (5-x)) \leq 0$

$(5-x)(x-1) \leq 0$ ✓



Solution: $x \leq 1$ or $x > 5$. ✓

(b) Let $P(x) = 4x^3 - x + p$.

If $P(x)$ is divisible by $(x+3)$ then $P(-3) = 0$

$P(-3) = -108 + 3 + p$

$-108 + 3 + p = 0$


$p = 105$ ✓

(c) $(a + \frac{1}{2})^5 = a^5 + 5a^4(\frac{1}{2}) + 10a^3(\frac{1}{2})^2 + 10a^2(\frac{1}{2})^3 + 5a(\frac{1}{2})^4 + (\frac{1}{2})^5$

$= a^5 + \frac{5}{2}a^4 + \frac{5}{2}a^3 + \frac{5}{4}a^2 + \frac{5}{16}a + \frac{1}{32}$

✓ for correct bin coeff 1, 5, 10, 10, 5, 1
 ✓ for all correct.

(d)



$x = \frac{lx_2 + kx_1}{l+k}$, $y = \frac{ly_2 + ky_1}{l+k}$

$= \frac{-3+5}{-2}$, $y = \frac{-12+2}{-2}$

$= -1$, $= 5$

The point is $(-1, 5)$ ✓ ✓

Give ✓ if they send the pt that divides AB internally.

(e) $\int x(1-x^2)^5 dx$

$= \int -\frac{1}{2} u^5 du$ ✓

$= -\frac{1}{2} \frac{u^6}{6} + c$

$= -\frac{1}{12} (1-x^2)^6 + c$ ✓

$u = 1-x^2$

$du = -2x dx$ ✓

$-\frac{1}{2} du = x dx$

Question 2

a) (i) $x = 4t$ and $y = 2t^2$
 $\frac{dx}{dt} = 4$ and $\frac{dy}{dt} = 4t$
 so $\frac{dy}{dx} = \frac{4t}{4} = t$
 so at $t=4$, the gradient is 4 ✓

(ii) When $t=4$, $x=16$, $y=32$ ✓

Tangent equation is $y - 32 = 4(x - 16)$
 $y - 32 = 4x - 64$
 $y = 4x - 32$ ✓

b) (i) Tangents to a circle from an external point are equal.
 $\therefore TX = XA$ ✓
 So $\triangle TXA$ is isosceles and $\angle XTA = \angle XAT = \theta$
 (base angles of isosceles triangle)

(ii) Similarly, $\triangle AYS$ is isosceles with base angles $\angle YAS$ and $\angle YSA$ equal.
 But $\angle YAS = \angle TPA = \theta$ (vertically opposite)
 So $\angle YSA = \theta$ and $\angle XTA = \theta$
 But these are alternate
 So $TX \parallel YS$ ✓

(c) (i) $(1 + mx)^n = 1 + nmx + \frac{n(n-1)}{2}(mx)^2 + \dots$ ✓

(ii) $1 - 4x + 7x^2 - \dots = 1 + nmx + \frac{n(n-1)}{2}(mx)^2 + \dots$

equate coefficients of x : $-4 = nm$ ①
 equate coefficients of x^2 : $7 = \frac{n(n-1)}{2}m^2$ ② ✓

From ①, $m = -\frac{4}{n}$, substitute this in ②.

$14 = n(n-1) \frac{16}{n^2}$

$14 = \frac{16(n-1)}{n}$

$14n = 16n - 16$

$2n = 16$

$n = 8$

$m = -\frac{4}{8}$

$= -\frac{1}{2}$


$n = 8$ and $m = -\frac{1}{2}$ ✓

✓ for a sensible method

(d) $\lim_{x \rightarrow 0} \frac{5x \cos 2x}{\sin 2x} = \lim_{x \rightarrow 0} \frac{5x}{\sin 2x} \times \lim_{x \rightarrow 0} \cos 2x$
 $= 5 \lim_{x \rightarrow 0} \frac{x}{\sin x} \times 1$ ✓
 $= 5 \times 1$
 $= 5$ ✓

Q3

a) (i)



(ii) $V = \frac{1}{3} \pi r^2 h$
 $= \frac{1}{3} \pi \left(\frac{1}{2}h\right)^2 h$ ✓
 $= \frac{1}{12} \pi h^3$

(iii) Find $\frac{dh}{dt}$, given $\frac{dV}{dt} = 10$

$\frac{dh}{dt} = \frac{dh}{dV} \frac{dV}{dt}$ ✓

now $V = \frac{1}{12} \pi h^3$

$\frac{dV}{dh} = \frac{\pi h^2}{4}$ so $\frac{dh}{dV} = \frac{4}{\pi h^2}$

so $\frac{dh}{dt} = \frac{4}{\pi h^2} \times 10$

$= \frac{4}{\pi \times 50^2} \times 10$ when $h = 50$

$= \frac{40}{\pi \times 50 \times 50}$ ✓

$= \frac{4}{250\pi}$ cm per minute

(b)

The general term is $\binom{22}{r} (2x^2)^{22-r} (-4x)^r$ ✓

The index of x is $22 - 2r + r = 0$ if the term is independent of x .
 $3r = 22$ ✓
 $r = \frac{22}{3}$ ✓

since r must be an integer, there is no term independent of x .

(c) $\int_{-1}^0 x\sqrt{1+x} dx$ $u = 1+x$
 $du = dx$
 when $x=0$, $u=1$ ✓
 when $x=-1$, $u=0$ ✓
 $= \int_0^1 (u-1)u^{1/2} du$
 $= \int_0^1 (u^{3/2} - u^{1/2}) du$ ✓
 $= \left[\frac{2u^{5/2}}{5} - \frac{2u^{3/2}}{3} \right]_0^1$ ✓
 $= \left(\frac{2}{5} - \frac{2}{3} \right) - (0)$
 $= -\frac{4}{15}$ ✓

(d) $\int \sin \cos^2 x dx$
 $= -\frac{1}{3} \cos^3 x + c$ ✓

Question 4

a) $y = \frac{1}{200} t e^{-t}$

$$\frac{dy}{dt} = \frac{1}{200} [t \times (-e^{-t}) + e^{-t}] \quad \checkmark$$

$$= \frac{1}{200} e^{-t} (1-t)$$

b) (i) $\frac{dA}{dt} = \frac{1}{200} (1-t) e^{-t}$

Now, $\frac{1}{200} (1-t) e^{-t} > 0$ when $1-t > 0$ or $t < 1$ ✓

so, for $0 < t < 1$, $\frac{dA}{dt} > 0$ and A is

increasing.

and $\frac{1}{200} (1-t) e^{-t} < 0$ when $1-t < 0$ or $t > 1$ ✓

so, for $t > 1$, $\frac{dA}{dt} < 0$ and A is

decreasing.

(ii) $t=0, A=0.0005$

$$A = \int \frac{1}{200} (1-t) e^{-t} dt$$

$$= \frac{1}{200} t e^{-t} + c \quad \text{from (a).} \quad \checkmark$$

when $t=0, 0.0005 = 0 + c$ so $c = 0.0005$

$$A = \frac{1}{200} t e^{-t} + 0.0005 \quad \checkmark$$

(iii) From (i), the maximum A is when $t=1$.

so, if $t=1, A = \frac{1}{200} e^{-1} + 0.0005$ ✓

$$= 0.001839 + 0.0005$$

$$= 0.002339$$

$$= 0.0023$$

(c) (i) $-\frac{\pi}{2} < x < \frac{3\pi}{2}$ ✓

(ii) $x = \frac{\pi}{2} - 2 \tan^{-1} y$

$$2 \tan^{-1} y = \frac{\pi}{2} - x$$

$$\tan^{-1} y = \frac{1}{2} (\frac{\pi}{2} - x)$$

$$y = \tan(\frac{\pi}{4} - \frac{1}{2}x), \quad -\frac{\pi}{2} < x < \frac{3\pi}{2}$$

(iii) $V = \pi \int x^2 dy$

or, using the inverse function

$$V = \pi \int y^2 dx$$

$$= \pi \int_0^{\frac{\pi}{2}} \tan^2(\frac{\pi}{4} - \frac{1}{2}x) dx \quad \checkmark$$

$$= \pi \int_0^{\frac{\pi}{2}} \sec^2(\frac{\pi}{4} - \frac{1}{2}x) - 1 dx \quad \checkmark$$

$$= \pi [-2 \tan(\frac{\pi}{4} - \frac{1}{2}x) - x]_0^{\frac{\pi}{2}} \quad \checkmark$$

$$= \pi \left[\frac{-2 \tan 0 - \frac{\pi}{2}}{\pi(0 - \frac{\pi}{2})} - \frac{(-2 \tan \frac{\pi}{4} - 0)}{1} \right] \quad \checkmark$$

question 5

$$1 \int_0^4 \frac{1}{3+\sqrt{x}} dx$$

$$x = (u-3)^2$$

$$dx = 2(u-3) du$$

when $x=4$, $u=5$
when $x=0$, $u=3$.

$$= \int_3^5 \frac{2(u-3) du}{u}$$

$$= 2 \int_3^5 \left(1 - \frac{3}{u}\right) du$$

$$= 2 \left[u - 3 \log_e u \right]_3^5$$

$$= 2 \left[(5 - 3 \log 5) - (3 - 3 \log 3) \right]$$

$$= 2 (2 - 3 \log 5 + 3 \log 3)$$

$$= 2 \left(2 + \log \frac{3}{5} \right)$$

$$(b)(i) (1+x)^n = 1 + \binom{n}{1}x + \binom{n}{2}x^2 + \binom{n}{3}x^3 + \dots + \binom{n}{n}x^n$$

$$n(1+x)^{n-1} = \binom{n}{1} + 2\binom{n}{2}x + 3\binom{n}{3}x^2 + \dots + n\binom{n}{n}x^{n-1}$$

(ii) let $x=1$

$$LHS = n(2)^{n-1}$$

$$RHS = \binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \dots + n\binom{n}{n}$$

(iii) let $x=1$ in expansion of $(1+x)^n$

$$2^n = 1 + \binom{n}{1} + \binom{n}{2} + \binom{n}{3} + \dots + \binom{n}{n}$$

$$\text{and } n2^{n-1} = \binom{n}{1} + 2\binom{n}{2} + 3\binom{n}{3} + \dots + n\binom{n}{n}$$

$$\text{adding } 2^n + n2^{n-1} = 1 + 2\binom{n}{1} + 3\binom{n}{2} + 4\binom{n}{3} + \dots + (n+1)\binom{n}{n}$$

$$\text{so } 2^n + n2^{n-1} - 1 = 2\binom{n}{1} + 3\binom{n}{2} + 4\binom{n}{3} + \dots + (n+1)\binom{n}{n}$$

$$(c) \sqrt{3} \sin \theta - \cos \theta = R \cos(\theta + \alpha)$$

$$= R \cos \theta \cos \alpha - R \sin \theta \sin \alpha$$

$$\text{so } -1 = R \cos \alpha \quad \text{and} \quad \sqrt{3} = -R \sin \alpha$$

$$R = \sqrt{3+1}$$

$$= 2$$

$$\text{and } \alpha = \frac{4\pi}{3}$$

(d)

If $n=1$

$$LHS = \frac{1}{2!} = \frac{1}{2}$$

$$RHS = \frac{2! - 1}{2!}$$

$$= \frac{1}{2}$$

so the statement is true when $n=1$.

Now suppose the statement is true for some value of n , k .

$$\text{i.e. } \frac{1}{2!} + \frac{2}{3!} + \dots + \frac{k}{(k+1)!} = \frac{(k+1)! - 1}{(k+1)!}$$

We now prove the result for $n = k+1$.

$$\text{That is prove that } \frac{1}{2!} + \frac{2}{3!} + \dots + \frac{k}{(k+1)!} + \frac{k+1}{(k+2)!} = \frac{(k+2)! - 1}{(k+2)!}$$

$$\text{Now, LHS} = \frac{(k+1)! - 1}{(k+1)!} + \frac{k+1}{(k+2)!} \quad \text{using induction hypothesis}$$

$$\begin{aligned}
 \text{LHS} &= \frac{(k+1)! - 1}{(k+1)!} + \frac{k+1}{(k+2)!} \\
 &= \frac{(k+2)(k+1)! - (k+2) + (k+1)}{(k+2)!} \quad \checkmark \\
 &= \frac{(k+2)! - 1}{(k+2)!} \\
 &= \text{RHS.}
 \end{aligned}$$

So, the statement is true for $k+1$ as long as it is true for k . Hence, by the principle of mathematical induction, it is true for all positive integers n .

Question 6.

(a) i) At A , $y = 2 \sin x$ and $y = \frac{1}{3}x$ ✓
 so we want $2 \sin x = \frac{1}{3}x$ ✓
 or $2 \sin x - \frac{1}{3}x = 0$

ii) Let $f(x) = 2 \sin x - \frac{1}{3}x$ }
 $f'(x) = 2 \cos x - \frac{1}{3}$ }
 $x_1 = x_0 = \frac{f(x_0)}{f'(x_0)}$, $x_0 = 3$
 $= 3 - \frac{2 \sin 3 - 1}{2 \cos 3 - \frac{1}{3}}$ ✓
 $= 3 - 0.3102$ ✓
 $= 2.7$ ✓

(b) (i) $T = S + Ae^{-kt}$
 $\frac{dT}{dt} = -kAe^{-kt}$, $Ae^{-kt} = T - S$
 $= -k(T - S)$

(i) (a) $T = 30^\circ + Ae^{-kt}$
 when $t=0$, $47.0^\circ = 30^\circ + Ae^0$ ✓
 $A = 17.0^\circ$

when $t=10$, $25.0 = 30 + 17.0e^{-10k}$
 $440e^{-10k} = 220$
 $e^{-10k} = \frac{1}{2}$

$-10k = \log_e \frac{1}{2}$
 $k = -\frac{1}{10} \log_e \frac{1}{2}$ ✓
 $= \frac{1}{10} \log_e 2$

(b) Find t when $T = 70^\circ$
 $70 = 30 + 440e^{-kt}$ $k = \frac{1}{10} \ln 2$
 $e^{-kt} = \frac{40}{440}$
 $= \frac{1}{11}$
 $-kt = \log_e \frac{1}{11}$
 $t = \frac{\log_e \frac{1}{11}}{-\frac{1}{10} \ln 2}$
 $\approx 35 \text{ min}$

(c) (i) $\angle AFB = \frac{\pi}{2}$ (the angle in a semicircle is a right angle).
 So $\angle ABF = \pi - (\frac{\pi}{2} + \beta)$ (the angle sum of $\triangle AFB$ is π)
 $= \frac{\pi}{2} - \beta$

(ii) $\angle BAD = \frac{\pi}{2}$ (angle between tangent and radius is $\frac{\pi}{2}$).
 So $\angle ADB = \pi - (\frac{\pi}{2} + (\frac{\pi}{2} - \beta))$ (angle sum of $\triangle ADB$ is π)
 $= \beta$

Now, $\angle BAF = \angle BEF$ (both subtended at the circumference by arc BF)
 $= \beta$

So, $\angle BEF = \angle CDF$
 So, CDFE is cyclic (exterior angle equals interior opposite angle)

Question 2

(a) (i) For car A
 $\ddot{x} = -b$, since the car is decelerating
 $\frac{1}{2}v^2 = -bx + c$

When $x=0$, $v = V_A$
 so $\frac{1}{2}V_A^2 = 0 + c$ making $c = \frac{1}{2}V_A^2$
 $v^2 = -2bx + V_A^2$
 and speed = $\sqrt{V_A^2 - 2bx}$

(ii) For car A:
 $\ddot{x} = -k$
 Integrating, $\dot{x} = -kt + c_1$
 When $t=0$, $\dot{x} = V_A$ making $c_1 = V_A$
 So $\dot{x} = -kt + V_A$
 Integrating, $x = -\frac{1}{2}kt^2 + tV_A + c_2$
 When $t=0$, $x=0$, taking the origin of displacement at car A, so $c_2 = 0$.
 We have $x = tV_A - \frac{1}{2}kt^2$

For car B:
 $\ddot{x} = 0$
 Integrating, $\dot{x} = c_3$
 When $t=0$, $\dot{x} = V_B$ making $c_3 = V_B$
 So $\dot{x} = V_B$
 Integrating, $x = tV_B + c_4$
 When $t=0$, $x=D$, car B is D metres in front of car A, making $c_4 = V_B$
 So $x = tV_B + D$

When the cars collide, their displacements are equal, so we have
 $tV_B + D = tV_A - \frac{1}{2}kt^2$

This is a quadratic in t .
 For t to have a real value, the
 discriminant must be positive
 $\frac{1}{2}kt^2 - tV_A + tV_B + D = 0$
 $kt^2 - 2t(V_A - V_B) + 2D = 0$


$$\Delta = 4(V_A - V_B)^2 - 8kD$$

$$4(V_A - V_B)^2 - 8kD > 0$$

$$(V_A - V_B)^2 > 2kD$$

$$V_A - V_B > \sqrt{2kD}$$

since $V_A > V_B$ and so $V_A - V_B$ is positive:




(ii)

$$x = a \sin nt$$

$$\dot{x} = na \cos nt$$

$$= \frac{2\pi a}{T} \cos \frac{2\pi}{T} t$$



At P we have $D = a \sin \frac{2\pi}{T} t$ ①

And $V = \frac{2\pi a}{T} \cos \frac{2\pi}{T} t$ ②

$$\textcircled{1} \div \textcircled{2} \quad \frac{D}{V} = \frac{aI}{2\pi a} \tan \frac{2\pi t}{T} \quad \checkmark$$

$$\frac{D \cdot 2\pi}{VT} = \tan \frac{2\pi t}{T}$$

Let t_1 and t_2 be the first two times
 upon the particle is at P.

Then $\frac{2\pi t_1}{T} = \tan^{-1} \frac{2\pi D}{VT}$

$$t_1 = \frac{T}{2\pi} \tan^{-1} \frac{2\pi D}{VT}$$

And $t_2 = \frac{T}{2} - \frac{T}{2\pi} \tan^{-1} \frac{2\pi D}{VT} \quad \checkmark$

So the difference in times is

$$t_2 - t_1 = \frac{T}{2} - \frac{2T}{2\pi} \tan^{-1} \frac{2\pi D}{VT}$$

$$= \frac{T}{\pi} \left(\frac{\pi}{2} - \tan^{-1} \frac{2\pi D}{VT} \right) \quad \checkmark$$

$$= \frac{T}{\pi} \tan^{-1} \frac{VT}{2\pi D}, \text{ using}$$

complementary angles.

