

Name: \_\_\_\_\_

Maths Class: \_\_\_\_\_

# SYDNEY TECHNICAL HIGH SCHOOL



## TRIAL HIGHER SCHOOL CERTIFICATE

2008

### EXTENSION 1 MATHEMATICS

**Instructions:**

**General Instructions**

- Reading time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- All necessary working should be shown in every question
- Start each question on a new page

**Total Marks – 84**

- Attempt Questions 1 – 7
- All questions are of equal value

(For markers use only)

Q1	Q2	Q3	Q4	Q5	Q6	Q7	Total

### Question 1

- a) Differentiate:
- i)  $x^2 \cos^{-1} x$  2
- ii)  $\log_{10} 3x$  2
- b) There is a remainder of 1 when  $P(x) = x^3 - 3x^2 + px - 14$  is divided by  $x - 3$ . Find the value of  $p$ . 2
- c) Find the simultaneous solution of:  $|x - 3| < 4$  and  $|x - 1| > 1$  3
- d) The point  $P(3, 5)$  divides the interval joining  $A(-1, 1)$  and  $B(5, 7)$  internally in the ratio  $m:n$ . Find  $m:n$ . 2
- e) Find  $\int \cos x \sin x \, dx$  1

### Question 2 ( Start a new page )

- a) Find  $\lim_{x \rightarrow \infty} \frac{3x^2 - 7x}{5 + x^2}$  1
- b) Find the acute angle, to the nearest degree, between the curve  $y = x^2$  and the line  $5x - y - 6 = 0$  at the point of intersection  $(3, 9)$  2
- c) i) Solve  $t^2 + 2t - 1 = 0$  1
- ii) Using your results from part i), and the expansion for  $\tan 2\theta$ , find the exact value of  $\tan 22.5^\circ$ . Simplify your answer. 2

- d) i) Express  $3 \cos x - 2 \sin x$  in the form  $A \cos(x + \alpha)$  where  $A > 0$  and  $0^\circ \leq \alpha \leq 90^\circ$  2
- ii) Hence find the smallest positive  $x$  degrees such that  $3 \cos x - 2 \sin x$  has a maximum value (do not use calculus). Give your answer correct to 1 decimal place. 1
- e) Express  $\sin(\tan^{-1}x + \tan^{-1}y)$  in terms of  $x$  and  $y$  only. 3

**Question 3** (Start a new page)


- a) Solve for  $0 \leq \theta \leq 2\pi$ :  $\cos 2\theta = \cos^2 \theta$  2
- b) Solve  $\frac{x^2}{x-4} < 0$  2
- c) Find  $\int \frac{x+4}{x^2+4} dx$  2
- d) Use the substitution  $u = e^x$  to find  $\int \frac{e^x}{\sqrt{9-4e^{2x}}} dx$  3
- e)  $\alpha, \beta, \gamma$  are the roots of the equation  $2x^3 + 5x - 3 = 0$  3
- Find the value of  $\alpha^2 + \beta^2 + \gamma^2$ .

Question 4 ( Start a new page )

a)

Not to scale

3




$O$  is the centre of the circle

$$\angle PXY = 35^\circ \text{ and } \angle PQY = 25^\circ$$

- i) Copy the diagram onto your answer paper
- ii) Find  $\angle MPY$  giving full reasons

b)




The points  $P(2p, p^2)$  and  $Q(2q, q^2)$   
 move on the parabola  $x^2 = 4y$  such that the  
 chord  $PQ$  subtends a right angle at the origin  $O$

- i) Show that  $pq = -4$  2
- ii)  $M$  is the midpoint of  $PQ$ . Derive the locus of  $M$  and show that it is the  
 parabola  $y = \frac{x^2+8}{2}$  2
- iii) Find the focus of the parabola for  $M$ . 1

c) Prove by mathematical induction, that

$$1 \times 2^0 + 2 \times 2^1 + 3 \times 2^2 + \dots + n \times 2^{n-1} = 1 + (n-1)2^n \text{ where } n \text{ is a positive integer} \quad 4$$

**Question 5 ( Start a new page )**



- i) Copy the diagram onto your answer page
- ii) Prove that  $BCED$  is a cyclic quadrilateral 3
- iii) Describe how to find the centre of the circle passing through  $B, C, E, D$ . 1

b) Given  $f(x) = \frac{2}{x+1}$  for  $x > -1$ :

- i) Find the equation of the inverse function  $f^{-1}(x)$  1
- ii) On the same diagram, sketch the graphs of  $y = f(x)$  and  $y = f^{-1}(x)$ . 3

Clearly show the coordinates of any points of intersection, intercepts on the coordinates axes and equations of any asymptotes.

- c) i) Sketch the curve  $y = \sin^{-1}\left(\frac{x}{2}\right)$  1
- ii) The area between the curve  $y = \sin^{-1}\left(\frac{x}{2}\right)$  and the  $y$  axis is rotated about the  $y$  axis. 3  
 Find the volume thus generated.

**Question 6 ( Start a new page )**

a) Differentiate  $y = \tan^{-1}(\sin 3x)$  2

b) In a population study, the population  $P$  of a town after  $t$  years is given by

$$P = 200 + Ae^{kt}.$$

The initial population was 300 and increased to 400 over 3 years.

i) Find the population after a further 2 years (nearest whole person) 3

ii) Find the rate of population growth after 10 years. 1

c) Kramer hits a golf ball from the top of the edge of a vertical cliff 25 metres above the sea. He hits it with an initial velocity of 50 m/s at a  $30^\circ$  angle of elevation.

The cliff top is taken as the point of origin.

i) Given  $\ddot{x} = 0$  and  $\ddot{y} = -10$ , derive the equations of the horizontal and vertical components of the motion for the golf ball. 2

ii) Find the maximum height of the golf ball above the cliff. 2

iii) Find the angle at which the golf ball hits the water (nearest degree). 2

**Question 7 ( Start a new page )**

a) A particle is moving according to the velocity equation  $v = 4 - 2t$  m/s. Find the total distance it travels in the first 5 seconds of its motion. 2

b) A particle is moving with simple harmonic motion in a straight line with velocity

$$v^2 = 108 + 36x - 9x^2 \text{ where } x \text{ cm is its displacement from a point } O.$$


Initially it is at rest at  $x = 6$  cm.

- i) Use differentiation to find its acceleration in terms of  $x$  and find its maximum acceleration. 2
- ii) Find the maximum speed of the particle and the time when this first occurs. 3
- iii) Write an expression for the particle's displacement in terms of time  $t$ . 1

c) A vertical pole, 2 metres high, casts a lengthening shadow as the sun sets. 4

At a particular instant, the shadow's length,  $x$ , is increasing by 0.3m/min.

Simultaneously, the angle of the Sun,  $\theta$ , is decreasing by 0.05 radians/min.



Find the angle  $\theta$  (to the nearest degree) when this is occurring.

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \quad \text{if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

**NOTE:**  $\ln x = \log_e x, \quad x > 0$



# 2008 Extension Solutions

$$\textcircled{1} \text{ a) i) } \frac{dy}{dx} = 2x \cos^{-1} x - \frac{x^2}{\sqrt{1-x^2}}$$

$$\text{ii) } y = \frac{\log 3x}{\log 10}$$

$$\frac{dy}{dx} = \frac{1}{x \log 10}$$

$$\text{b) } P(3) = 1$$

$$\therefore 27 - 27 + 3p - 14 = 1$$

$$\therefore 3p = 15$$

$$\therefore p = 5$$

$$\text{c) } |x-3| < 4 \Rightarrow -4 < x-3 < 4$$
$$-1 < x < 7$$

$$|x-1| > 1 \Rightarrow x-1 > 1 \text{ or } x-1 < -1$$
$$x > 2 \text{ or } x < 0$$

$\therefore$  Simultaneous sol. is

$$-1 < x < 7 \text{ or } -1 < x < 0$$

$$\text{d) } 3 = \frac{-n + 5m}{m+n}$$

$$3m + 3n = -n + 5m$$

$$-2m = -4n$$

$$m = 2n$$

$$\frac{m}{n} = 2$$

$$\therefore m:n = 2:1$$

$$\text{e) } \frac{\sin^2 x}{2} + c$$

$$\textcircled{2} \text{ a) } \lim_{x \rightarrow \infty} \frac{3 - \frac{7}{x}}{\frac{5}{x^2} + 1} = \frac{3-0}{0+1}$$

$$= 3$$

$$\text{b) } \frac{dy}{dx} = 2x \Rightarrow m_1 = 6$$
$$m_2 = 5$$

$$\tan \theta = \left| \frac{6-5}{1+30} \right|$$

$$= \frac{1}{31}$$

$$\therefore \theta \approx 2^\circ$$

$$\text{c) i) } t = \frac{-2 \pm \sqrt{4+4}}{2}$$
$$= \frac{-2 \pm 2\sqrt{2}}{2}$$
$$= -1 \pm \sqrt{2}$$

$$\text{ii) } \tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

$$\tan 45^\circ = \frac{2 \tan 22.5^\circ}{1 - \tan^2 22.5^\circ}$$

$$\therefore 1 - \tan^2 22.5^\circ = 2 \tan 22.5^\circ$$

$$\therefore \tan^2 22.5^\circ + 2 \tan 22.5^\circ - 1 = 0$$

$$\therefore \tan 22.5^\circ = -1 + \sqrt{2} (> 0)$$

(from i) above)

$$d) i) 3\cos x - 2\sin x = A \cos(x+\alpha)$$

$$(A = \sqrt{13})$$

$$= \sqrt{13} \cos(x+\alpha)$$

$$\therefore \frac{3}{\sqrt{13}} \cos x - \frac{2}{\sqrt{13}} \sin x = \cos(x+\alpha)$$

$$= \cos x \cos \alpha - \sin x \sin \alpha$$

$$\therefore \cos \alpha = \frac{3}{\sqrt{13}} \quad \therefore \alpha = 33.7^\circ$$

$$\sin \alpha = \frac{2}{\sqrt{13}}$$

$$\therefore 3\cos x - 2\sin x = \sqrt{13} \cos(x+33.7^\circ)$$

ii) max. value of  $3\cos x - 2\sin x$   
 $=$  max value of  $\sqrt{13} \cos(x+33.7^\circ)$


$$= \sqrt{13}, \text{ and this occurs}$$

$$\text{when } \cos(x+33.7^\circ) = 1$$

$$\therefore x + 33.7^\circ = 360^\circ (\text{not } 0^\circ)$$

$$\therefore x = 326.3^\circ$$

e) Let  $\tan^{-1} x = \alpha, \tan^{-1} y = \beta$   
 $x = \tan \alpha, y = \tan \beta$



$$\sin(\tan^{-1} x + \tan^{-1} y)$$

$$= \sin(\alpha + \beta)$$

$$= \sin \alpha \cos \beta + \cos \alpha \sin \beta$$

$$= \frac{x}{\sqrt{1+x^2}} \cdot \frac{1}{\sqrt{1+y^2}} + \frac{1}{\sqrt{1+x^2}} \cdot \frac{y}{\sqrt{1+y^2}}$$


$$= \frac{x+y}{\sqrt{(1+x^2)(1+y^2)}}$$

$$\frac{x+y}{\sqrt{(1+x^2)(1+y^2)}}$$

3) a)  $2\cos^2 \theta - 1 = \cos^2 \theta$   
 $\cos^2 \theta = 1$   
 $\cos \theta = \pm 1$   
 $\theta = 0, \pi, 2\pi$

b)  $\frac{x^2}{x-4} \times (x-4)^2 < 0$

$$x^2(x-4) < 0$$



$$\therefore x < 0 \text{ or } 0 < x < 4$$

$$\text{or } x < 4 (x \neq 0)$$

c)  $\int \frac{x}{x^2+4} dx + \int \frac{4}{x^2+4} dx$

$$= \frac{1}{2} \log(x^2+4) + 2 \tan^{-1}\left(\frac{x}{2}\right) + c$$

d)  $\int \frac{e^x}{\sqrt{1-4e^{2x}}} dx = \int \frac{u}{\sqrt{4-4u^2}} \frac{du}{u}$

$$u = e^x \quad = \int \frac{1}{2\sqrt{4-u^2}} du$$

$$\frac{du}{dx} = e^x$$

$$dx = \frac{du}{e^x} = \frac{du}{u}$$

$$= \frac{du}{u}$$

$$= \frac{1}{2} \sin^{-1}\left(\frac{u}{2}\right) + c$$


$$= \frac{1}{2} \sin^{-1}\left(\frac{e^x}{2}\right) + c$$

e)  $(\alpha+\beta+\gamma)^2 = \alpha^2 + \beta^2 + \gamma^2 + 2\alpha\beta + 2\beta\gamma + 2\alpha\gamma$

$$\therefore \alpha^2 + \beta^2 + \gamma^2 = (\alpha+\beta+\gamma)^2 - 2(\alpha\beta + \beta\gamma + \alpha\gamma)$$

$$= (\alpha+\beta+\gamma)^2 - 2(\alpha\beta + \beta\gamma + \alpha\gamma)$$

4) a) i)



$\angle PMQ = 35^\circ$  (angles standing on same chord PY)

$\angle XPY = 90^\circ$  (angle in a semicircle)

$\angle YPQ = 120^\circ$  (angle sum of  $\triangle MPQ$ )

$\therefore \angle MPY = 30^\circ$

b) i)  $m_{OP} = \frac{p}{2p}$ ,  $m_{OQ} = \frac{q}{2}$   
 $= \frac{p}{2}$

$m_{OP} \times m_{OQ} = -1$  for perpend. lines

$\frac{p}{2} \times \frac{q}{2} = -1$

$\therefore pq = -4$  as reqd.

ii) M has coords  $\left( \frac{2p+2q}{2}, \frac{p^2+q^2}{2} \right)$

$\therefore x = p+q, y = \frac{p^2+q^2}{2}$   
 $= \frac{(p+q)^2 - 2pq}{2}$   
 $= \frac{x^2 - 2(-4)}{2}$

iii)  $2y = x^2 + 8$

$x^2 = 2y - 8$

$(x-0)^2 = 2(y-4)$

$\therefore$  vertex at  $(0, 4)$  and  $4a = 2$   
 $\therefore a = \frac{1}{2}$

$\therefore$  focus at  $(0, 4\frac{1}{2})$ .

c) Test  $n=1 \Rightarrow$  LHS =  $1 \times 2^0$ , RHS =  $1 + 0 \times 2^1$   
 $= 1$   $= 1$

$\therefore$  result is true for  $n=1$

Assume result is true for  $n=k$ ,

$\therefore$  assume that  $S_k = 1 + (k-1)2^k$

Prove true for  $n=k+1$ ,

$\therefore$  prove that  $S_{k+1} = 1 + k \cdot 2^{k+1}$


Now  $S_{k+1} = S_k + T_{k+1}$   
 $= 1 + (k-1)2^k + (k+1)2^k$   
 $= 1 + 2^k(k-1+k+1)$   
 $= 1 + 2^k \cdot 2k$   
 $= 1 + k \cdot 2^k \cdot 2$   
 $= 1 + k \cdot 2^{k+1}$

(shown)

So, if the result is true for  $n=k$ , then it has been proved true for  $n=k+1$

Since the result is true for  $n=1$ , then from above it must be true for  $n=1+1=2$  and so on for

5 a) i)



ii)  $\angle MAN = \angle ADE$  (alt. angles  
 $MN \parallel DE$ )

$\angle MAN = \angle BCA$  (angle in alt.  
 segment)

$\therefore \angle ADE = \angle BCA$

$\therefore BCED$  is a cyc. quad since  
 exterior angle equals interior  
 opposite angle.

iii) Perpendicular bisectors  
 of at least 2 sides of  $BCED$   
 meet at the centre of the  
 circle.


b) i)  $x = \frac{2}{y+1}$

$$xy + x = 2$$


$$xy = 2 - x$$

$\therefore f^{-1}(x) \Rightarrow y = \frac{2-x}{x}$  or  $\frac{2}{x} - 1$   
 (for  $x > 0$ )

ii)



c) i)



ii)  $V = 2\pi \int_0^{\pi/2} (2 \sin y)^2 dy$   
 $= 8\pi \int_0^{\pi/2} \sin^2 y dy$   
 $= 8\pi \int_0^{\pi/2} \frac{1}{2} (1 - \cos 2y) dy$   
 $= 4\pi \left[ y - \frac{\sin 2y}{2} \right]_0^{\pi/2}$   
 $= 4\pi \left[ \frac{\pi}{2} - 0 - (0 - 0) \right]$   
 $= 2\pi^2 u^3$

6

a)  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$  ( $u = \sin 3x$ )  
 $= \frac{1}{1+u^2} \times 3 \cos 3x$   
 $= \frac{3 \cos 3x}{1 + \sin^2 3x}$

b) i)  $P = 300, t = 0$ :

$$300 = 200 + A \quad (A = 100)$$

$$\therefore P = 200 + 100e^{kt}$$

$P = 400, t = 3$

$$400 = 200 + 100e^{3k}$$

$$200 = 100e^{3k}$$

$$\therefore e^{3k} = 2$$


$$3k = \log 2$$

$$\therefore P = 200 + 100e^{\frac{t \log 2}{3}}$$

$$\text{When } t = 5, P = 200 + 100e^{\frac{5 \log 2}{3}} \\ \approx 517 \text{ people}$$

$$(ii) \frac{dP}{dt} = 100e^{\frac{t \log 2}{3}} \times \frac{\log 2}{3}$$

$$\text{When } t = 10, \frac{dP}{dt} = 233 \text{ people per year}$$



i)

$$\ddot{x} = 0 \quad | \quad \ddot{y} = -10$$

$$\dot{x} = c \quad | \quad \dot{y} = -10t + c$$

$$\text{When } t=0, \dot{x} = 25\sqrt{3} \quad | \quad \text{When } t=0, \dot{y} = 25$$

$$\therefore \dot{x} = 25\sqrt{3}t + k \quad | \quad \therefore 25 = 0 + c$$

$$\text{When } t=0, \dot{x} = 0 \quad | \quad (c = 25)$$

$$(k=0) \quad | \quad \dot{y} = -10t + 25$$

$$\therefore \underline{\underline{x = 25\sqrt{3}t}} \quad | \quad \underline{\underline{y = -5t^2 + 25t + k}}$$

$$\text{When } t=0, y=0 \quad | \quad (k=0)$$

$$\therefore \underline{\underline{y = -5t^2 + 25t}}$$

(ii) Max. height when  $\dot{y} = 0$  (i.e.  $y = \frac{dy}{dt}$ )

$$\therefore -10t + 25 = 0$$

$$\therefore t = 2.5 \text{ seconds}$$

$$(iii) y = -25 \Rightarrow -25 = -5t^2 + 25t$$

$$5t^2 - 25t - 25 = 0$$

$$t^2 - 5t - 5 = 0$$


$$t = \frac{5 \pm \sqrt{25+20}}{2}$$

$$= \frac{5 \pm \sqrt{45}}{2}$$

$$= \frac{5 + 3\sqrt{5}}{2} \quad (t > 0)$$

$$\dot{y} = -10 \left( \frac{5 + 3\sqrt{5}}{2} \right) + 25$$


$$\approx -33.54 \quad \text{and } \dot{x} = 25\sqrt{3}$$



$$\tan \alpha = \frac{33.54}{25\sqrt{3}}$$

$$\therefore \alpha \approx 38^\circ$$

7 a) dist travelled = total area under velocity graph.



$\therefore$  total dist. travelled = 13 metres

b) i)  $\dot{x} = \frac{dx}{dt} \left( \frac{1}{2} v^2 \right)$

$$= \frac{d}{dx} \left( 54 + 18x - \frac{9}{2} x^2 \right)$$

$$= 18 - 9x \text{ or } -9(x-2)$$

i)  $V_{\max}$  when  $x =$  centre of oscillation  
( $x=2$ )

$$\therefore v^2 = 108 + 72 - 36$$

$$= 144$$

$\therefore$  max. speed = 12 cm/s.

and time taken =  $\frac{1}{4}$  of period  
 $= \frac{1}{4} \times \frac{2\pi}{3}$   
 $= \frac{\pi}{6}$  seconds

iii)  $x = b + a \cos(\omega t + \alpha)$

$$\therefore x = 2 + 4 \cos 3t$$

c)  $\frac{dx}{dt} = 0.3, \frac{d\theta}{dt} = -0.05$

$$\tan \theta = \frac{2}{x}$$

$$\therefore x = \frac{2}{\tan \theta}$$

$$\frac{dx}{d\theta} = \frac{-2 \sec^2 \theta}{\tan^2 \theta}$$

$$= \frac{-2}{\sin^2 \theta}$$

$$\frac{dx}{d\theta} = \frac{dx}{dt} \times \frac{dt}{d\theta}$$

$$\frac{-2}{\sin^2 \theta} = 0.3 \times -0.05$$

$$= -0.015$$

$$\sin^2 \theta = \frac{1}{3}$$

$$\therefore \sin \theta = +\frac{1}{\sqrt{3}} \quad (\theta \text{ is acute})$$

OR  $\theta = \tan^{-1}\left(\frac{2}{x}\right)$

$$\frac{d\theta}{dx} = \frac{1}{1 + \left(\frac{2}{x}\right)^2} \times (-2x^{-2})$$

$$= \frac{1}{1 + \frac{4}{x^2}} \times \frac{-2}{x^2}$$

$$= \frac{-2}{x^2 + 4}$$

$$\frac{d\theta}{dx} = \frac{d\theta}{dt} \times \frac{dt}{dx}$$

$$\frac{-2}{x^2 + 4} = -0.05 \times \frac{1}{0.3}$$

$$= -\frac{1}{6}$$

$$12 = x^2 + 4$$

$$x^2 = 8$$

$$x = +\sqrt{8} \quad (x > 0)$$

$$\tan \theta = \frac{2}{\sqrt{8}}$$