


Hunters Hill High School  
**Extension 2, Mathematics**

Trial Examination, 2015



**Hunters Hill**  

---

**High School**

**General Instructions**

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- The marks for each question are shown on the paper
- Show all necessary working in questions 11-16

**Total Marks: 100**

**Section I** Page 3 – 7

**10 marks**

- Attempt Questions 1-10
- Allow about 15 minutes for this section


**Section II** Pages 8 – 14

**90 marks**

- Attempt Questions 11-16
- Allow about 2 hour 45 minutes for this section


**Section I MULTIPLE CHOICE:** Write the correct alternative on your writing paper.

1 The diagram shows the graph of the function  $y = f(x)$ .




Which of the following is the graph of  $y = f(x)^2$ ?


(A)




(B)



(C)



(D)




- 2 Consider the hyperbola with the equation  $\frac{x^2}{144} - \frac{y^2}{25} = 1$ .

What are the equations of the directrices?

- (A)  $x = \pm \frac{13}{144}$
- (B)  $x = \pm \frac{13}{25}$
- (C)  $x = \pm \frac{25}{13}$
- (D)  $x = \pm \frac{144}{13}$

- 3 Consider the Argand diagram below.



Which inequality could define the shaded area?

- (A)  $|z - i| \leq 2$  and  $0 \leq \arg(z - 1) \leq \frac{3\pi}{4}$
- (B)  $|z + i| \leq 2$  and  $0 \leq \arg(z - 1) \leq \frac{3\pi}{4}$
- (C)  $|z - i| \leq 2$  and  $0 \leq \arg(z - 1) \leq \frac{\pi}{4}$
- (D)  $|z + i| \leq 2$  and  $0 \leq \arg(z - 1) \leq \frac{\pi}{4}$

4 Which of the following is an expression for  $\int \frac{2}{x^2 + 4x + 13} dx$ ?


(A)  $\frac{1}{3} \tan^{-1} \frac{(x+2)}{3} + c$

(B)  $\frac{2}{3} \tan^{-1} \frac{(x+2)}{3} + c$

(C)  $\frac{1}{9} \tan^{-1} \frac{(x+2)}{9} + c$

(D)  $\frac{2}{9} \tan^{-1} \frac{(x+2)}{9} + c$

5 The point  $P$  lies on the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  where  $a > b > 0$ . The tangent at  $P$  meets the tangents at the ends of the major axis at  $R$  and  $T$ .



What is the equation of the tangent at  $P$ ?


(A)  $\frac{ax}{\cos \theta} - \frac{by}{\sin \theta} = a^2 - b^2$

(B)  $\frac{ax}{\sec \theta} + \frac{by}{\tan \theta} = a^2 + b^2$

(C)  $\frac{x}{a} \sec \theta - \frac{y}{b} \tan \theta = 1$

(D)  $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$

- 6 The region enclosed by the ellipse  $(x-1)^2 + \frac{y^2}{4} = 1$  is rotated about the  $y$  axis to form a solid.



What is the correct expression for volume of this solid using the method of slicing?

- (A)  $V = \int_{-2}^2 \pi \sqrt{1-y^2} dy$
- (B)  $V = \int_{-2}^2 2\pi \sqrt{1-y^2} dy$
- (C)  $V = \int_{-2}^2 \pi \sqrt{4-y^2} dy$
- (D)  $V = \int_{-2}^2 2\pi \sqrt{4-y^2} dy$
- 7 The polynomial equation  $x^3 - 3x^2 - x + 2 = 0$  has roots  $\alpha$ ,  $\beta$  and  $\gamma$ . Which of the following polynomial equations have roots  $\frac{1}{\alpha}$ ,  $\frac{1}{\beta}$  and  $\frac{1}{\gamma}$ ?
- (A)  $x^3 - x^2 - 3x + 1 = 0$
- (B)  $x^3 - 2x^2 - 3x + 1 = 0$
- (C)  $2x^3 - x^2 - 3x + 1 = 0$
- (D)  $2x^3 - 2x^2 - 3x + 1 = 0$

- 8 What is the derivative of  $\sin^{-1} x - \sqrt{1-x^2}$  ?
- (A)  $\frac{\sqrt{1+x}}{\sqrt{1-x}}$
- (B)  $\frac{\sqrt{1+x}}{1-x}$
- (C)  $\frac{1+x}{\sqrt{1-x}}$
- (D)  $\frac{1+x}{1-x}$
- 9 What are the values of real numbers  $p$  and  $q$  such that  $1-i$  is a root of the equation  $z^3 + pz + q = 0$  ?
- (A)  $p = -2$  and  $q = -4$
- (B)  $p = -2$  and  $q = 4$
- (C)  $p = 2$  and  $q = 4$
- (D)  $p = 2$  and  $q = 4$
- 10 A particle of mass  $m$  is moving in a straight line under the action of a force.

$$F = \frac{m}{x^3}(6-10x)$$

What of the following is an expression for its velocity in any position, if the particle starts from rest at  $x = 1$  ?

- (A)  $v = \pm \frac{1}{x} \sqrt{(-3+10x-7x^2)}$
- (B)  $v = \pm x \sqrt{(-3+10x-7x^2)}$
- (C)  $v = \pm \frac{1}{x} \sqrt{2(-3+10x-7x^2)}$
- (D)  $v = \pm x \sqrt{2(-3+10x-7x^2)}$

**Total Marks – 90****Section II****Attempt Questions 11-16****All Questions are of equal value****Marks****Question 11 (15 marks)** Begin a NEW sheet of paper.

(a) By using the method of partial fractions, show that

**4**

$$\int \frac{dx}{x^2 - 1} = \ln \sqrt{\frac{x-1}{x+1}} + c$$

(b) Use the substitution  $u = \cos x$  to evaluate**4**

$$\int_0^1 \sqrt{1-x^2} dx$$

(c) If  $I = \int e^x \sin x dx$ Find  $I$  using the method of integration by parts.**4**

(d) Evaluate

$$\int_0^{\frac{\pi}{2}} \cos x \sin^3 x dx$$

**3****End of Question 11**

**Question 12 (15 marks)** Begin a NEW sheet of paper.**Marks**(a) Given  $A = 3 - 4i$  and  $B = \sqrt{3} + i$ .(i) Find  $AB$  in  $x + iy$  form

1

(ii) Find  $\frac{A}{B}$  in  $x + iy$  form

1

(iii) Find  $\sqrt{A}$  in  $x + iy$  form

3

(iv) Find  $B$  in modulus- argument form

2

(v) Hence find  $B^4$  in  $x + iy$  form

1

(b) On separate Argand diagrams sketch the following loc

(i)  $2 \geq |z| \geq 1$ 

1

(ii)  $\frac{3\pi}{4} > \arg z > \frac{\pi}{4}$ 

1

(iii)  $3 \geq \operatorname{Re} Z \geq 0$  and  $3 \geq \operatorname{Im} Z \geq 1$ 

2

(c) On the Argand diagram shown OABC is a rectangle with the length OA being twice OC. OC represents the complex number  $x + iy$ .

Find the complex number represented by

(i) OA


1

(ii) OB

1

(iii) BC

1

**End of Question 12**




**Question 13 (15 marks)** Begin a NEW sheet of paper.

**Marks**

- (a) For the curve with equation  $x^2 + 3xy - y^2 = 13$ , determine the gradient of the tangent at the point (2, 3) on the curve. 4

(b)




The diagram above shows a solid which has the circle  $x^2 + y^2 = 16$  as its base. The cross-section perpendicular to the x axis is an equilateral triangle.

- (i) Show that the cross-sectional area of the equilateral triangle is given by  $A(x) = \sqrt{3}(16 - x^2)$  3
- (ii) Calculate the volume of the solid. 3

- (c) Shown are two circles centres  $H$  and  $K$  which touch at  $M$ .  $PQ$  and  $RM$  are common tangents.

COPY  
DIAGRAM



- (i) Show that quadrilaterals  $HPRM$  and  $MRQK$  are cyclic. 2
- (ii) Prove that triangles  $PRM$  and  $MKQ$  are similar. 3

**End of Question 13**

- Question 14 (15 marks)** Begin a NEW sheet of paper. **Marks**
- (a) The equation  $P(x) = x^3 + 3x^2 - 24x + k = 0$  has a double root.  
Find the possible values of  $k$ . **2**
- (b) The roots of  $x^3 + 3px + q = 0$  are  $\alpha, \beta$  and  $\gamma$ , (none of which are equal to 0).
- (i) Find the monic equation with roots  $\frac{\beta\gamma}{\alpha}, \frac{\alpha\gamma}{\beta}$  and  $\frac{\alpha\beta}{\gamma}$ , giving the coefficients in terms of  $p$  and  $q$ . **4**
- (ii) Deduce that if  $\gamma = \alpha\beta$  then  $(3p - q)^2 + q = 0$  **2**
- (c) The polynomial  $P(x)$  leaves a remainder of 9 when divided by  $(x - 2)$  and a remainder of 4 when divided by  $(x - 3)$ . Find the remainder when  $P(x)$  is divided by  $(x - 2)(x - 3)$ . **3**
- (d) (i) Solve  $Z^5 = 1$  over the complex field giving your answers in modulus-argument form. **2**
- (ii) Hence write  $Z^5 - 1$  as the product of linear and quadratic factors. **2**


**End of Question 14**

- Question 15 (15 marks)** Begin a NEW sheet of paper. **Marks**
- (a) A mass of 1kg is falling under gravity ( $g$ ) through a medium in which the resistance to the motion is proportional to the square of the velocity. ( $k$  is the constant of proportionality)
- (i) Draw a sketch showing all forces acting. 1
- (ii) Write an equation for the acceleration of this mass. 1
- (iii) Show that the mass reaches a terminal velocity 1
- given by  $v = \sqrt{\frac{g}{k}}$ .
- (iv) Show that the distance it has fallen when it reaches 4
- a velocity  $v$  m/s is given by  $x = \frac{1}{2k} \ln\left(\frac{g}{g - kv^2}\right)$
- (b) (i) Show that the recurrence (reduction) formula for 4
- $I_n = \int \sec^n x dx$
- is  $I_n = \frac{1}{n-1} \tan x \sec^{n-2} x + \frac{n-2}{n-1} I_{n-2}$
- (ii) Hence evaluate  $\int_0^{\frac{\pi}{4}} \sec^4 x dx$  2
- (c) A cubic equation in  $z$  has all real coefficients. If two of the roots are 3 and  $2 + i$  determine the equation. 2

**End of Question 15**

**Question 16 (15 marks)** Begin a NEW sheet of paper.**Marks**

(a)



The sketch shows the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  and the circle  $x^2 + y^2 = a^2$

with  $a, b \geq 0$ . T lies on the circle where  $\angle TOX = \theta$  and  $0 \leq \theta \leq \frac{\pi}{2}$ . The tangent at T meets OX at M and MP is perpendicular to OX with P on the hyperbola. Coordinates of T are  $(a \cos \theta, a \sin \theta)$

- (i) Find the equation of the tangent TM and hence the coordinates of M. 3
- (ii) Hence show that the coordinates of P are  $(a \sec \theta, b \tan \theta)$  1
- (iii) If  $Q(a \sec \beta, b \tan \beta)$  is another point on the hyperbola, where  $\theta + \beta = \frac{\pi}{2}$  and  $\theta \neq \frac{\pi}{4}$ , show that the equation of PQ is  $ay = b(\cos \theta + \sin \theta)x - ab$ . 3
- (iv) Every such chord PQ passes through a fixed point, find its coordinates. 2
- (v) Show that as  $\theta$  approaches  $\frac{\pi}{2}$ , PQ approaches a line parallel to an asymptote. 2

**Question 7 continues on page 9.**

**Question 16 continued.****Marks**

(b) (i) By letting  $Z = \cos \theta + i \sin \theta$  show that

**1**

$$Z^n + \frac{1}{Z_n} = 2 \cos n\theta .$$

(ii) Hence express  $\cos^4 \theta$  in terms of  $\cos n\theta$

**3**

**End of Question 16**

**End of Examination**

## STANDARD INTEGRALS

2

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln(x + \sqrt{x^2 - a^2}), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln(x + \sqrt{x^2 + a^2})$$

NOTE :  $\ln x = \log_e x, \quad x > 0$

**HHHS**

2015  
TRIAL HSC  
EXAMINATION

**Mathematics**  
Extension 2

**SOLUTIONS**

Section I	Trial HSC Examination- Mathematics Extension 2	2015
1		1 Mark: D
2	$b^2 = a^2(e^2 - 1) \qquad a^2 = 144 \text{ and } b^2 = 25.$ $25 = 144(e^2 - 1) \qquad a = 12 \qquad b = 5$ $(e^2 - 1) = \frac{25}{144} \text{ or } e^2 = \frac{169}{144} \text{ or } e = \frac{13}{12}$ <p>Equation of the directrices are <math>x = \pm \frac{a}{e} = \pm \frac{144}{13}</math>.</p>	1 Mark: D
3	<p><math> z - i  \leq 2</math> represents a region with a centre is <math>(0, 1)</math> and radius is less than or equal to 2.</p> <p><math>0 \leq \arg(z - 1) \leq \frac{3\pi}{4}</math> represents a region between angle 0 and <math>\frac{3\pi}{4}</math> whose vertex is <math>(1, 0)</math>, not including the vertex</p> <p><math> z - i  \leq 2</math> and <math>0 \leq \arg(z - 1) \leq \frac{3\pi}{4}</math></p>	1 Mark: A
4	$\int \frac{2}{x^2 + 4x + 13} dx = 2 \int \frac{dx}{(x+2)^2 + 3^2}$ $= \frac{2}{3} \tan^{-1} \frac{(x+2)}{3} + c$	1 Mark: B



5	<p>To find the equation of tangent through <math>P</math></p> $x = a \cos \theta \qquad y = b \sin \theta$ $\frac{dx}{d\theta} = -a \sin \theta \qquad \frac{dy}{d\theta} = b \cos \theta$ $\frac{dy}{dx} = \frac{dy}{d\theta} \times \frac{d\theta}{dx}$ $= b \cos \theta \times \frac{1}{-a \sin \theta} = \frac{-b \cos \theta}{a \sin \theta}$ <p>Equation of the tangent</p> $y - y_1 = m(x - x_1)$ $y - b \sin \theta = \frac{-b \cos \theta}{a \sin \theta} (x - a \cos \theta)$ $ay \sin \theta - ab \sin^2 \theta = -bx \cos \theta + ab \cos^2 \theta$ $bx \cos \theta + ay \sin \theta = ab(\sin^2 \theta + \cos^2 \theta)$ $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$	1 Mark: D
6	<p>The base is an annulus.</p> $A = \pi(r_2^2 - r_1^2)$ $= \pi(r_2 + r_1)(r_2 - r_1)$ $(r-1)^2 + \frac{y^2}{4} = 1$ $r^2 - 2r + \frac{y^2}{4} = 0$ $r = \frac{2 \pm \sqrt{(-2)^2 - 4 \times 1 \times \frac{y^2}{4}}}{2}$ $r = 1 \pm \sqrt{1 - \frac{y^2}{4}}$ <p>Therefore <math>r_2 + r_1 = 2</math> and <math>r_2 - r_1 = 2\sqrt{1 - \frac{y^2}{4}}</math></p> $V = \lim_{\delta y \rightarrow 0} \sum_{y=-2}^2 \pi \times 2 \times 2\sqrt{1 - \frac{y^2}{4}} \delta y$ $= \int_{-2}^2 2\pi \sqrt{4 - y^2} dy$	1 Mark: D

7	$x = \frac{1}{\alpha}, \frac{1}{\beta}, \frac{1}{\gamma}$ $\alpha = \frac{1}{x} \text{ satisfies } x^3 - 3x^2 - x + 2 = 0$ $\left(\frac{1}{x}\right)^3 - 3\left(\frac{1}{x}\right)^2 - \frac{1}{x} + 2 = 0$ $1 - 3x - x^2 + 2x^3 = 0$ $2x^3 - x^2 - 3x + 1 = 0$	1 Mark: C
8	$y = \sin^{-1} x - \sqrt{1-x^2}$ $\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}} - \frac{1}{2}(1-x^2)^{-\frac{1}{2}} \times -2x$ $= \frac{1}{\sqrt{1-x^2}} + \frac{x}{\sqrt{1-x^2}}$ $= \frac{1+x}{\sqrt{1-x^2}}$ $= \frac{1+x}{\sqrt{(1+x)(1-x)}}$ $= \frac{\sqrt{1+x}}{\sqrt{1-x}}$ <p>Result defined for <math>-1 \leq x \leq 1</math></p>	1 Mark: A
9	<p>Using the conjugate root theorem <math>1+i</math> and <math>1-i</math> are both roots of the equation <math>z^3 + pz + q = 0</math>.</p> <p><math>(1+i) + (1-i) + \alpha = 0</math> (sum of the roots)</p> $\alpha = -2$ <p><math>(1+i) \times (1-i) \times -2 = -q</math> (product of the roots)</p> $(1+1) \times -2 = -q$ $q = 4$ <p><math>(1+i)(1-i) + (1-i) - 2 + (1+i) - 2 = p</math></p> $p = -2$ <p>Therefore <math>p = -2</math> and <math>q = 4</math></p>	1 Mark: B




10	$F = \frac{m}{x^3}(6-10x)$ $ma = \frac{m}{x^3}(6-10x)$ $v \frac{dv}{dx} = \frac{6}{x^3} - \frac{10}{x^2}$ $\int v dv = \int \left( \frac{6}{x^3} - \frac{10}{x^2} \right) dx$ $\frac{1}{2} v^2 = \left( \frac{6x^{-2}}{-2} - \frac{10x^{-1}}{-1} \right) + c$ $\frac{1}{2} v^2 = \left( \frac{-3}{x^2} + \frac{10}{x} \right) + c$ <p>When <math>v = 0</math> and <math>x = 1</math></p> $\frac{1}{2} 0^2 = \left( \frac{-3}{1^2} + \frac{10}{1} \right) + c$ $c = -7$ $\frac{1}{2} v^2 = \left( \frac{-3}{x^2} + \frac{10}{x} \right) - 7$ $v^2 = \left( \frac{-6}{x^2} + \frac{20}{x} \right) - 14$ $= \frac{-6 + 20x - 14x^2}{x^2}$ $v = \pm \frac{1}{x} \sqrt{2(-3 + 10x - 7x^2)}$	1 Mark: C
----	--	-----------

Question 11	Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment
(a)	$\text{Let } \frac{A}{x+1} + \frac{B}{x-1} = \frac{1}{x^2-1}$ $\therefore A(x-1) + B(x+1) = 1$ $\text{If } x=1 \quad 2B=1$ $B = \frac{1}{2}$ $\text{If } x=-1 \quad 2A=-1$ $A = -\frac{1}{2}$ $\frac{dx}{x^2-1} = \frac{1}{2} \int \frac{1}{x-1} - \frac{1}{x+1} dx$ $\therefore \int = \frac{1}{2} [\ln(x-1) - \ln(x+1)] + C$ $= \frac{1}{2} \ln \left( \frac{x-1}{x+1} \right) + C$ $= \ln \sqrt{\frac{x-1}{x+1}} + c$	<p>1</p> <p>1</p> <p>1</p> <p>1</p>	<p>Any method to find A and B</p> <p>No mark for C</p> <p>Total = 4</p>

Question 11	Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment
(b)	<p>Let <math>x = \cos \theta</math>      When <math>x = 1</math>    <math>\theta = 0</math>  <math>dx = -\sin \theta d\theta</math>      <math>x = 0</math>    <math>\theta = \frac{\pi}{2}</math></p> $\therefore \int_0^1 \sqrt{1-x^2} dx = \int_{\frac{\pi}{2}}^0 \sqrt{1-\cos^2 \theta} (-\sin \theta d\theta)$ $= \int_{\frac{\pi}{2}}^0 -\sin^2 \theta d\theta$ $= \frac{1}{2} \int_{\frac{\pi}{2}}^0 (\cos 2\theta - 1) d\theta$ $= \frac{1}{2} \left[ \frac{1}{2} \sin 2\theta - \theta \right]_{\frac{\pi}{2}}^0$ $= \frac{1}{2} \left[ (0) - \left( 0 - \frac{\pi}{2} \right) \right]$ $= \frac{\pi}{4}$	<p>1</p> <p>1</p> <p>1</p> <p>1</p>	<p>Can use alternate methods ie <math>x = \sin \theta</math></p> <p>Total = 4</p>
(c)	<p><math>I = \int e^x \sin x dx</math></p> <p><math>= e^x \sin x - \int e^x \cos x dx</math></p> <p><math>= e^x \sin x - (e^x \cos x - \int e^x [-\sin x] dx)</math></p> <p><math>= e^x \sin x - e^x \cos x - I</math></p> <p><math>2I = e^x (\sin x - \cos x) + C</math></p> <p><math>\therefore I = \frac{e^x}{2} (\sin x - \cos x) + C</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>	<p>Total = 4</p>

Question 11		Trial HSC Examination- Mathematics Extension 2		2015	
Part	Solution	Marks	Comment		
(d)	$\int_0^{\frac{\pi}{2}} \cos x \sin^3 x \, dx = \left[ \frac{1}{4} \sin^4 x \right]_0^{\frac{\pi}{2}}$ $= \frac{1}{4} \left[ \sin^4 \left( \frac{\pi}{2} \right) - \sin^4 (0) \right]$ $= \frac{1}{4} [1 - 0]$ $= \frac{1}{4}$	1  1  1	May be done by a substitution of $u = \sin x$ $du = \cos x \, dx$   Total = 3		

Question 12		Trial HSC Examination- Mathematics Extension 2		2015	
Part	Solution	Marks	Comment		
(a) (i)	$AB = (3 - 4i)(\sqrt{3} + i)$ $= 3\sqrt{3} + 4 + (3 - 4\sqrt{3})i$	1			
(ii)	$\frac{A}{B} = \frac{(3 - 4i)(\sqrt{3} - i)}{(\sqrt{3} + i)(\sqrt{3} - i)}$ $= \frac{3\sqrt{3} - 4 + (-4\sqrt{3} - 3)i}{4}$	1			
(iii)	$x + iy = \sqrt{3 - 4i}(x + y\text{real})$ $\therefore x^2 - y^2 + 2xyi = 3 - 4i$ $\therefore x^2 - y^2 = 3 \dots \dots \dots (\alpha)$ $2xy = -4 \dots \dots \dots (\beta)$	1	Other methods okay		
	Squaring $x^4 - 2x^2y^2 + y^4 = 9, 4x^2y^2 = 16$ $x^4 + 2x^2y^2 + y^4 = 25$ $(x^2 + y^2)^2 = 25$ $x^2 + y^2 = 5 \dots \dots \dots (\gamma)$	1			
	Adding $(\alpha + \gamma) 2x^2 = 8$ $x = \pm 2$ $\therefore y = \mp 1$ $\therefore \sqrt{A} = \pm(2 - i)$	1			
(iv)	$B = \sqrt{3} + i$ $= 2\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right)$ $ B  = 2 \quad Arg B = \frac{\pi}{6}$ $= 2\left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}\right)$	1 for mod  1 for arg	Total = 2		

Question 12	Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment
(v)	$B^4 = 2^4 \left( \cos \frac{\pi}{6} + i \sin \frac{\pi}{6} \right)$ $= 16 \left( \cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right)$ $= -8 + 8\sqrt{3}i$	1	
(b) (i)		1	
(ii)		1	Dotted lines
(iii)		2	
(c) (i)	$OA = 2(-y + ix)$	1	
(ii)	$OB = OA + AB$ $= -2y + 2ix + x + iy$ $= (x - 2y) + (2x + y)i$	1	
(iii)	$BC = -OA$ $= 2y - 2xi$	1	



Question 13		Trial HSC Examination- Mathematics Extension 2		2015
Part	Solution	Marks	Comment	
(a)	$x^2 + 3xy - y^2 = 13$ $2x + 3y + 3x \cdot \frac{dy}{dx} - 2y \cdot \frac{dy}{dx} = 0$ $\frac{dy}{dx}(3x - 2y) = -(2x + 3y)$ $\frac{dy}{dx} = \frac{-(2x + 3y)}{(3x - 2y)}$ $\therefore \text{at } (2,3) \quad \frac{dy}{dx} = -\frac{(4+9)}{6-6}$ <p><math>\therefore</math> Gradient infinite</p> <p><math>\therefore</math> Tangent is vertical</p>	<p>2</p> <p>2</p>	<p>Total = 4</p>	

(b)  
(i)

$$a = b = 2y \text{ and } \theta = 60^\circ$$

$$\text{Area} = \frac{1}{2} a b \sin C$$

$$\text{Area} = \frac{1}{2} \times 2y \times 2y \times \sin 60^\circ$$

$$\text{Area} = \frac{1}{2} \times 4y^2 \times \frac{\sqrt{3}}{2}$$

$$\text{Area} = \sqrt{3} y^2$$

$$\text{Area} = \sqrt{3} (16 - x^2)$$

1

1

1

Total = 3

(ii)

$$V = \lim_{\delta x \rightarrow 0} \sum_{x=-4}^4 \sqrt{3}(16 - x^2) \delta x$$

$$V = \int_{-4}^4 \sqrt{3}(16 - x^2) dx$$

$$V = 2\sqrt{3} \int_0^4 (16 - x^2) dx$$

$$V = 2\sqrt{3} \left[ 16x - \frac{x^3}{3} \right]_0^4$$

$$V = \frac{128\sqrt{3}}{3} \text{ units}^3$$


1

1

1

Total = 3

(c)  
(i)



In  $HPRM$

$$\angle HPR = \angle RMH = 90^\circ \text{ (Angle between radius and tangent)}$$

$\therefore HPRM$  is a cyclic quadrilateral

(Opposite angles supplementary)

Similarly for  $MRQK$

1

1

Total = 2

(ii)

Join  $PM$  and  $QM$ .

in  $\triangle PRM$  and  $\triangle MQK$

$$\angle PRM = \angle MKQ \text{ (Exterior angle of cyclic quad } MRQK)$$

$$PR = RM \text{ (Tangent from external point)}$$

$$KM = KQ \text{ (radii)}$$

$\therefore$  Triangle isosceles.

$$\therefore \angle RPM = \angle RMP = \angle KMQ$$

$$\therefore \triangle PRM \cong \triangle MQK \text{ (Equiangular)}$$

1


1

Total = 2

Question 14		Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment	
(a)	$P(x) = x^3 + 3x^2 - 24x + k$ $P'(x) = 3x^2 + 6x - 24$ $P'(x) = 3(x^2 + 2x - 8)$ $P'(x) = 3(x+4)(x-2)$ Therefore $P(-4) = 0 \text{ and } P(2) = 0$ So $k = -80, 28$	1      1	Total = 2	
(b) (i)	From $x^3 + 3px + q = 0$ $\alpha + \beta + \gamma = 0, \alpha\beta + \alpha\gamma + \beta\gamma = 3p, \alpha\beta\gamma = -q$ $\therefore \frac{\beta\gamma}{\alpha} + \frac{\alpha\gamma}{\beta} + \frac{\alpha\beta}{\gamma} = \frac{(\beta\gamma)^2 + (\alpha\gamma)^2 + (\alpha\beta)^2}{\alpha\beta\gamma}$ $= \frac{(\beta\gamma + \alpha\gamma + \alpha\beta)^2 - 2(\alpha\beta\gamma^2 + \alpha\beta^2\gamma + \alpha^2\beta\gamma)}{\alpha\beta\gamma}$ $= \frac{(\beta\gamma + \alpha\gamma + \alpha\beta)^2 - 2\alpha\beta\gamma(\alpha + \beta + \gamma)}{\alpha\beta\gamma}$ $= \frac{(3p)^2 + 2q(0)}{-q} = -\frac{9p^2}{q}$ $\frac{\beta\gamma}{\alpha} \cdot \frac{\alpha\gamma}{\beta} + \frac{\alpha\gamma}{\beta} \cdot \frac{\alpha\beta}{\gamma} + \frac{\beta\gamma}{\alpha} \cdot \frac{\alpha\beta}{\gamma} = \gamma^2 + \alpha^2 + \beta^2$ $= (\gamma + \alpha + \beta)^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma)$ $= 0 - 2.3p$ $= -6p$ $\frac{\beta\gamma}{\alpha} \cdot \frac{\alpha\gamma}{\beta} \cdot \frac{\alpha\beta}{\gamma} = \alpha\beta\gamma$ $= -q$ $\therefore$ Required equation is $x^3 + \frac{9p^2}{q}x^2 - 6px + q = 0$	1          1  1  1	Total = 4	

Question 14		Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment	
(b) (ii)	<p>For <math>\gamma = \alpha\beta</math></p> $\frac{\alpha\beta}{\gamma} = 1 \text{ is a root}$ $\therefore 1 + \frac{9p^2}{q} - 6p + q = 0$ $q + 9p^2 - 6pq + q^2 = 0$ $\therefore (3p - q)^2 + q = 0$	1       1	Total = 2	
(c)	$P(x) = (x - 2)(x - 3)Q(x) + (ax + B)$ $P(2) = 9 \quad P(3) = 4$ <p>Therefore <math>2a + b = 9</math></p> $3a + b = 4$ $a = -5, b = 19$	1   1  1	Total = 3	
(d)	<p>Let <math>Z = cis\theta</math></p> $Z^5 = cis5\theta$ $\therefore \cos 5\theta + i \sin 5\theta = 1$ $\theta = 0, \frac{2\pi}{5}, \frac{4\pi}{5}, \frac{6\pi}{5}, \frac{8\pi}{5}$ $\text{i.e. } = 0, \frac{2\pi}{5}, \frac{4\pi}{5}, \frac{-4\pi}{5}, \frac{-2\pi}{5}$ $\therefore \text{Roots are } Z_1 = 1, \quad Z_2 = cis \frac{2\pi}{5}, \quad Z_3 = cis \frac{4\pi}{5}$ $Z_4 = cis\left(-\frac{4\pi}{5}\right) \quad Z_5 = cis\left(-\frac{2\pi}{5}\right)$ $= \bar{Z}_3 \quad = \bar{Z}_2$	1     1	Total = 2	

Question 14		Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment	
(e) ii)	$z^5 - 1$ $= (z - z_1)(z - z_2)(z - z_5)(z - z_3)(z - z_4)$ $= (z - z_1)(z^2 - (z_2 + z_5)z + z_2z_5)(z^2 - (z_3 + z_4)z + z_3z_4)$ $= (z - 1)(z^2 - 2z \cos \frac{2\pi}{5} + 1)(z^2 - 2z \cos \frac{4\pi}{5} + 1)$	1 1	Total = 2	

Question 15	Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment
(a) (i)		1	
(a) (ii)	$\ddot{x} = g - kv^2$	1	
(a) (iii)	<p>When <math>\ddot{x} = 0</math></p> $g = kv^2$ $\therefore v = \sqrt{\frac{g}{k}}$	1	
(a) (iv)	$\ddot{x} \frac{d}{dx} \left( \frac{1}{2} v^2 \right) = g - kv^2$ $= \frac{d}{dv} \left( \frac{1}{2} v^2 \right) \cdot \frac{dv}{dx} = g - kv^2$ $= v \cdot \frac{dv}{dx} = g - kv^2$ $\therefore \frac{dv}{dx} = \frac{g - kv^2}{v}$ $\frac{dx}{dv} = \frac{v}{g - kv^2}$ $\therefore x = -\frac{1}{2k} \ln(g - kv^2) + c$ <p>When <math>x = 0</math> <math>v = 0</math></p> $\therefore c = \frac{1}{2k} \ln g$ $\therefore x = -\frac{1}{2k} \ln(g - kv^2) + \frac{1}{2k} \ln g$ $= \frac{1}{2k} \ln \left( \frac{g}{g - kv^2} \right)$	<p>1</p> <p>1</p> <p>1</p> <p>1</p>	<p>Total = 4</p>



Question 15		Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment	
(b) (i)	$I_n = \int \sec^n x dx$ $= \int \sec^{n-2} x \cdot \sec^2 x dx$ $= \tan x \cdot \sec^{n-2} x - \int (n-2) \sec^{n-3} x \cdot \tan x \cdot \sec x \tan x dx$ $= \tan x \sec^{n-2} x - (n-2) \int \sec^{n-2} x \cdot \tan^2 x dx$ $= \tan x \sec^{n-2} x - (n-2) \int \sec^{n-2} x \cdot (\sec^2 x - 1) dx$ $= \tan x \sec^{n-2} x - (n-2) \int (\sec^n x - \sec^{n-2} x) dx$ $= \tan x \sec^{n-2} x - (n-2) I_n + (n-2) I_{n-2}$ $I_n + (n-2) I_n = (n-1) I_n = \tan x \sec^{n-2} x + (n-2) I_{n-2}$ $I_n = \frac{1}{n-1} \tan x \sec^{n-2} x + \frac{n-2}{n-1} I_{n-2}$	1  1  1  1		
			Total = 4	
(b) (ii)	$\int_0^{\frac{\pi}{4}} \sec^4 x dx = I_4 = \left[ \frac{1}{3} \tan x \sec^2 x + \frac{2}{3} \int \sec^2 x \right]_0^{\frac{\pi}{4}}$ $= \left[ \frac{1}{3} \tan x \sec^2 x + \frac{2}{3} \tan x \right]_0^{\frac{\pi}{4}}$ $= \left( \frac{1}{3} \tan \frac{\pi}{4} \sec^2 \frac{\pi}{4} + \frac{2}{3} \tan \frac{\pi}{4} \right) - \left( \frac{1}{3} \tan 0 \sec^2 0 + \frac{2}{3} \tan 0 \right)$ $= \left( \frac{1}{3} \times 1 \times 2 + \frac{2}{3} \times 1 \right) - 0$ $= \frac{4}{3}$	1     1		
			Total = 2	

Question 15		Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment	
(c)	<p>If one root is <math>2 + i</math>, another is <math>2 - i</math></p> $\therefore (z - (2 + i))(z - (2 - i))(z - 3) = 0$ $(z - 2 - i)(z - 2 + i)(z - 3) = 0$ $(z^2 - 4z + 5)(z - 3) = 0$ $z^3 - 7z^2 + 17z - 15 = 0$	1		
		1	Total = 2	

Question 16	Trial HSC Examination- Mathematics Extension 2	2015	
Part	Solution	Marks	Comment
(a) (i)	<p>Coordinates of T are <math>(a \cos \theta, a \sin \theta)</math></p> $x^2 + y^2 = a^2$ $\therefore 2x + 2y \frac{dy}{dx} = 0$ $\therefore \frac{dy}{dx} = -\frac{x}{y}$ <p>at T <math>\frac{dy}{dx} = -\frac{\cos \theta}{\sin \theta}</math></p> <p><math>\therefore</math> Equation <math>TM</math></p> $y - a \sin \theta = -\frac{\cos \theta}{\sin \theta}(x - a \cos \theta)$ $x \cos \theta + y \sin \theta = a$ <p>When <math>y = 0</math> <math>x = a \sec \theta</math></p> <p><math>\therefore</math> Coordinates <math>M</math> are <math>(a \sec \theta, 0)</math></p>	<p>1</p> <p>1</p> <p>1</p>	<p>Total = 3</p>
(a) (ii)	<p>On <math>\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1</math> when <math>x = a \sec \theta</math></p> $a^2 \frac{\sec^2 \theta}{a^2} - \frac{y^2}{b^2} = 1$ $\frac{y^2}{b^2} = \sec^2 \theta - 1$ $\frac{y^2}{b^2} = \tan^2 \theta$ <p><math>\therefore y = b \tan \theta</math></p> <p>Coordinates of <math>P</math> are <math>(a \sec \theta, b \tan \theta)</math></p>	<p>1</p>	

<p>(a) iii)</p>	$a \sec \beta = a \sec\left(\frac{\pi}{2} - \theta\right)$ $= a \operatorname{cosec} \theta$ $b \tan \beta = b \tan\left(\frac{\pi}{2} - \theta\right)$ $= b \cot \theta$ $\text{Gradient } PQ = \frac{b \cot \theta - b \tan \theta}{a \operatorname{cosec} \theta - a \sec \theta}$ $= \frac{b}{a} \left\{ \frac{\frac{\cos \theta}{\sin \theta} - \frac{\sin \theta}{\cos \theta}}{\frac{1}{\sin \theta} - \frac{1}{\cos \theta}} \right\}$ $= \frac{b}{a} \left\{ \frac{\frac{\cos^2 \theta - \sin^2 \theta}{\sin \theta \cos \theta}}{\frac{\cos \theta - \sin \theta}{\sin \theta \cos \theta}} \right\}$ $= \frac{b}{a} \left( \frac{\cos^2 \theta - \sin^2 \theta}{\cos \theta - \sin \theta} \right)$ $= \frac{b}{a} (\cos \theta + \sin \theta)$ <p><math>\therefore</math> Equation <math>PQ</math> is</p> $y - b \tan \theta = \frac{b}{a} (\cos \theta + \sin \theta) (x - a \sec \theta)$ $y - b \tan \theta = \frac{b}{a} (\cos \theta + \sin \theta) x - b \cos \theta \sec \theta - b \sin \theta \sec \theta$ $y - b \tan \theta = \frac{b}{a} (\cos \theta + \sin \theta) x - b - b \tan \theta$ $y = \frac{b}{a} (\cos \theta + \sin \theta) x - b$ $ay = b (\cos \theta + \sin \theta) x - ab$	<p>1</p> <p>1</p> <p>1</p>	<p>Total = 3</p>
<p>(a) iv)</p>	<p>All of the lines have the same <math>y</math> intercept. i.e. <math>y = -b</math> <math>\therefore</math> The fixed point is the intercept <math>(0, -b)</math></p>	<p>1</p> <p>1</p>	<p>Total = 2</p>

<p>(a) (v)</p>	<p>Equations of Asymptotes are <math>y = \pm \frac{b}{a}x</math></p> <p>Gradients of asymptotes are <math>m = \pm \frac{b}{a}</math></p> <p>As <math>\theta \rightarrow \frac{\pi}{2}</math> the equation of <math>PQ \rightarrow ay = b(0+1)x - ab</math></p> <p><math>\therefore</math> Gradient of <math>PQ \rightarrow \frac{b}{a}</math></p> <p><math>\therefore PQ</math> approaches a line which is parallel to an asymptote.</p>	<p>1</p> <p>1</p>	<p>Total = 2</p>
<p>(b) i)</p>	<p><math>z = \cos \theta + i \sin \theta</math></p> <p><math>\frac{1}{z} = z^{-1} = \cos \theta - i \sin \theta</math></p> <p>By De Moivres Theorem</p> <p><math>z^n = \cos n\theta + i \sin n\theta</math></p> <p><math>z^{-n} = \cos n\theta - i \sin n\theta</math></p> <p><math>z^n + z^{-n} = \cos n\theta + i \sin n\theta + \cos n\theta - i \sin n\theta</math></p> <p><math>z^n + \frac{1}{z^n} = 2 \cos n\theta</math></p>	<p>1</p>	
<p>ii)</p>	<p><math>\left(z + \frac{1}{z}\right)^4 = z^4 + 4z^3 \frac{1}{z} + 6z^2 \frac{1}{z^2} + 4z \frac{1}{z^3} + \frac{1}{z^4}</math></p> <p><math>= \left(z^4 + \frac{1}{z^4}\right) + 4\left(z^2 + \frac{1}{z^2}\right) + 6</math></p> <p><math>(2 \cos \theta)^4 = 2 \cos 4\theta + 4(2 \cos 2\theta) + 6</math></p> <p><math>2^4 \cos^4 \theta = 2 \cos 4\theta + 8 \cos 2\theta + 6</math></p> <p><math>\cos^4 \theta = \frac{1}{8} \cos 4\theta + \frac{1}{2} \cos 2\theta + \frac{3}{8}</math></p>	<p>1</p> <p>1</p> <p>1</p>	<p>Total = 3</p>