

JRAHS Ext 2 Trial 2002

QUESTION 1 (15 MARKS) Start a new page

a) (i) Given $f(x)$ is an odd function, show that $\int_{-a}^a f(x)dx=0$ using $x = -t$.

(ii) Hence, evaluate $\int_{-2}^2 x^4(1+\sin^3 x)dx$.

b) (i) Let $I_n = \int_1^e x(\ln x)^n dx, n = 0,1,2,3,\dots$

Using integration by parts, show that $I_n = \frac{e^2}{2} - \frac{n}{2}I_{n-1}, n = 1,2,3,\dots$

(ii) The area bounded by the curve $y = \sqrt{x}(\ln x)^2, x \geq 1$, the x -axis and the line $x = e$ is rotated about the x -axis through 2π radians.

Find the exact volume of the solid of revolution so formed.

c) (i) Sketch the curve $f(x) = \frac{x^2 - x - 6}{x - 1}$.

(ii) Hence, sketch the graph of $y^2 = f(x)$.

QUESTION 2 (15 MARKS) Start a new page

a) Find the following:

(i) $\int \frac{e^{-x}}{1+e^x} dx$.

(ii) $\int \frac{x}{\sqrt{1-2x-x^2}} dx$.

b) Evaluate in exact form:

(i) $\int_0^{\frac{\pi}{6}} \frac{d\theta}{9-8\cos^2 \theta}$ using the substitution $t = \tan \theta$.

(ii) $\int_0^{\frac{\pi}{6}} \sec^3 2\theta d\theta$.

c) Using the method of cylindrical shells, find the volume generated by revolving the area bounded by the lines $x = \pm 2$ and the hyperbola

$$\frac{y^2}{9} - \frac{x^2}{4} = 1 \text{ about the } y\text{-axis.}$$

QUESTION 3 (15 MARKS) Start a new page

- a) Let α, β, γ be the roots of the cubic equation $x^3 + px^2 + q = 0$, where p and q are real. The equation $x^3 + ax^2 + bx + c = 0$ has roots α^2, β^2 and γ^2 . Find a, b, c as functions of p and q .
- b) (i) Find the complex square roots of $5 - 12i$, expressing your answer in the form $a+bi$, where a and b are real.
- (ii) Hence, solve the equation: $z^2 + 4z - 1 + 12i = 0$
- c) Given that $z = \cos \theta + i \sin \theta$,
- (i) Show that $z^n + z^{-n} = 2 \cos n\theta$ using De Moivre's Theorem.
- (ii) Hence, solve the equation: $2z^4 - z^3 + 3z^2 - z + 2 = 0$.

QUESTION 4 (15 MARKS) Start a new page

- a) The sequence U_n , is defined such that
 $U_{n+2} = 4U_{n+1} - U_n$, $n \geq 1$ and $U_1 = 2, U_2 = 4$.
Prove by mathematical induction that:
 $U_n = (2 + \sqrt{3})^{n-1} + (2 - \sqrt{3})^{n-1}$.
- b) $P(\cos \theta, \sin \theta)$ and $Q(\sec \theta, \tan \theta)$ lie on the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$, respectively as shown.

M and N are the feet of the perpendicular from P and Q respectively to the x -axis. $0 < \theta < \frac{\pi}{2}$, and QP meets the x -axis at K . A is the point $(a,0)$.

- (i) Given $\triangle KPM \parallel \triangle KQN$, show that $\frac{KM}{KN} = \cos \theta$.
- (ii) Hence, show that K has coordinates $(-a,0)$.
- (iii) **Show** that the tangent to the ellipse at P has equation $\frac{x \cos \theta}{a} + \frac{y \sin \theta}{b} = 1$, **and** deduce it passes through N .
- (iv) **Given** that the tangent to the hyperbola at Q has equation $\frac{x \sec \theta}{a} - \frac{y \tan \theta}{b} = 1$, show that the tangent passes through M .
- (v) Show that the tangents PN , QM and the common tangent at A are concurrent. Find the point of concurrence.

QUESTION 5 (15 MARKS) Start a new page

- a) (i) If $\omega = i - 1$, evaluate the following points $\bar{\omega}$, $i\omega$, $\frac{1}{\omega}$
- (ii) Hence, indicate ω , $\bar{\omega}$, $i\omega$, $\frac{1}{\omega}$ on the Argand diagram.
- b) Sketch the region R in the Argand diagram consisting of the points z for which:
- $$|\arg z| < \frac{\pi}{3}, \quad z + \bar{z} < 4, \quad |z| > 2$$
- c) On the Argand diagram, P represents the complex number z , and R the number $\frac{1}{z}$. A square $PQRS$ is drawn in the plane with PR as a diagonal. If P lies on the circle $|z| = 2$,
- (i) Prove that Q will lie on the ellipse whose equation has the form $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$;
- (ii) Hence, specify the numerical values for a and b .

QUESTION 6 (15 MARKS) Start a new page

a) The base of a solid is the region bounded by the line $y = 2x$ and the parabola $y = 4x - x^2$ and the x -axis. Cross-sections parallel to the x -axis are right-angled isosceles triangles with the hypotenuse in the base of the solid.

(i) Show that the volume of the solid is given by:

$$V = \frac{1}{16} \int_0^4 (4 + 2\sqrt{4-y} - y)^2 dy$$

(ii) Hence, find the exact volume of the solid.

b) A particle of mass, m kg, moves in a straight line with velocity v metres/second, under a constant force, P Newtons, and a resistance, R Newtons. Initially, the particle has a speed v_0 metres/second. If $R = 5 + 3v$ and $P = 10$.

(i) Show that velocity, $v = \frac{5}{3}(1 - e^{-\frac{3t}{m}}) + v_0 e^{-\frac{3t}{m}}$.

(ii) Find the terminal velocity of the particle.

(iii) When the particle accelerates from v_0 to v_1 , show that the distance travelled, x metres, is given by :

$$x = \frac{m}{9} \left[3(v_0 - v_1) + 5 \ln \left(\frac{5 - 3v_0}{5 - 3v_1} \right) \right].$$

QUESTION 7 (15 MARKS) Start a new page

a) With respect to the x and y axis, the line $x = 1$ is a directrix, and the point $(2,0)$ is a focus of a conic of eccentricity $\sqrt{2}$.

Find the equation of the conic, and sketch the curve indicating its asymptotes, foci, and directrices.

- b) A circular cone of semi vertical angle θ is fixed with its vertex upwards as shown. A particle P , of mass $2m$ kg, is attached to the vertex V by a light inextensible string of length $2a$ metres.

The particle P rotates with uniform angular velocity ω radians/second in a horizontal circle on the outside surface of the cone and in contact with it.

- (i) Find the tension (T) in the string, in Newtons.
- (ii) Find the normal force (N) on P , in Newtons.
- (iii) Show that, for the particle to remain in uniform circular

motion on the surface of the cone, then $\omega < \left[\frac{g}{2a \cos \theta} \right]^{\frac{1}{2}}$

where g is acceleration due to gravity.

- c) A car of mass, m kg, with speed v metres/second travels around a circular track of radius R metres, inclined at angle θ to the horizontal and g is the acceleration due to gravity.

- (i) Show that if there is a tendency for the car to slip that

$$\tan \theta = \frac{v^2}{gR}.$$

- (ii) If the speed of the car is now halved, prove that the sideways frictional force F , on the wheels, exerted by the track is given:

$$F = \frac{3mgv^2}{4\sqrt{v^4 + g^2 R^2}}.$$

QUESTION 8 (15 MARKS) Start a new page

- a) Two particles of mass 4kg and 6kg are attached at either end of a light inextensible string of length 7 metres, which passes through a small vertical frictionless ring R . The heavier particle A hangs vertically at a distance of 4 metres below the ring while the other particle B describes a horizontal circle whose centre is O . Let θ be the acute angle which particle B makes with the vertical.

Find:

- (i) The distance OR and the radius OB , of the horizontal circle.
 - (iii) The angular velocity of B about O in revolutions/minute to 2 decimal places (use $g = 9.8 \text{ m/s}^2$).
- b) In the diagram below, EC and ED are perpendicular to BA and AC at G and H respectively. The lines AC and BD meet at I . Let $\angle ECA = \alpha$.

- (i) Copy the diagram, then show that $FGCH$ is a cyclic quadrilateral.
- (ii) Prove $\triangle BCD$ is isosceles.
- (iii) Prove $\triangle CID \parallel \triangle CDA$.
- (iv) Given that $\triangle CIB \parallel \triangle CBA$ and $AB + AD = 2BC$, prove that $2CI = BD$.