


Student Number:	
Class:	



**TRIAL HIGHER SCHOOL CERTIFICATE  
EXAMINATION 2015**

**MATHEMATICS  
EXTENSION 2**

**General Instructions:**

- Reading Time: 5 minutes.
- Working Time: 3 hours.
- Write in black pen.
- Board approved calculators & templates may be used
- A Standard Integral Sheet is provided.
- In Question 11 - 16, show all relevant mathematical reasoning and/or calculations.
- Marks may not be awarded for careless or badly arranged working.

**Total Marks 100**

**Section I: 10 marks**

- Attempt Question 1 – 10.
- Answer on the Multiple Choice answer sheet provided.
- Allow about 15 minutes for this section.

**Section II: 90 Marks**


- Attempt Question 11 - 16
- Answer on lined paper provided. Start a new page for each new question.
- Allow about 2 hours & 45 minutes for this section.

**The answers to all questions are to be returned in separate *stapled* bundles clearly labelled Question 11, Question 12, etc. Each question must show your Candidate Number.**

## Section 1 (10 Marks)

Attempt questions 1 – 10. Use the Multiple Choice answer sheet supplied.

1. A circle with centre  $(0,2)$  and radius 3 units is shown below on an Argand diagram.



Which of the following inequalities represents the shaded region ?

- A)  $\text{Im}(z) \leq 0$  and  $|z - 2| \leq 3$       B)  $\text{Re}(z) \leq 0$  and  $|z - 2| \leq 3$   
C)  $\text{Im}(z) \leq 0$  and  $|z - 2i| \leq 3$       D)  $\text{Re}(z) \leq 0$  and  $|z - 2i| \leq 3$
2. The equation  $x^2 - xy + y^2 = 3$  defines  $y$  implicitly in terms of  $x$ .

The expression for  $\frac{dy}{dx}$  is :

- A)  $\frac{y-2x+3}{2y-x}$       B)  $\frac{y-2x+3}{x-2y}$       C)  $\frac{y-2x}{2y-x}$       D)  $\frac{y-2x}{x-2y}$
3. In how many ways can five letters be chosen from the letters ARRANGE ?
- A) 9      B) 10      C) 12      D) 21
4. Consider a polynomial  $P(x)$  of degree 3.

Two real numbers  $a$  and  $b$  are such that :

$$a < b$$


$$P(a) > P(b) > 0$$

$$P'(a) = P'(b) = 0$$

The polynomial has :

- A) 3 real zeroes      B) 1 real zero  $\gamma$  such that  $\gamma < a$   
C) 1 real zero  $\gamma$  such that  $a < \gamma < b$       D) 1 real zero  $\gamma$  such that  $\gamma > b$





5. The graph shows a part of the hyperbola  $x = ct, y = c/t$ .



Which pair of parametric equations precisely describes the graph as shown?

- A)  $x = c(t^2 + 1), y = c/(t^2 + 1)$     B)  $x = c(1 - t^2), y = c/(1 - t^2)$   
 C)  $x = c\sqrt{1 - t^2}, y = c/\sqrt{1 - t^2}$     D)  $x = c \sin t, y = c/\sin t$
6. The region enclosed by  $y = x^3, y = 0$  and  $x = 2$  is rotated about the  $y$ -axis to produce a solid. What is the volume of that solid ?  
 A)  $\frac{8\pi}{5}$  units<sup>3</sup>    B)  $\frac{32\pi}{5}$  units<sup>3</sup>    C)  $\frac{64\pi}{5}$  units<sup>3</sup>    D)  $\frac{16\pi}{5}$  units<sup>3</sup>
7. The equation  $|z - 4| + |z + 4| = 10$  defines an ellipse. What is the length of the semi minor axis ?  
 A)  $2\frac{2}{5}$     B) 3    C) 4    D) 5
8. A particle is moving in a circle of radius 80cm with a linear speed of  $4\pi$  m/s. It has a constant angular speed (in revolutions per minute) of :  
 A)  $3/8$  rpm    B)  $3/2$  rpm    C)  $37\frac{1}{2}$  rpm    D) 150 rpm

9. Which of the diagrams below best represents the graph of  $y = \sin^{-1}(\sin x)$  ?



10. A particle of mass  $m$  moves in a straight line under the action of a resultant force  $F$  where  $F=F(x)$ . Given that the velocity  $v$  is  $v_0$  when the position  $x$  is  $x_0$ , and that  $v$  is  $v_1$  when  $x$  is  $x_1$ , it follows that  $|v_1| =$

A)  $\sqrt{\frac{2}{m} \int_{x_0}^{x_1} \sqrt{F(x)} dx} + v_0$

B)  $\sqrt{2} \int_{\sqrt{x_0}}^{\sqrt{x_1}} F(x) dx + v_0$

C)  $\sqrt{\frac{2}{m} \int_{x_0}^{x_1} \sqrt{F(x)} dx + (v_0)^2}$

D)  $\sqrt{\frac{2}{m} \int_{x_0}^{x_1} \{F(x) + (v_0)^2\} dx}$

End of Section 1


## Section 2 (90 Marks)

Attempt all questions 11 – 16. Start each question on a new sheet of paper.

**QUESTION 11** (Start a new sheet of paper)


Marks

- a) Find  $\int \frac{1}{\sqrt{5+4x-x^2}} dx$  2
- b) i) Find the real numbers  $a$  and  $b$  such that
- $$\frac{3x^2-3x+7}{(x-2)(x^2+9)} \equiv \frac{a}{(x-2)} + \frac{bx+1}{(x^2+9)}$$
- 2
- ii) Find  $\int \frac{3x^2-3x+7}{(x-2)(x^2+9)} dx$  3
- c) i) Prove that  $\int_0^a f(x)dx = \int_0^a f(a-x)dx$  1
- ii) Hence find the value of  $\int_0^2 x(2-x)^5 dx$ . 2
- d) Consider the equation  $z^2 + az + (2+i) = 0$ .
- i) Find the complex number  $a$ , given that  $i$  is a root of the equation. 1
- ii) Also write down the second root of the equation. 1
- e) In the Argand diagram,  $OABC$  is a rectangle, where  $3OC = OA$ . The vertex at  $A$  represents the complex number  $w$ .
- i) What complex number corresponds to the vertex  $C$ ? 1
- ii) What complex number corresponds to the point of intersection  $D$  of the diagonals  $OB$  and  $AC$ ? 2




**QUESTION 12** (Start a new sheet of paper)

- a) The diagram below shows the graph of  $y = f(x)$ .



Draw separate one-third page sketches of the following graphs:


- |      |                      |   |
|------|----------------------|---|
| i)   | $y = \frac{1}{f(x)}$ | 2 |
| ii)  | $y^2 = f(x)$         | 2 |
| iii) | $y = (f(x))^2$       | 2 |
- b) i) The polynomial  $P(x) = x^4 - 6x^3 + 13x^2 - ax - b$  has two double zeroes. Find  $a$  and  $b$ . 3
- ii) Hence determine the equation of the line which touches the curve  $y = x^4 - 6x^3 + 13x^2$  at two distinct points. 1
- c) The base of a solid  $S$  is the region in the  $xy$  plane enclosed by the parabola  $y^2 = 4x$  and the line  $x = 4$ . Each cross section perpendicular to the  $x$  axis is a semi-ellipse with the major axis in the  $xy$  plane and with the major and minor axes in the ratio  $a:b$ .



- |      |   |   |
|------|---|---|
| i)   | Assuming that the area of an ellipse with semi-axes $A$ and $B$ is $\pi AB$ , show that the area of the semi-ellipse shown at $x = h$ is $2\pi hb/a$ .  | 1 |
| ii)  | Find the volume of the solid $S$ .  | 3 |
| iii) | The solid $T$ is obtained by rotating the region enclosed by the parabola and the line $x = 4$ about the $x$ axis. Using (with a little care) your result from (ii), or otherwise, find the volume of $T$ . | 1 |

**QUESTION 13** (Start a new sheet of paper)

a)




In the diagram above  $ABCD$  is a cyclic quadrilateral whose diagonals are perpendicular and intersect at  $Q$ . Let  $M$  be the midpoint of  $BC$  and suppose that  $MQ$  produced meets  $AD$  at  $N$ . Let  $\angle QBC = \alpha$ .

- i) Explain why  $BM = QM$ . 1
- ii) Prove that  $MN \perp AD$ . 3

- b) At a point on a railway line where the radius of curvature is 200m, the track is designed so that a train travelling at 50 km/hr exerts no lateral force on the track. What lateral force would a stationary locomotive, of mass 40 tonnes, exert on the track at this point? (Take  $g$  to be 9.81m/s/s.) 4

- c) The diagram shows the side view of a vertical cylindrical water cooler of constant cross sectional area  $A$ . Water drains through a hole at the bottom of the cooler. It is known that the volume of water decreases at a rate given by  $\frac{dv}{dt} = -k\sqrt{y}$  where  $k$  is a positive constant and  $y$  is the depth of the water. Initially the cooler is full and it would take  $T$  seconds to drain completely.



- i) Show that  $\frac{dy}{dt} = -\frac{k}{A}\sqrt{y}$  1
- ii) Show that  $y = y_0 \left(1 - \frac{t}{T}\right)^2$  for  $0 \leq t \leq T$ . 4
- iii) If it takes 10 seconds for half the water to drain, evaluate  $T$ . 2

QUESTION 14 (Start a new sheet of paper)

- a) A particle moves in Simple Harmonic Motion, the period being 2 seconds and the amplitude 3 metres. Find the maximum speed and the maximum acceleration during the motion. 2

- b) i) Use de Moivre's Theorem to show that

$$(\cot\theta + i)^n + (\cot\theta - i)^n = \frac{2\cos n\theta}{\sin^n\theta} \quad 2$$

- ii) Show that the equation  $(x + i)^5 + (x - i)^5 = 0$  has roots  $0, \pm\cot\frac{\pi}{10}, \pm\cot\frac{3\pi}{10}$ . 2

- iii) Hence show that the equation  $x^4 - 10x^2 + 5 = 0$  has roots  $\pm\cot\frac{\pi}{10}, \pm\cot\frac{3\pi}{10}$ . 2

- iv) Hence show that  $\cot\frac{\pi}{10} = \sqrt{5 + 2\sqrt{5}}$  2

- c) A ball is projected so as to just clear two walls. The first wall is at a height  $b$  at a horizontal distance  $a$  from the point of projection and the second is of height  $a$  at a horizontal distance  $b$  from the point of projection. ( $b > a$ )

It may be assumed that, if the ball is projected from the origin with velocity  $V$  at an angle  $\alpha$  to the horizontal ( $x$  axis), then the equation of the path is given by

$$y = x\tan\alpha - \frac{gx^2\sec^2\alpha}{2V^2}$$

- i) Show that the range on the horizontal plane is  $\frac{a^2+ab+b^2}{a+b}$ . 3
- ii) Show that the angle of projection must exceed  $\tan^{-1}3$ . 2



**QUESTION 15** (Start a new sheet of paper)


- a) The diagram shows a segment of the circle  $x^2 + y^2 = r^2$  which is rotated about the  $y$  axis to form a collar. This collar is thus a sphere with a symmetrical hole through it. Let the hole be of height  $2h$  as shown.

Use the method of cylindrical shells to show that the volume of the material in the collar is given by the integral

$$4\pi \int_{\sqrt{r^2-h^2}}^r x\sqrt{r^2-x^2} dx$$

Evaluate the integral to show that the volume of material in the collar is a function of  $h$  only and independent of  $r$ .

4




- b) Consider the equation  $z^5 + 32 = 0$ .
- Write down the roots of this equation in modulus argument form. 2
  - Illustrate these roots on an Argand diagram. 1
  - If the points  $A, B, C, D$  and  $E$  cyclically represent these roots, find the area of triangle  $ACD$ . (Give your answer to 2 decimal places) 3

Question 15 is continued on the next page ....

Question 15 (continued)

- c) i) Consider the diagram below. The tangent at  $T(2t, \frac{2}{t})$  to the rectangular hyperbola  $xy = 4$  meets the ellipse  $x^2 + 4y^2 = 4$  at  $P$  and  $Q$ . The tangents to the ellipse at  $P$  and  $Q$  intersect at  $M$ . Find the equation of the locus of  $M$ . (You may use standard forms of tangents and such things without proof.) 3
- ii) Describe briefly the locus and any restrictions that it may have. 2



QUESTION 16 (Start a new sheet of paper)

- a) A box contains  $n$  jellybeans, some white and some black. Alan and Betty take turns picking a jellybean from the box, without looking, until the box is empty. Alan picks first.
- i) If there is 1 black and  $n - 1$  white jellybeans and  $n$  is odd, find the probability that Alan picks the black jellybean. 1
- ii) If there are 2 black and  $n - 2$  white jellybeans and  $n$  is even, find the probability that Alan is the first to pick a black jellybean. 2
- iii) If there are 2 black and  $n - 2$  white jellybeans and  $n$  is odd, find the probability that Alan is the first to pick a black jellybean. 2

Question 16 is continued on the next page....

Question 16 (continued)


b) Consider the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ .

i) Show that lines with equations  $y = mx \pm \sqrt{a^2m^2 + b^2}$  are always tangent to the ellipse for any real value of  $m$ . 2

ii) From any point external to the ellipse, two tangents may be drawn. By considering the above as a quadratic equation in  $m$ , or otherwise, show that the locus of points where the two tangents are perpendicular to each other is a circle with equation  $x^2 + y^2 = a^2 + b^2$ . 2

iii) Show that the area of the ellipse's circumscribing rectangle of which  $y = mx \pm \sqrt{a^2m^2 + b^2}$  are two parallel sides, is given by

$$A = \frac{4}{(1+m^2)} \sqrt{(a^2 + m^2b^2)(a^2m^2 + b^2)} \quad 3$$



You are given that this formula for the area can be rearranged to give the following form for the square of the area:  $A^2 = 16a^2b^2 + \frac{16(a^2 - b^2)^2}{(m + \frac{1}{m})^2}$

iv) Show that, for any  $m > 0$ ,  $m + \frac{1}{m} \geq 2$ . 1

v) Hence, or otherwise, find the maximum and minimum areas of rectangles which circumscribe the ellipse. 2

END OF EXAM


JRAHS EXTENSION 2 MATHS TRIAL  
2015 (SOLUTIONS)

2/  $2x - y - x \frac{dy}{dx} + 2y \frac{dy}{dx} = 0$


$\frac{dy}{dx} (2y - x) = y - 2x$

$\frac{dy}{dx} = \frac{y - 2x}{2y - x}$


- 3/ 2R 2A 3 ways  
1R 2A 3 "  
2R 1A 3 "  
1R 1A 1 "  
0R 2A 1 "  
2R 0A 1 "  
12 ways



- 1/ B  
2/ C  
3/ C  
4/ B  
5/ D  
6/ C  
7/ B  
8/ D  
9/ D  
10/ C



$\pi \cdot 2^2 \cdot 8 - \int_0^8 \pi x^2 dy$   
 $= \pi \left( 32 - \int_0^8 y^{2/3} dy \right)$   
 $= \pi \left( 32 - \left[ \frac{3y^{5/3}}{5} \right]_0^8 \right)$   
 $= \pi \left( 32 - 32 \times \frac{3}{5} \right)$   
 $= \frac{64\pi}{5}$



8/  $0.8 \omega = 4\pi$  (m/sec)  
 $\omega = 4\pi / 0.8 = 5\pi$  (rad/sec)  
 $= 300\pi$  rad/min  
 $= 150$  rev/min

10/  $F(x) = m \ddot{x}$   
 $= m \frac{d}{dx} \left( \frac{v^2}{2} \right)$   
 $\int_{x_0}^{x_1} F(x) dx = m \left[ \frac{v^2}{2} \right]_{v_0}^{v_1}$   
 $\frac{2}{m} \int_{x_0}^{x_1} F(x) dx = v_1^2 - v_0^2$   
 $v_1^2 = \frac{2}{m} \int_{x_0}^{x_1} F(x) dx + v_0^2$   
 $|v_1| = \sqrt{\frac{2}{m} \int_{x_0}^{x_1} F(x) dx + v_0^2}$

## Question 11

$$a) \int \frac{dx}{\sqrt{9-(x-2)^2}} = \frac{\sin^{-1}\left(\frac{x-2}{3}\right) + k}{3}$$

$$b) i) a(x^2+9) + (bx+1)(x-2) \\ \equiv 3x^2 - 3x + 7$$

Set  $x=2$

$$13a = 13 \quad \underline{a=1}$$

$$\text{Coeff of } x^2: a+b=3 \therefore \underline{b=2}$$

$$(ii) \int \frac{3x^2-3x+7}{(x-2)(x^2+9)} dx = \int \frac{dx}{x-2} + \int \frac{2x+1}{x^2+9} dx$$

$$= \int \frac{1}{x-2} dx + \int \frac{2x}{x^2+9} dx + \int \frac{dx}{x^2+9}$$

$$= \ln|x-2| + \ln(x^2+9) + \frac{1}{3} \tan^{-1}\left(\frac{x}{3}\right) + k$$

$$c) \text{ Let } I = \int_0^a f(x) dx$$

$$\text{Set } u = a-x \rightarrow x = a-u \\ \text{"} du = -dx \text{"}$$

$$\text{When } x=a, u=0 \\ x=0, u=a$$

$$I = \int_a^0 f(a-u) \cdot (-du)$$

$$= - \int_a^0 f(a-u) du$$

$$= \int_0^a f(a-u) du$$

$$= \int_0^a f(a-x) dx$$

(Dummy variable)

$$ii) I = \int_0^2 x(2-x)^5 dx$$

$$= \int_0^2 (2-x)x^5 dx \quad (\text{Using (i)})$$

$$= \int_0^2 2x^5 - x^6 dx$$

$$= \left[ \frac{x^6}{3} - \frac{x^7}{7} \right]_0^2 = \frac{64}{3} - \frac{128}{7}$$

$$= \frac{7 \times 64 - 3 \times 128}{21} = \frac{64}{21}$$

d) i) By factor thm:

$$(i)^2 + a(i) + 2 + i = 0$$

$$i(a+1) = -1$$

$$a+1 = -1/i = i$$

$$\underline{a = i-1}$$

ii) Let other root be  $\beta$

$$\text{Sum of roots} = -a = \beta + i$$

$$\therefore (1-i) = \beta + i$$

$$\beta = (1-i) - i$$

$$= \underline{1-2i}$$

e) i) Rotation  $\equiv$  Multiply by  $i$   
The length is shortened by a factor of 3.

$$\therefore C \text{ represents } \underline{i\omega/3}$$

ii) Diagonals bisect each other.


$$|OD| = \frac{1}{2} |OB|$$

$$OB = OA + AB = OA + OC$$


$$OD = \frac{1}{2} (\omega + \frac{i\omega}{3}) = \omega \left( \frac{1}{2} + \frac{i}{6} \right)$$

Question 12


a) i)



ii)



iii)



## Question 12 (cont)

b) i) Let the zeros of  $P(x)$  be  $\alpha, \alpha, \beta, \beta$ .

$$\therefore 2(\alpha + \beta) = 6 \quad (\text{sum of roots})$$

$$\underline{\alpha + \beta = 3} \quad (1)$$

$$\alpha^2 + \beta^2 + 4\alpha\beta = 13 \quad (\text{sum of roots 2 at a time})$$

$$(\alpha + \beta)^2 + 2\alpha\beta = 13$$

$$\therefore 2\alpha\beta = 4 \quad (\text{Using (1) above})$$

$$\underline{\alpha\beta = 2}$$

Now  $b = -\alpha^2\beta^2$  (Product of roots)  $a = 2\alpha^2\beta + 2\beta^2\alpha$  (Sum of roots 3 at a time)


$$= -(\alpha\beta)^2 = 2 \times \beta(\alpha + \beta)$$

$$\underline{b = -4} \quad \underline{a = 12}$$

ii) Two touching points is the above scenario if we solve  $x^4 - 6x^3 + 13x^2$  against  $ax + b$

$\therefore$  line is  $y = 12x - 4$

c) i)



If  $x = h$ ,  $y = \pm 2\sqrt{h}$


- Length of major axis is  $4\sqrt{h}$ .
- Length of minor axis is  $4b\sqrt{h}/a$ .
- Semi-Major, Semi-Minor are  $2\sqrt{h}$ ,  $2b\sqrt{h}/a$ .
- Area of Full Ellipse =  $4\pi hb/a$ .
- Area of Half Ellipse shown =  $2\pi hb/a$

ii)  $\delta V \doteq \frac{2\pi x b}{a} \delta x$

$$V = \lim_{\delta x \rightarrow 0} \sum_{x=0}^4 \frac{2\pi x b}{a} \delta x$$

$$= \frac{2\pi b}{a} \int_0^4 x dx = \frac{2\pi b}{a} \left[ \frac{x^2}{2} \right]_0^4$$

$$= \underline{\underline{16\pi b/a \text{ units}^3}}$$




iii) This will generate circles rather than semi ellipses  
Hence ratio  $b/a = 1$ . But it will include two halves

$$\therefore V_{\Omega} = \underline{\underline{32\pi \text{ units}^3}}$$

## Question 13

- a) i)  $\angle BQC$  is right angle  
 (Given that diagonals perpendicular)  
 $\triangle BQC$  is an angle in a semicircle with  $BC$  as diameter and  $M$  as the circle centre.  
 $MB$  and  $QM$  are radii of this circle  
 Thus equal. i.e.  $BM = QM$



- ii)  $\therefore \angle BQM = \alpha$  (Equal angles opposite equal sides in  $\triangle BQM$ )  
 $\therefore \angle NQD = \alpha$  (Vertically opposite angles are equal)

Now let  $\angle ACB = \beta$

Then  $\angle ADQ = \beta$  (Angles standing on the same arc are equal)


In  $\triangle BQC$ ,  $\alpha + \beta + 90^\circ = 180^\circ$

In  $\triangle QND$ ,  $\alpha + \beta + \angle QND = 180^\circ$  (Angle sum of triangle is  $180^\circ$ )

$\therefore \angle QND = 90^\circ$

i.e.  $MN \perp AD$


- b) Let angle of rail inclination be  $\alpha$ . There is no lateral force at the design speed, 50 km/hr  
 $50 \text{ km/h} = \frac{50000 \text{ m/s}}{3600} = 13.89 \text{ m/s}$



Resolving vertically  $R \cos \alpha = mg$  ①

horizontally  $R \sin \alpha = \frac{mv^2}{r}$  ②

②  $\div$  ①  $\tan \alpha = \frac{v^2}{rg} = \frac{(13.89)^2}{200 \times 9.81} = 0.0983$



No movement or acceleration

Resolve vertically  $N \cos \alpha + T \sin \alpha = mg$

horizontally  $N \sin \alpha - T \cos \alpha = 0$

$\therefore N = T \cot \alpha$

$\therefore \frac{T \cos^2 \alpha}{\sin \alpha} + T \sin \alpha = mg$

$T (\cos^2 \alpha + \sin^2 \alpha) = mg \sin \alpha$

$\therefore$  With  $m = 40000$ ,  $\sin \alpha = 0.09785$

$T = 38396 \text{ N} = 38.4 \text{ kN}$  (to 3sf)



### Question 13 (cont)

$$\begin{aligned} \text{c) i)} \quad V &= Ay \\ \therefore \frac{dV}{dt} &= A \frac{dy}{dt} \\ \therefore \frac{dy}{dt} &= \frac{1}{A} \frac{dV}{dt} = \frac{1}{A} (-k\sqrt{y}) = \underline{\underline{-\frac{k\sqrt{y}}{A}}} \end{aligned}$$

$$\begin{aligned} \text{ii)} \quad \frac{dt}{dy} &= \frac{-A}{k} y^{-1/2} \\ t &= -\frac{2A}{k} y^{1/2} + c \end{aligned}$$

$$\text{When } t=T, y=0 \quad \therefore c=T$$

$$\therefore t = -\frac{2A\sqrt{y}}{k} + T \quad *$$

$$\text{When } t=0, y=y_0$$

$$\therefore T = \frac{2A\sqrt{y_0}}{k} \Rightarrow \frac{A}{k} = \frac{T}{2\sqrt{y_0}}$$

Sub this into \*

$$t = -\frac{2T}{2\sqrt{y_0}} \sqrt{y} + T$$

$$\frac{t}{T} = 1 - \frac{\sqrt{y}}{\sqrt{y_0}}$$

$$\sqrt{\frac{y}{y_0}} = 1 - \frac{t}{T}$$

$$\frac{y}{y_0} = \left(1 - \frac{t}{T}\right)^2$$

$$\underline{\underline{y = y_0 \left(1 - \frac{t}{T}\right)^2}}$$

$$\text{iii)} \quad t=10 \text{ when } y=y_0/2, \quad \therefore \frac{1}{2} = \left(1 - \frac{10}{T}\right)^2 \Rightarrow 1 - \frac{10}{T} = \frac{1}{\sqrt{2}}$$

$$\begin{aligned} \Rightarrow \frac{10}{T} &= 1 - \frac{1}{\sqrt{2}} = \frac{\sqrt{2}-1}{\sqrt{2}} \Rightarrow T = \frac{10\sqrt{2}}{\sqrt{2}-1} \text{ sec} = 10\sqrt{2}(\sqrt{2}+1) \text{ sec} \\ &= \underline{\underline{20 + 10\sqrt{2} \text{ sec}}} \end{aligned}$$

### Question 14 (cont)

iii)  $(x+i)^5 + (x-i)^5 = 0$

$$(x^5 + {}^5C_1 x^4(i) + {}^5C_2 x^3(i)^2 + {}^5C_3 x^2(i)^3 + {}^5C_4 x(i)^4 + i^5)$$

$$+ (x^5 + {}^5C_1 x^4(-i) + {}^5C_2 x^3(-i)^2 + {}^5C_3 x^2(-i)^3 + {}^5C_4 x(-i)^4 + (-i)^5) = 0$$

The imaginary terms cancel

$$2x^5 + 2{}^5C_2 x^3 i^2 + 2{}^5C_4 x i^4 = 0$$

But  ${}^5C_2 = 10$ ,  ${}^5C_4 = 5$ ,  $i^2 = -1$ .  $\therefore x^5 - 10x^3 + 5x = 0$  is the equivalent equation

$$x(x^4 - 10x^2 + 5) = 0$$

But the 0 root is given by  $x$  factor.


$\therefore$  roots of  $x^4 - 10x^2 + 5 = 0$  are  $\pm \cot \frac{\pi}{10}$ ,  $\pm \cot \frac{3\pi}{10}$

iv) solve:  $x^2 = \frac{10 \pm \sqrt{100 - 20}}{2} = 5 \pm 2\sqrt{5}$

$\therefore$  Four roots are  $\pm \sqrt{5 \pm 2\sqrt{5}}$ .

But the positive roots are  $\cot \frac{\pi}{10}$ ,  $\cot \frac{3\pi}{10}$   
The largest positive root is  $\cot \frac{\pi}{10}$ .

$$\therefore \underline{\underline{\cot \frac{\pi}{10} = \sqrt{5 + 2\sqrt{5}}}}$$



The points  $(a,b)$  and  $(b,a)$  lie on the path

$$\left. \begin{aligned} \therefore b &= a \tan \alpha - \frac{g a^2 \sec^2 \alpha}{2v^2} & (1) \\ a &= b \tan \alpha - \frac{g b^2 \sec^2 \alpha}{2v^2} & (2) \end{aligned} \right\} \text{ to be solved for } v \text{ and } \alpha.$$

## Question 14

a)  $\ddot{x} = -n^2 x$  ( $2 = \frac{2\pi}{n} \rightarrow n = \pi$ )

$$\therefore \frac{d}{dx} \left( \frac{v^2}{2} \right) = -\pi^2 x$$

$$\frac{v^2}{2} = -\frac{\pi^2 x^2}{2} + k$$

When  $x = 3, v = 0 \therefore k = 9\pi^2/2$

$$\therefore v^2 = (9 - x^2)\pi^2$$

Max speed when  $x = 0$   $|v|_{\max} = \underline{\underline{3\pi \text{ m/sec}}}$

Max acc. at  $x = -3$   $\ddot{x}_{\max} = \underline{\underline{3\pi^2 \text{ m/sec}^2}}$

b) i)  $(\cot \theta + i)^n + (\cot \theta - i)^n = \left( \frac{\cos \theta + i \sin \theta}{\sin \theta} \right)^n + \left( \frac{\cos \theta - i \sin \theta}{\sin \theta} \right)^n$   
 $= \frac{1}{\sin^n \theta} \left\{ (\cos \theta + i \sin \theta)^n + (\cos(-\theta) + i \sin(-\theta))^n \right\}$   
(de Moivre)  $= \frac{1}{\sin^n \theta} (\cos n\theta + i \sin n\theta + \cos(-n\theta) + i \sin(-n\theta))$   
 $= \frac{1}{\sin^n \theta} (\cos n\theta + i \sin n\theta + \cos n\theta - i \sin n\theta)$   
 $= \underline{\underline{\frac{2 \cos n\theta}{\sin^n \theta}}}$

ii) Put  $n=5, x = \cot \theta$  where  $\theta$  are roots of  $\frac{2 \cos 5\theta}{\sin^5 \theta} = 0$

i.e.  $\cos 5\theta = 0$

$$5\theta = 2k\pi \pm \frac{\pi}{2} \quad (\text{general solution for } \cos)$$
$$= \pm \frac{\pi}{2} \quad (k=0),$$

$$2\pi \pm \frac{\pi}{2} \quad (k=1) \quad \text{or} \quad -2\pi \pm \frac{\pi}{2} \quad (k=-1)$$

$$\therefore \theta = \pm \frac{\pi}{10}, \frac{5\pi}{10}, \frac{3\pi}{10}, -\frac{5\pi}{10}, -\frac{3\pi}{10}$$

The roots of equation are  $\cot(\frac{\pi}{2}), \cot(\pm \frac{\pi}{10}), \cot(\pm \frac{3\pi}{10})$   
( $\cot(\pm \frac{5\pi}{10})$  is same as  $\cot(\frac{\pi}{2})$ )

i.e. since  $\cot(-\theta) = -\cot \theta$ , roots are  $\underline{\underline{0, \pm \cot \frac{\pi}{10}, \pm \cot \frac{3\pi}{10}}}$   
 $\uparrow$   
 $\cot(\frac{\pi}{2})$

### Question 14 (cont)

$$\textcircled{1} \times b^2 \quad b^3 = ab^2 \tan \alpha - \frac{ga^2 b^2 \sec^2 \alpha}{2V^2} \quad \textcircled{3}$$

$$\textcircled{2} \times a^2 \quad a^3 = a^2 b \tan \alpha - \frac{gab^2 \sec^2 \alpha}{2V^2} \quad \textcircled{4}$$

$$\textcircled{3} - \textcircled{4} \quad b^3 - a^3 = \tan \alpha \cdot ab(b-a)$$

$$\therefore \tan \alpha = \frac{b^3 - a^3}{ab(b-a)} = \frac{b^2 + ab + a^2}{ab} \quad *$$

$$\textcircled{1} \times b \quad b^2 = ab \tan \alpha - \frac{ga^2 b \sec^2 \alpha}{2V^2} \quad \textcircled{5}$$

$$\textcircled{2} \times a \quad a^2 = ab \tan \alpha - \frac{gab^2 \sec^2 \alpha}{2V^2} \quad \textcircled{6}$$

$$\textcircled{5} - \textcircled{6} \quad b^2 - a^2 = \frac{g \sec^2 \alpha}{2V^2} \cdot ab(b-a)$$

$$\frac{2V^2}{g \sec^2 \alpha} = \frac{ab}{a+b}$$

If the range is  $r$ , then  $(r, 0)$  lies on the curve

$$\therefore 0 = r \tan \alpha - \frac{gr^2 \sec^2 \alpha}{2V^2}$$

Rejecting the  $r=0$ , solution

$$\frac{gr \sec^2 \alpha}{2V^2} = \tan \alpha$$

$$r = \tan \alpha \frac{2V^2}{g \sec^2 \alpha} = \frac{b^2 + ab + a^2}{ab} \frac{ab}{a+b} = \frac{b^2 + ab + a^2}{a+b}$$

$$\text{ii) From } * \quad \tan \alpha = \frac{b^2 + ab + a^2}{ab}$$

But since  $(a-b)^2 > 0$  (since  $b \neq a$ , given)  
 $a^2 + b^2 - 2ab > 0$   
 $a^2 + b^2 > 2ab$



$$\therefore \tan \alpha > \frac{2ab + ab}{ab}$$

$$\tan \alpha > 3$$

$$\alpha > \tan^{-1}(3)$$

Since  $\alpha$  is acute and  $\tan^{-1}$  is an increasing function.

# Question 15



$$\delta V \doteq 4\pi xy \delta x$$

$$\therefore V = \lim_{\delta x \rightarrow 0} \sum_{x \in d} 4\pi xy \delta x$$

where  $d$  as in diagram

$$= 4\pi \int_d^r xy \, dx$$

But, by Pythagoras,  $d = \sqrt{r^2 - h^2}$ ,  $y = \sqrt{r^2 - x^2}$


$$\therefore V = 4\pi \int_{\sqrt{r^2 - h^2}}^r x \sqrt{r^2 - x^2} \, dx$$

$$= \left[ \frac{4\pi}{3} (-1) (r^2 - x^2)^{3/2} \right]_{\sqrt{r^2 - h^2}}^r$$


$$= \frac{4\pi}{3} \left\{ -0 + (r^2 - (r^2 - h^2))^{3/2} \right\} = \frac{4\pi h^3}{3} \quad (\text{independent of } r)$$

b) i)  $z^5 = 2^5(-1)$  Roots are

- $2(\cos \pi/5 + i \sin \pi/5)$  C
- $2(\cos 3\pi/5 + i \sin 3\pi/5)$  B
- $2(\cos \pi + i \sin \pi) = -2$  A
- $2(\cos(-3\pi/5) + i \sin(-3\pi/5))$  E
- $2(\cos(-\pi/5) + i \sin(-\pi/5))$  D



Question 15 (cont)



$$\begin{aligned} \text{Total area} &= \Delta COB + \Delta AOC + \Delta AOD \\ &= \frac{1}{2} \times 2 \times 2 \sin \frac{2\pi}{5} + 2 \left( \frac{1}{2} \times 2 \times 2 \sin \frac{4\pi}{5} \right) \\ &= 2 \sin \frac{2\pi}{5} + 4 \sin \frac{4\pi}{5} \\ &= 1.902113 + 2.35114 \\ &= \underline{\underline{4.25 \text{ u}^2 \text{ (to 2DP)}}} \end{aligned}$$

c) Let M be the point (X, Y)

The chord of contact from M to the ellipse is given by


$$xX + 4yY = 4 \quad (1)$$

The tangent from T has equation

$$x \left( \frac{2}{t} \right) + 2ty = 8$$

$$\text{or } \frac{x}{t} + ty = 4 \quad (2)$$

(1) and (2) are the same line



$$\left. \begin{aligned} X &= 1/t \\ Y &= t/4 \end{aligned} \right\} \text{Eliminate } t \Rightarrow XY = \frac{1}{4} \quad \text{Locus of M is } \underline{\underline{xy = \frac{1}{4}}}$$

ii) This is a rectangular hyperbola, centred on origin. It will be "tighter" to the axes.

Chords of contact cannot be drawn from inside ellipse. The curve crosses ellipse when

$$x^2 + \frac{4}{16x^2} = 4$$

$$\left. \begin{aligned} \text{Restrictions on } x: & \quad x > \sqrt{\frac{2 + \sqrt{15}}{2}} \\ & \quad 0 < x < \sqrt{\frac{2 - \sqrt{15}}{2}} \end{aligned} \right\}$$

$$\begin{aligned} 4x^4 - 16x^2 + 1 &= 0 \\ x^2 &= \frac{16 \pm \sqrt{240}}{8} \quad / 8 \\ &= 2 \pm \frac{\sqrt{15}}{2} \end{aligned}$$

And similar on negative side.

## Question 16

a) Alan picks first:

$$\text{Prob. of success} = \frac{1}{n} + \frac{(n-1)(n-2)}{n(n-1)(n-2)} \frac{1}{n} + \frac{(n-1)(n-2)(n-3)(n-4)}{n(n-1)(n-2)(n-3)(n-4)} \frac{1}{n} + \dots$$

↑            ↑    ↑    ↑            ... +  $\frac{(n-1)(n-2)}{n(n-1)} \frac{1}{n} \frac{1}{5} \frac{1}{4} \frac{1}{3} \frac{1}{2} \frac{1}{1}$

1st pick    A fails   B fails   A picks

Each term is  $\frac{1}{n}$ , and there are  $(n+1)/2$  terms

$$\therefore \text{Prob} = \frac{n+1}{2n}$$

ii) Prob =  $\frac{2}{n} + \frac{n-2}{n} \frac{n-3}{n-1} \frac{2}{n} + \dots + \frac{(n-2)(n-3)}{n(n-1)} \frac{4}{6} \frac{3}{5} \frac{2}{4} \frac{1}{3} \frac{2}{2}$

$$= \frac{2}{n} \left( \frac{n-1}{n-1} + \frac{n-3}{n-1} + \frac{n-5}{n-1} + \dots + \frac{1}{n-1} \right)$$

$$= \frac{2}{n(n-1)} (1+3+5+\dots+(n-1))$$

↑ A.P. with  $a=1, d=2, "n" = \frac{n}{2}$

$$= \frac{2}{n(n-1)} \frac{n}{4} \left( 2 + \left( \frac{n}{2} - 1 \right) 2 \right) = \frac{2}{n(n-1)} \frac{n^2}{4}$$

$$= \frac{n}{2(n-1)}$$

iii) Prob =  $\frac{2}{n} + \frac{(n-2)(n-3)}{n(n-1)(n-2)} \frac{2}{n} + \dots + \frac{(n-2)(n-3)}{n(n-1)} \frac{5}{7} \frac{4}{6} \frac{3}{5} \frac{2}{4} \frac{2}{3}$

$$= \frac{2}{n} \left( \frac{n-1}{n-1} + \frac{n-3}{n-1} + \dots + \frac{2}{n-1} \right)$$

$$= \frac{2}{n(n-1)} (2+4+6+\dots+(n-1))$$

↑ AP  $a=2, d=2, "n" = \frac{n-1}{2}$

$$= \frac{2}{n(n-1)} \frac{(n-1)}{4} \left( 4 + \left( \frac{(n-1)}{2} - 1 \right) 2 \right)$$

$$= \frac{2}{n(n-1)} \frac{(n-1)(n+1)}{4}$$

$$= \frac{n+1}{2n}$$

## Question 16 (cont)

b) i) Solve  $y = mx \pm \sqrt{a^2 m^2 + b^2}$  against  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$

Sub for  $y$  to get quadratic in  $x$ .

$$\frac{x^2}{a^2} + \frac{(mx \pm \sqrt{a^2 m^2 + b^2})^2}{b^2} = 1$$

$$\frac{x^2}{a^2} + \frac{m^2 x^2 + a^2 m^2 + b^2 \pm 2mx\sqrt{a^2 m^2 + b^2}}{b^2} - 1 = 0$$

$$b^2 x^2 + a^2 m^2 x^2 + a^4 m^2 + a^2 b^2 \pm 2a^2 m \sqrt{a^2 m^2 + b^2} x - a^2 b^2 = 0$$

$$x^2(a^2 m^2 + b^2) \pm 2a^2 m \sqrt{a^2 m^2 + b^2} x + a^4 m^2 = 0$$

$$(x\sqrt{a^2 m^2 + b^2} \pm a^2 m)^2 = 0$$

As this always gives a double root, the lines  $y = mx \pm \sqrt{a^2 m^2 + b^2}$  are always tangents to the ellipse.

ii) Let  $P(x, y)$  be external to the ellipse. Then the gradients of the 2 tangents from  $P$  are given by

$$Y = mX \pm \sqrt{a^2 m^2 + b^2}$$

$$Y - mX = \sqrt{a^2 m^2 + b^2}$$

$$Y^2 + m^2 X^2 - 2mXY = a^2 m^2 + b^2$$

$$m^2(X^2 - a^2) - 2mXY + (Y^2 - b^2) = 0$$

If these lines are perpendicular, the two roots  $m_1, m_2$  must give  $m_1 m_2 = -1$ . This is the product of the roots.

$$\therefore \frac{Y^2 - b^2}{X^2 - a^2} = -1$$

$$Y^2 - b^2 = -X^2 + a^2$$

$$\underline{X^2 + Y^2 = a^2 + b^2}$$

Thus the locus of points where the two tangents are perpendicular is  $x^2 + y^2 = a^2 + b^2$ , a circle.





$$\begin{aligned} \therefore \text{Area is } CA \times CH &= \sqrt{\frac{4(a^2+m^2b^2)}{1+m^2}} \sqrt{\frac{4(a^2m^2+b^2)}{1+m^2}} \\ &= \frac{4\sqrt{(a^2+m^2b^2)(a^2m^2+b^2)}}{1+m^2} \end{aligned}$$

$$\begin{aligned} \text{iv)} \quad (a-b)^2 &\geq 0 \\ a^2+b^2-2ab &\geq 0 \\ a^2+b^2 &\geq 2ab \end{aligned}$$

Substitute  $a = \sqrt{m}$ ,  $b = \frac{1}{\sqrt{m}}$  (Requires  $m > 0$ )

$$\therefore m + \frac{1}{m} \geq \frac{2\sqrt{m}}{\sqrt{m}}, \text{ i.e. } \underline{\underline{m + \frac{1}{m} \geq 2}} \text{ for } m > 0.$$

$$\text{v)} \quad A^2 = 16a^2b^2 + \frac{16(a^2-b^2)^2}{(m + \frac{1}{m})^2}$$

Smallest  $A$  will be as  $m + \frac{1}{m} \rightarrow \infty$ , no contribution from second term

$$A_{\min}^2 = 16a^2b^2 \quad \underline{\underline{A_{\min} = 4ab}}$$

Largest  $A$  will be when  $m + \frac{1}{m} = 2$ .

$$\begin{aligned} A_{\max}^2 &= 16a^2b^2 + \frac{16(a^2-b^2)^2}{4} = 16a^2b^2 + 4(a^4+b^4-2a^2b^2) \\ &= 8a^2b^2 + 4a^4 + 4b^4 \\ &= 4(a^4+b^4+2a^2b^2) \\ &= 4(a^2+b^2)^2 \end{aligned}$$

$$\therefore A_{\max} = \underline{\underline{2(a^2+b^2)}}$$