


The King's School

4 unit mathematics

Trial HSC Examination 1994

1. (a) Evaluate $\int_0^1 \frac{x^3}{(x^2+1)^2} dx$ by using the substitution $u = x^2 + 1$.
- (b) Evaluate $\int_0^1 2x \tan^{-1} x dx$
- (c) (i) Show that $\int_0^a f(x) dx = \int_0^a f(a-x) dx$
- (ii) Prove that $I = \int_0^{\frac{\pi}{2}} \frac{A \sin x + B \cos x}{\sin x + \cos x} dx = \frac{\pi}{4}(A+B)$, A, B constants.
- (d) Find $\int \frac{2 \tan x}{\tan 2x + \sin 2x} dx$ by using the substitution $t = \tan x$.

2. (a) (i) Find the square roots of $-35 + 12i$
- (ii) Solve $z^2 - (5 + 4i)z + 11 + 7i = 0$
- (b) Describe in the Argand diagram the locus of the complex numbers z if
- (i) $|z - 1| = 4$
- (ii) $|z - 1| = |z + 1|$
- (iii) $|z - 1| + |z + 1| = 4$
- (iv) $\Re\left(\frac{i}{z}\right) = \frac{1}{2}, z \neq 0$.
- (c)



(Figure not to scale)

$\triangle ABC$ is drawn in the Argand diagram. $\hat{BAC} = 45^\circ$, $A = (10, 2)$, $B = (6, 8)$. The length of side AC is twice the length of side AB .

- Find (i) the complex number that the vector AB represents.
- (ii) the complex number that point C represents.

3. (a) Sketch carefully the hyperbola $3x^2 - y^2 = 12$, showing on your diagram the foci, the directrices and the asymptotes in their correct positions.

(b) A tangent to the parabola $y^2 = 2ax$ meets the hyperbola $xy = c^2$ in the points P, Q .

(i) Show that the equation of the tangent at $R(x_1, y_1)$ on the parabola is $y_1y = a(x + x_1)$

(ii) Show that the x coordinates of P and Q are given by the equation $ax^2 + ax_1x - c^2y_1 = 0$

(iii) Deduce the cartesian equation of the locus of the midpoint M of the interval PQ .

4. (a) (i) Sketch the line $y = x - 1$ and the rectangular hyperbola $y = \frac{1}{x-1}$ on the same axes, showing their points of intersection.

(ii) On separate diagrams and using (i), sketch the graphs of the following functions and relations. For each graph label any asymptote.

(α) $y = x - 1 + \frac{1}{x-1}$

(β) $y = |x - 1 + \frac{1}{x-1}|$

(γ) $y^2 = x - 1 + \frac{1}{x-1}$

(δ) $y = x - 1 - \frac{1}{x-1}$

(b) Consider two functions f and g for which we know the following facts:

$f(c) = g(c) = 0$, $f'(c)$ and $g'(c)$ exist, $g'(c) \neq 0$.

By considering $f'(c) = \lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c}$ and a similar result for $g'(c)$, show that

$\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \frac{f'(c)}{g'(c)}$. Hence, or otherwise, show that

(i) $\lim_{x \rightarrow 1} \frac{\ln x}{x^2 - 1} = \frac{1}{2}$

(ii) $\lim_{x \rightarrow 0} \frac{1 - \cos^7 x}{x^2} = \frac{7}{2}$

5. (a) A particle of mass m moves in the x axis under the influence of a force $\frac{mn^2}{x^3}$, n a positive constant, directed **away** from the origin, O . Initially, the particle is at rest at $x = a > 0$.

(i) Prove that the velocity v is given by $v^2 = n^2(\frac{1}{a^2} - \frac{1}{x^2})$

(ii) Deduce that $ax = \sqrt{n^2t^2 + a^4}$

(b) A particle of mass m is projected from a point O on horizontal ground with speed u at an angle of elevation α . It hits the ground again at a distance $2a$ from O and in its flight reaches a maximum height of b . The acceleration due to gravity is g and no forces other than the gravitational force act on the particle. At time t , the horizontal and vertical displacements from O are x and y , respectively

(i) Prove that $y^2 = (u \sin \alpha)^2 - 2gy$

(ii) Deduce that $\tan \alpha = \frac{2b}{a}$ and $u^2 = \frac{g}{2b}(4b^2 + a^2)$

6. (a) u, v, w are the roots of the equation $P(x) = 8x^3 + 28x^2 + 14x - 15 = 0$

(i) Form the equation with roots $2u + 3, 2v + 3$ and $2w + 3$

(ii) Hence, or otherwise, solve $P(x) = 0$.

(b) Consider the polynomial equation $f(x) = x^n + nkx + (n - 1) = 0$, $n > 1$. For what values of k will $f(x) = 0$ have a double root if

(i) n is odd

- (ii) n is even?
 (c) If $a > b > 0$, show that $P(x) = x^3 + x^2 - ax - b$ always has
 (i) two distinct stationary points, and
 (ii) 3 distinct real zeros.

7. (a) Consider the region between the line $y = x$ and the curve $y = x^3$ in the first quadrant. Take $P(x, x^3)$ as any point on the curve $y = x^3$.


(i) The region is rotated about the line $x = 2$. Use the method of cylindrical shells to find the volume of the solid of revolution.

(ii) The region is rotated about the line $y = x$. By taking a slice in the region perpendicular to $y = x$, find the volume of the solid of revolution.

(b) (i) Find a, b, c if $\frac{4n-2}{n(n+1)(n+2)} \equiv \frac{a}{n} + \frac{b}{n+1} + \frac{c}{n+2}$

(ii) Use (i) to deduce that the sum to infinity of the series $\frac{2}{1 \times 2 \times 3} + \frac{6}{2 \times 3 \times 4} + \frac{10}{3 \times 4 \times 5} + \dots$ is $1\frac{1}{2}$.

8. (a) (i)




(Figure not to scale)

In the diagram, $QR \parallel DS$ and $\frac{PD}{DQ} = \frac{PS}{SQ}$. Copy the diagram. Prove that

(α) $SQ = SR$

(β) $\hat{QSD} = \hat{STD}$

(ii)



In the diagram, PQ is a chord of an ellipse with eccentricity e . PQ produced meets a directrix at D and PP' , QQ' are drawn perpendicular to this directrix. S is the corresponding focus of the ellipse. Copy the diagram.

(α) Prove that $\frac{PP'}{QQ'} = \frac{PD}{QD}$

(β) Deduce that DS bisects $Q\hat{S}T$.

(b) Consider the series of n terms

$$S_n = 1 + \frac{2n-2}{2n-3} + \frac{(2n-2)(2n-4)}{(2n-3)(2n-5)} + \dots + \frac{(2n-2)(2n-4)\dots \times 4 \times 2}{(2n-3)(2n-5)\dots \times 3 \times 1}$$

(i) Show that $S_3 = 5$

(ii) Prove by induction, for $n \geq 1$, that $S_n = 2n - 1$.