

Section I



10 Marks

Attempt Question 1 – 10.

Allow approximately 15 minutes for this section.

Use the multiple-choice answer sheet for Question 1 – 10

Question 1

The diagram illustrates the graph of $y = f(x)$.The graph of $y = f(|x|)$ is most likely to like:

Question 2

The curve represented by the equation $\frac{x^2}{25} + \frac{y^2}{16} = 1$ is an ellipse with:

- (A) eccentricity given by $e = \frac{3}{5}$ and foci at $(\pm 3, 0)$.
- (B) eccentricity given by $e = \frac{5}{3}$ and foci at $(\pm 5, 0)$.
- (C) eccentricity given by $e = \frac{3}{5}$ and foci at $(\pm 5, 0)$.
- (D) eccentricity given by $e = \frac{4}{5}$ and foci at $(\pm 3, 0)$.

Question 3

Given that $w^3 = -1$ and that w is complex, the value of $(1 + w - w^2)^3$ is:

- (A) -8
- (B) 8
- (C) 1
- (D) -1

Question 4

The value of $\left[\frac{-1 + \sqrt{-3}}{2}\right]^{29} + \left[\frac{-1 - \sqrt{-3}}{2}\right]^{29}$ is:

- (A) -1
- (B) 1
- (C) $-\sqrt{3}$
- (D) -2

Question 5

What is the solution to the inequation: $\frac{x(5-x)}{x-4} \geq -3$?

- (A) $2 \leq x < 4$ or $x \geq 6$.
- (B) $1 \leq x < 4$ or $x \geq 5$.
- (C) $4 < x \leq 6$ or $x \leq 2$.
- (D) $4 > x \leq 5$ or $x \leq 1$.


Question 6

Consider the points $P\left(ct, \frac{c}{t}\right)$ and $Q\left(\frac{ct}{2}, \frac{2c}{t}\right)$ on the rectangular hyperbola $xy = c^2$. The locus of the midpoint, M , of PQ is:

- (A) $xy = \frac{9}{8}c^2$
- (B) $t = \frac{4x}{3c}$
- (C) $xy = 8c^2$
- (D) $xy = \frac{c^2}{2}$

Question 7

Consider the Argand diagram below.



Which inequality could define the shaded area?

- (A) $|z-1| \leq \sqrt{2}$ and $0 \leq \text{Arg}(z-i) \leq \frac{\pi}{4}$.
- (B) $|z-1| \leq \sqrt{2}$ and $0 \leq \text{Arg}(z+i) \leq \frac{\pi}{4}$.
- (C) $|z-1| \leq 1$ and $0 \leq \text{Arg}(z-i) \leq \frac{\pi}{4}$.
- (D) $|z-1| \leq 1$ and $0 \leq \text{Arg}(z+i) \leq \frac{\pi}{4}$.

Question 8

The gradient of the tangent to the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ at the point $P(a \sec \theta, b \tan \theta)$ is given by:

- (A) $\frac{a \tan \theta}{b \sec \theta}$
- (B) $-\frac{b \sec \theta}{a \tan \theta}$
- (C) $-\frac{a \tan \theta}{b \sec \theta}$
- (D) $\frac{b \sec \theta}{a \tan \theta}$

Question 9


Which of the following is an expression for $\int \frac{1}{\sqrt{x^2 - 6x + 10}} dx$?

- (A) $\ln(x - 3 - \sqrt{x^2 - 6x + 10}) + c$
- (B) $\ln(x + 3 - \sqrt{x^2 - 6x + 10}) + c$
- (C) $\ln(x - 3 + \sqrt{x^2 - 6x + 10}) + c$
- (D) $\ln(x + 3 + \sqrt{x^2 - 6x + 10}) + c$

Question 10

How many rectangles are in the chess board illustrated?

- (A) 1024
- (B) 2304
- (C) 2025
- (D) 1296



END OF SECTION I

Section II**Total Marks 90****Attempt Question 11 – 16.****Allow approximately 2 hours & 45 minutes for this section.**

Answer all questions, starting each new question on a new booklet with your **student number** and the question number on the front cover.

All necessary working must be shown in each and every question.

	Marks
Question 11. (15 marks) Use a SEPARATE writing booklet.	
(a) Given $z = 2 + i$ and $w = -1 + 2i$, find:	
(i) $ w $	1
(ii) \bar{z}	1
 (b) Illustrate the loci in the complex plane given by these equations:	
(i) $ z = z - 6 - 3i $	2
(ii) $\arg(z + 2) = \arg(z - 2 - 5i)$	3
(iii) $\operatorname{Re}(z^2) = 4$	2
 (c) Suppose z_1, z_2 and z_3 are three complex numbers, each of modulus 1, such that $z_1 + z_2 + z_3 = 0$. Suppose also that z is a complex number of modulus 3. Show that:	
(i) $ z - z_1 ^2 = 10 - (z\bar{z}_1 + \bar{z}z_1)$	3
(ii) $ z - z_1 ^2 + z - z_2 ^2 + z - z_3 ^2 = 30$	3

Question 12. (15 marks) Use a SEPARATE writing booklet.

Marks

(a) Solve the equation $x^4 + 2x^3 + x^2 - 1 = 0$, given that one root is

3

$$-\frac{1}{2} + i\frac{\sqrt{3}}{2}.$$

(b) Show that if $f(x) = x^n - 1$ (for $n > 1$) then $f(x)$ has no multiple roots.

3

(c) Given that $f(x) = x^2(x-1)$, draw a neat sketch showing any intercepts and asymptotes.

(i) $y = f(x)$

1

(ii) $y^2 = f(x)$

2

(iii) $y = [f(x)]^2$

3


(iv) $y = \frac{1}{f(x)}$

3

Question 13. (15 marks) Use a SEPARATE writing booklet.

Marks

(a)



$\triangle ABC$ is inscribed a circle. The bisector of $\angle BAC$ meets BC at X and the circle at D .

Prove (i) $\triangle ABX$ is similar to $\triangle ADC$

1

(ii) $AB.AC = AD.AX$

2

(iii) $AB.AC = [AX]^2 + BX.XC$

2

(b)

(i) Evaluate $\int_0^{\frac{\pi}{2}} \cos^3 x \, dx$

3

(ii) Use the method of partial fractions to find $\int \frac{9x+1}{x^2-2x-3} \, dx$.

3

(iii) If $I_n = \int x^n e^{ax} \, dx$,

(α) show that $I_n = \frac{1}{a} x^n e^{ax} - \frac{n}{a} I_{n-1}$

2

(β) Hence, or otherwise, evaluate $\int_1^3 x^2 e^{3x} \, dx$, giving your answer

2

correct to 2 decimal places.


- Question 14.** (15 marks) Use a SEPARATE writing booklet. **Marks**
- (a) The region bounded by the graphs of $y = x^2$ and $y = x + 2$ is revolved about the line $x = 3$. Using the method of cylindrical shells, express the volume of the resulting solid as a definite integral. **(Note: Do not evaluate this integral).** **3**
- (b) (i) Show that the cubic $f(x) = x^3 - 3x + 1$ has **exactly** one root, $x = \alpha$, between 0 and 1. **4**
- (ii) Beginning with the approximation $x = 0$, use two applications of Newton's method to gain a better approximation to $x = \alpha$. Give your answer correct to three decimal places.
- (c) Find the equation with rational coefficients whose roots are the reciprocals of the squares of the roots of $x^3 - x^2 - 14x + 24 = 0$ **3**
- (d) Let $f(x) = \frac{1}{2}(1 - \sqrt{1-x})$ for $0 \leq x \leq 1$ and define: **5**
- $f_1(x) = f(x)$ and
- $f_{n+1}(x) = f(f_n(x))$, for $n = 1, 2, 3, \dots$
- Show by mathematical induction that, for any θ in the interval $0 \leq \theta \leq \frac{\pi}{2}$,
- $f_n(\sin^2 \theta) = \sin^2\left(\frac{\theta}{2^n}\right)$, for all positive integers n .

- Question 15.** (15 marks) Use a SEPARATE writing booklet. **Marks**
- (a) (i) Prove that the equation of the tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ at the point $P(x_1, y_1)$, is given by $\frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$. **2**
- (ii) Hence, or otherwise, find the point of contact of the tangent $y = mx + \sqrt{a^2m^2 + b^2}$ and the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. **3**
- (b) The foci of a hyperbola coincide with the foci of the ellipse $\frac{x^2}{25} + \frac{y^2}{9} = 1$. **4**
Find the equation of the hyperbola if its eccentricity is 2.
- (c) (i) Prove that $\int_0^a F(x) dx = \int_0^a F(a-x) dx$. **1**
- (ii) Hence, or otherwise, evaluate $\int_0^{\frac{\pi}{4}} \log_e(1 + \tan \theta) d\theta$ **5**

Question 16. (15 marks) Use a SEPARATE writing booklet.

Marks

(a)



$ABCD$ is a cyclic quadrilateral. The diagonals AC and BD intersect at right angles at X . M is the midpoint of BC . MX produced meets AD at N .


- | | | |
|-------|---|----------|
| (i) | Copy the diagram showing the above information. | 0 |
| (ii) | Show that $\angle MBX = \angle MXB$ | 3 |
| (iii) | Show that MN is perpendicular to AD . | 4 |

Question 16 continues on page 7

Question 16 (continued)

Marks

(b)



The diagram shows a hemispherical bowl of radius r . The bowl has been tilted so that its axis is no longer vertical, but at an angle θ to the vertical. At this angle, it can hold a volume V of water.

The vertical line from the centre O meets the surface of the water at W and meets the bottom of the bowl at B . Let P be any point between W and B and let h be the distance OP .

- (i) Explain why $V = \int_{r \sin \theta}^r \pi(r^2 - h^2) dh$. 2
- (ii) Hence show $V = \frac{r^3 \pi}{3} (2 - 3 \sin \theta + \sin^3 \theta)$ 3
- (iii) The bowl has been tilted so that it is exactly half full. Using the result from Q 14 (b), find the angle θ correct to one tenth of a degree. 3

END OF PAPER