



NORTH SYDNEY BOYS HIGH SCHOOL

2007
TRIAL HIGHER SCHOOL CERTIFICATE
EXAMINATION

Mathematics Extension 2

General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write on one side of the paper (with lines) in the booklet provided
- Write using blue or black pen
- Board approved calculators may be used
- All necessary working should be shown in every question
- Each new question is to be started on a new page.

- Attempt all questions

Class Teacher:
(Please tick or highlight)

Mr Ee
 Mr Trenwith
 Mr Weiss

Student Number: _____

(To be used by the exam markers only.)

Question No	1	2	3	4	5	6	7	8	Total	Total
Mark	$\frac{15}{15}$	$\frac{15}{15}$	$\frac{15}{15}$	$\frac{15}{15}$	$\frac{15}{15}$	$\frac{15}{15}$	$\frac{15}{15}$	$\frac{15}{15}$	$\frac{120}{120}$	$\frac{100}{100}$

Mark

QUESTION 1 (15 marks)

- (a) Find $\int \frac{x}{\sqrt{16-x^2}} dx$ 2
- (b) By completing the square, find $\int \frac{8}{x^2+4x+13} dx$ 2
- (c) Use integration by parts to evaluate $\int_1^e x^4 \log_e x dx$. 4
- (d) Use the substitution $u = \cos x$ to find $\int \cos^2 x \sin^5 x dx$ 3
- (e) Express $\frac{3x+7}{(x+1)(x+2)(x+3)}$ in partial fractions and hence prove that $\int_0^1 \frac{3x+7}{(x+1)(x+2)(x+3)} dx = \ln 2$. 4

QUESTION 2 (15 marks) Start a new page

- (a) Let $z = 2+i$ and $w = 1-i$. Find, in the form $x+iy$,
- (i) $3z+iw$ 1
- (ii) $z\bar{w}$ 1
- (iii) $\frac{5}{z}$ 1
- (b) Let $\alpha = -\sqrt{3}+i$.
- (i) Express α in modulus-argument form. 2
- (ii) Express α^4 in modulus-argument form. 2
- (iii) Hence express α^4 in the form $x+iy$. 1

QUESTION 2 (Continued)


- (c) If $z_1 = 4 + i$ and $z_2 = 1 + 2i$ show geometrically how to construct the vectors representing.
- (i) $z_1 + z_2$. 1
- (ii) $z_1 - z_2$. 1
- (d) Consider the hyperbola with the equation $\frac{x^2}{16} - \frac{y^2}{9} = 1$.
- (i) Find the coordinates of the foci and x -intercepts of the hyperbola. 2
- (ii) Find the equations of the directrices and the asymptotes of the hyperbola. 2
- (iii) What are the parametric equations of this hyperbola? 1

QUESTION 3 (15 marks) Start a new page


- (a) If α , β and γ are the roots of the cubic equation $x^3 + mx + n = 0$, find in terms of m and n , the values of
- (i) $\frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$ 3
- (ii) $\alpha^3 + \beta^3 + \gamma^3$ 2
- (iii) Determine the cubic equation whose roots are α^2 , β^2 and γ^2 . 3
- (b) Given that the equation $x^4 - 5x^3 - 9x^2 + 81x - 108 = 0$ has a triple root, find all the roots of the equation. 4
- (c) If $y = e^{-x}(A \sin 2x + B \cos 2x)$, prove that 3
- $$\frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + 5y = 0$$

QUESTION 4 (15 marks) Start a new page

- (a) Given $f(x) = x^2 - 2x$. On separate diagrams sketch the graphs of the following. Indicate clearly any asymptotes, intercepts with the axes and local maxima and minima.



- (i) $y = |f(x)|$ 1
- (ii) $y = f(|x|)$ 1
- (iii) $y = \frac{1}{f(x)}$ 2
- (iv) $y^2 = f(x)$ 2
- (v) $y = [f(x)]^2$ 1
- (vi) $y = \ln[f(x)]$ 2
- (b) The region bounded by $y = \ln x$, $x = e$ and the x -axis is rotated about the y -axis. Find the volume of rotation. Use the method of cylindrical shells. 3



- (b) First differentiating both sides of the formula


$$1 + x + x^2 + x^3 + \dots + x^n = \frac{x^{n+1} - 1}{x - 1}$$

then find an expression for

$$1 + 2 \times 2 + 3 \times 4 + 4 \times 8 + \dots + n 2^{n-1}$$

QUESTION 5 (15 marks) Start a new page

- (a) In the circle shown below, the diameter AB meets the chord CD at right angles at M. The tangents at C and D meet at T.



(i) Show that BC bisects $\angle MCT$. 2

(ii) Show that triangle BCM is similar to triangle CAM. Hence, show that $CM^2 = AM \times BM$. 2

(iii) Show that $TB \times TA = MB \times TA + TB \times TM$. 4

(iv) Hence, or otherwise, show that M divides the interval AB internally in the same ratio that T divides AB externally. 2


(c) If $I_n = \int \tan^n x \, dx$

(i) Show that $I_n = \frac{1}{n-1} \tan^{n-1} x - I_{n-2}$. 3

(ii) Find $\int \tan^6 x \, dx$. 2

QUESTION 6 (15 marks) Start a new page


- (a) The solid shown has a semicircular base of radius 2 units. Vertical cross-sections perpendicular to the diameter are right-angled triangles whose height is bounded by the parabola $z = 4 - x^2$.



By slicing at right angles to the x -axis, show that the volume of the solid is given by

$$V = \int_0^2 (4 - x^2)^{3/2} dx, \text{ and hence calculate this volume.}$$

- (b) The tangents at $P(x_1, y_1)$, $Q(x_2, y_2)$ on the hyperbola $xy = 1$ intersect at the point $T(x_0, y_0)$.



(i) Show that the tangent at $P(x_1, y_1)$ has equation $xy_1 + yx_1 = 2$. 2

(ii) Show that the chord of contact PQ has equation $xy_0 + yx_0 = 2$. 2

(iii) Show that x_1 and x_2 are the roots of the quadratic equation $y_0 x^2 - 2x + x_0 = 0$. 2

(iv) Hence, or otherwise, show that the midpoint R, of PQ has coordinates $\left(\frac{1}{y_0}, \frac{1}{x_0}\right)$. 2

(v) Hence, or otherwise, show that as T moves on the hyperbola $xy = c^2$, $0 < c < 1$, R moves on the hyperbola $xy = \frac{1}{c^2}$. 2

QUESTION 7 (15 marks) Start a new page

- (a) (i) Show that $\tan\left(A + \frac{\pi}{2}\right) = -\cot A$ 1
- (ii) Use mathematical induction to prove that $\tan\left[(2n+1)\frac{\pi}{4}\right] = (-1)^n$ for all integer $n \geq 1$. 4
- (b) Two stones are thrown simultaneously from the same point in the same direction with the same non-zero angle of projection (upward inclination to the horizontal), α , but with different velocities U, V metres per second ($U < V$).
- The slower stone hits the ground at a point P on the same level as the point of projection. At that instant the faster stone just clears a wall of height h metres above the level of projection and its (downward) path makes an angle β with the horizontal.
- (i) Show that while the stones are in flight, the line joining them has a gradient of $\tan \alpha$. 3
- (ii) Hence, express the horizontal distance from P to the foot of the wall in terms of h and α . 2
- (iii) Show that $V(\tan \alpha + \tan \beta) = 2U \tan \alpha$. 3
- (iv) Hence, deduce that, if $\beta = \frac{1}{2}\alpha$, then $U < \frac{3}{4}V$. 2

QUESTION 8 (15 marks) Start a new page

- (a) Show that the locus in the Argand plane represented by the equation $|z-1| + |z+1| = 4$ is a conic and find its cartesian equation. 3
- (b) A particle of mass m is projected against a constant gravitational force mg and resistance $\frac{mv}{k}$, where v is the velocity of the particle and k is a constant. Let x be the distance traveled in time t . Initially the particle has zero displacement and $v_0 = k(h - g)$, where h is a constant.
- (i) Show that the equation of motion of the particle is $\ddot{x} = -\left[\frac{kg+v}{k}\right]$ 1
- (ii) Show that $t = k \log\left(\frac{kh}{kg+v}\right)$ 2
- (iii) Find the time taken by the particle to reach the maximum height, H , and determine the height of that point. 3
- (c) A polynomial $P(x)$ is divided by $x^2 - a^2$ where $a \neq 0$, and the remainder is $px + q$.
- (i) Show that $p = \frac{1}{2a}[P(a) - P(-a)]$ 3
- and $q = \frac{1}{2}[P(a) + P(-a)]$
- (ii) Find the remainder when $P(x) = x^n - a^n$, for n a positive integer, is divided by $x^2 - a^2$. 3

2007 Trial HSC Ext 2 Suggested Solutions

1(a) Let $u = 16 - x^2$
 $\frac{du}{dx} = -2x$
 $dx = -\frac{du}{2x}$

$\int \frac{x}{\sqrt{16-x^2}} dx = \int \frac{x}{\sqrt{u}} \cdot \frac{du}{-2x}$
 $= -\frac{1}{2} \int u^{-\frac{1}{2}} du$
 $= -\frac{1}{2} (2\sqrt{u}) + C$
 $= -\sqrt{16-x^2} + C$

1(b) $\int \frac{8}{x^2+4x+13} dx$
 $= \int \frac{8}{(x+2)^2+9} dx$
 $= \frac{8}{3} \tan^{-1}\left(\frac{x+2}{3}\right) + C$

1(c) $\int x^4 \ln x dx$
 $= \left[\frac{x^5}{5} \ln x \right] - \int \frac{x^5}{5} \cdot \frac{1}{x} dx$
 $= \frac{e^5}{5} - \frac{1}{5} \left[\frac{x^5}{5} \right]_1^e$
 $= \frac{e^5}{5} - \frac{e^5}{25} + \frac{1}{25} = \frac{4e^5+1}{25}$

1(d) Let $u = \cos x$
 $\frac{du}{dx} = -\sin x$ $du = -\sin x dx$
 $\int \cos^2 x \sin^5 x dx$
 $= \int \cos^2 x \sin^4 x \sin x dx$
 $= \int u^2 (1-u)^2 (-du)$
 $= -\int u^2 - 2u^3 + u^4 du$

$= -\frac{1}{3} u^3 + \frac{2}{5} u^5 - \frac{1}{7} u^7 + C$
 $= -\frac{1}{3} \cos^3 x + \frac{2}{5} \cos^5 x - \frac{1}{7} \cos^7 x + C$

1(e) $\frac{3x+7}{(x+1)(x+2)(x+3)} \equiv \frac{A}{x+1} + \frac{B}{x+2} + \frac{C}{x+3}$
 $3x+7 \equiv A(x+2)(x+3) + B(x+1)(x+3) + C(x+1)(x+2)$
 $7 = 2A + 3A + 3B + 3B + 3C + 2C + 2C + 3C$
 $7 = 5A + 6B + 7C$
 $7 = 5A + 6B + 7C$
 $7 = 5A + 6B + 7C$
 $7 = 5A + 6B + 7C$

$\int_0^1 \frac{3x+7}{(x+1)(x+2)(x+3)} dx$
 $= \int_0^1 \left(\frac{2}{x+1} - \frac{1}{x+2} - \frac{1}{x+3} \right) dx$
 $= [2 \ln(x+1) - \ln(x+2) - \ln(x+3)]_0^1$
 $= (2 \ln 2 - \ln 3 - \ln 4) - (2 \ln 1 - \ln 2 - \ln 3)$
 $= \ln \frac{4}{2} + \ln 6 = \ln \frac{24}{2} = \ln 12$

Q2. (a) $z = 2+i$, $w = 1-i$

(i) $3z+iw$
 $= 3(2+i) + i(1-i)$
 $= 6+3i+i+i+1$
 $= 7+4i$

(ii) $z\bar{w}$
 $= (2+i)(1-i)$
 $= 2+i+i-i-1$
 $= 1+3i$


(iii) $\frac{5}{z}$
 $= \frac{5}{2+i} \cdot \frac{2-i}{2-i}$
 $= \frac{10-5i}{4+i} = 2-i$

(b) $\alpha = -\sqrt{5} + i$

(i) $|\alpha| = (-\sqrt{5})^2 + 1^2$
 $= \sqrt{4} = 2$
 $\arg \alpha = \tan^{-1} \frac{1}{\sqrt{5}}$
 $= \frac{5\pi}{6}$

(ii) $\alpha^4 = 2^4 \operatorname{cis}\left(4 \times \frac{5\pi}{6}\right)$
 $= 16 \operatorname{cis}\left(-\frac{2\pi}{3}\right)$

(iii) $\alpha^4 = 16\left(\cos\left(-\frac{2\pi}{3}\right) + i \sin\left(-\frac{2\pi}{3}\right)\right)$
 $= 16x - \frac{1}{2} + i(16x - \frac{\sqrt{3}}{2})$
 $= -8 - 8\sqrt{3}i$



(d) (i) $\frac{x^2}{16} - \frac{y^2}{9} = 1$
 $a=4, b=3$
 $b^2 = a^2(e^2 - 1)$
 $9 = 16(e^2 - 1)$
 $e^2 - 1 = \frac{9}{16}$
 $e^2 = \frac{9}{16} + 1 = \frac{25}{16}$
 $e = \frac{5}{4}$
 $S \equiv (ae, 0), S' \equiv (-ae, 0)$
 $\equiv (5, 0) \equiv (-5, 0)$

When $y=0, \frac{x^2}{16} = 1$
 $x = \pm 4$
 $\therefore x$ -intercepts are 4 & -4

(ii) Directrices are
 $x = \pm \frac{a}{e}$
 $= \pm \frac{16}{5}$
 Eqn of asymptotes are
 $y = \pm \frac{b}{a} x$
 $= \pm \frac{3}{4} x$

(iii) $x = 4 \sec \theta$
 $y = 3 \tan \theta$

$$3(a) \quad x^3 + mx + n = 0.$$

$$\alpha\beta\gamma = -n$$

$$\alpha + \beta + \gamma = 0$$

$$\alpha\beta + \alpha\gamma + \beta\gamma = m.$$

$$(i) \quad \frac{1}{\alpha} + \frac{1}{\beta} + \frac{1}{\gamma}$$

$$= \frac{\beta\gamma + \alpha\gamma + \alpha\beta}{\alpha\beta\gamma} = \frac{m}{-n}$$

$$(ii) \quad \alpha^3 + m\alpha + n = 0 \quad \text{--- (1)}$$

$$\beta^3 + m\beta + n = 0 \quad \text{--- (2)}$$

$$\gamma^3 + m\gamma + n = 0 \quad \text{--- (3)}$$

(1) + (2) + (3)

$$\alpha^3 + \beta^3 + \gamma^3 + m(\alpha + \beta + \gamma) + 3n = 0.$$

$$\alpha^3 + \beta^3 + \gamma^3 = -3n.$$

$$(iii) \quad (\sqrt{x})^3 + m\sqrt{x} + n = 0$$

$$\sqrt{x}(x+m) + n = 0$$

$$\sqrt{x} = \frac{-n}{x+m}$$

$$x = \frac{n^2}{(x+m)^2}$$

$$x(x+m)^2 - n^2 = 0.$$

$$x^3 + 2mx^2 + m^2x - n^2 = 0.$$

$$(b) \quad P(x) = x^4 - 5x^3 - 9x^2 + 81x - 108$$

$$P'(x) = 4x^3 - 15x^2 - 18x + 81$$

$$P''(x) = 12x - 30x - 18 \quad \text{--- (1)}$$

$$\text{When } P''(x) = 0 \quad 12x - 30x - 18 = 0.$$

$$x = -\frac{1}{2} \text{ or } x = 3 \quad \text{--- (2)}$$

$$P'(3) + P(3) = 0.$$

$\therefore x = 3$ is a triple root of

$$P(x) = 0 \quad \text{--- (3)}$$

If the other root is α

$$\text{Sum of roots} = 9 + \alpha = 5.$$

$$\alpha = -4. \quad \text{--- (4)}$$

\therefore Roots of $P(x) = 0$ are $3, 3, 3, + -4$.

$$(c) \quad y = e^{-x} (A \sin 2x + B \cos 2x)$$

$$e^x y = A \sin 2x + B \cos 2x.$$

$$e^x \frac{dy}{dx} + y e^x = 2A \cos 2x - 2B \sin 2x.$$

$$e^x \frac{dy}{dx} + \frac{dy}{dx} e^x + y e^x + e^x \frac{dy}{dx}$$


$$= -4A \sin 2x - 4B \cos 2x \quad \text{--- (1)}$$

$$= -4e^x y.$$



$$e^x \left(\frac{d^2y}{dx^2} + 2 \frac{dy}{dx} + 5y \right) = 0.$$

$$\therefore \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} + 5y = 0. \quad \text{--- (2)}$$


$$Q4. (a) (i) \quad y = |f(x)|$$




$$(ii) \quad y = f(|x|)$$




$$(iv) \quad y^2 = f(x)$$




$$(v) \quad y = [f(x)]^2$$



$$(vi) \quad y = \ln[f(x)]$$



(b)



$$\delta V = [\pi(x+\delta x)^2 - \pi x^2] \delta x$$

$$= 2\pi x y \delta x + \pi y \delta x^2$$

$$\approx 2\pi x y \delta x.$$

$$V = \lim_{\delta x \rightarrow 0} \sum_{x=1}^e 2\pi x y \delta x$$

$$= \int_1^e 2\pi x y dx \quad \text{--- (1)}$$

$$= 2\pi \int_1^e x \ln x dx$$

$$= 2\pi \left[\frac{x^2}{2} \ln x \right]_1^e - 2\pi \int_1^e \frac{x}{2} dx$$

$$= \pi \left[x^2 \ln x - \frac{x^2}{2} \right]_1^e$$

$$= \pi \left[(e^2 - \frac{e^2}{2}) - (0 - \frac{1}{2}) \right]$$

$$= \frac{\pi}{2} (e^2 + 1) \text{ unit}^3. \quad \text{--- (2)}$$

(c) $1+x+x^2+\dots+x^n = \frac{x^{n+1}-1}{x-1}$

f. both sides w.r.t. x

$$2x+3x^2+\dots+n x^n = \frac{(n+1)x^n(x-1) - (x-1)}{(x-1)^2}$$


$$= \frac{(n+1)x^{n+1} - (n+1)x^n - x + 1}{(x-1)^2}$$

$$= \frac{n x^{n+1} - (n+1)x^n + 1}{(x-1)^2}$$

Put $x=2$

$$2 \times 2 + 3 \times 4 + 4 \times 8 + \dots + n \times 2^n = \frac{n \times 2^{n+1} - (n+1)2^n + 1}{(2-1)^2}$$

$$= 2^n [2n - (n+1)] + 1 = 2^n (n+1) + 1$$



(d) Join AC + BC
 $\angle ACB = 90^\circ$ (\angle in a semi-c)
 $\angle TCB = \angle BAC$ (\angle bet. tangent + chord = \angle in alt. seg)
 $\angle ACM = 90^\circ - \alpha$ (\angle sum of Δ)
 But $\angle ACB = 90^\circ$
 $\therefore \angle BCM = 90^\circ - (90^\circ - \alpha) = \alpha$
 $\therefore \angle TCB = \angle BCM$
 i.e. BC bisects $\angle TCM$.

(ii) In $\Delta BCM + \Delta CAM$.
 $\angle BMC = \angle AMC = 90^\circ$ ($CM \perp AB$)
 $\angle CAM = \angle BCM = \alpha$ (proved above)
 $\therefore \Delta BCM \sim \Delta CAM$ (equiv. angles)
 $\frac{CM}{AM} = \frac{BM}{CM}$ (corresp. sides of similar Δ s)
 $\therefore CM^2 = AM \times BM$
 (iii) $CT^2 = TA \times TB$
 (The sq. on the tangent equal the product of the segs. of any chord through the pt.)
 In ΔTCM
 $CT^2 = CM^2 + MT^2$
 $= AM \times BM + MT^2$
 $TA \times TB = CT^2$
 $= AM \times BM + MT^2$
 $= (TA - TM) \times MB + MT^2$
 $= TA \times MB - TM \times MB + MT^2$
 $= TA \times MB - TM(TM - MB)$
 $= TA \times MB - TM \times TB$
 (iv) $TA \times TB - TM \times TB = TA \times MB$
 $TB(TA - TM) = TA \times MB$
 $TB \times AM = TA \times MB$
 $\frac{AM}{MB} = \frac{TA}{TB}$
 i.e. M divides AB internally in the same ratio as T divides BR externally.
 (e) $I_n = \int \tan^n x dx$
 $= \int \tan^{n-2} (\sec^2 x - 1) dx$
 $= \int \tan^{n-2} \sec^2 x dx - \int \tan^{n-2} dx$
 $= \frac{\tan^{n-1}}{n-1} - I_{n-2}$
 (ii) $I_6 = \frac{1}{5} \tan^5 x - I_4$
 $= \frac{1}{5} \tan^5 x - \left[\frac{1}{3} \tan^3 x - I_2 \right]$
 $= \frac{1}{5} \tan^5 x - \frac{1}{3} \tan^3 x + \tan x - I_0$
 $= \frac{1}{5} \tan^5 x - \frac{1}{3} \tan^3 x + \tan x - x + C$

Q6 (w) $\delta V = \frac{1}{2} z y \delta x$

$$= \frac{1}{2} (4-x^2) \sqrt{4-x^2} \delta x$$

$$= \frac{1}{2} (4-x^2)^{3/2} \delta x$$

$$V = \lim_{\delta x \rightarrow 0} \sum_{x=2}^0 \frac{1}{2} (4-x^2)^{3/2} \delta x$$

$$= \int_{-2}^2 \frac{1}{2} (4-x^2)^{3/2} dx$$

$$= \int_0^2 (4-x^2)^{3/2} dx \quad \left[(4-x^2)^{3/2} \text{ is an even function} \right]$$

Let $x = 2 \sin \theta$ $\frac{dx}{d\theta} = 2 \cos \theta$

$$4-x^2 = 4(1-\sin^2 \theta) = 4 \cos^2 \theta$$

$$= 4 \cos^4 \theta$$

$$x=0, 2 \sin \theta = 0 \Rightarrow \theta = 0$$

$$x=2, 2 \sin \theta = 2 \Rightarrow \theta = \frac{\pi}{2}$$

$$V = \int_0^{\pi/2} (4 \cos^2 \theta)^{3/2} \cdot 2 \cos \theta d\theta$$

$$= 16 \int_0^{\pi/2} \cos^4 \theta d\theta$$

$$= 16 \int_0^{\pi/2} \frac{\cos 2\theta + 1}{2} d\theta$$

$$= 4 \int_0^{\pi/2} (\cos 2\theta + 2 \cos 2\theta + 1) d\theta$$

$$= 4 \int_0^{\pi/2} \frac{\cos 4\theta + 1}{2} + 2 \cos 2\theta + 1 d\theta$$

$$= 2 \int_0^{\pi/2} \cos 4\theta + \cos 2\theta + 3 d\theta$$

$$= 2 \left[\frac{\sin 4\theta}{4} + \frac{\sin 2\theta}{2} + 3\theta \right]_0^{\pi/2}$$

$$= 2 \left[(0+0+3\frac{\pi}{2}) - (0+0+0) \right]$$

$$= 3\pi \text{ unit}^2$$

(b) (i) $xy=1 \Rightarrow \frac{dy}{dx} = -\frac{1}{x^2}$

At (x_1, y_1) $m_T = -\frac{1}{x_1^2}$

Eqn of tangent at (x_1, y_1) is

$$y - y_1 = -\frac{1}{x_1^2} (x - x_1)$$

But $x_1 y_1 = 1 \Rightarrow y_1 = \frac{1}{x_1}$

$$x_1 y - x_1 y_1 = -\frac{x}{x_1} + 1$$

$$\therefore x_1 y - 1 = -\frac{x}{x_1} + 1$$

$$x_1 y + x_1 y = 2$$

(ii) Tangent at P + Q are

$$x_1 y_0 + x_0 y = 2$$

$$x_2 y_0 + x_0 y = 2$$

Egn $x_1 y_0 + x_0 y = 2$ satisfied by both tangent P + Q.

$$\therefore x_1 y_0 + x_0 y = 2$$
 is eqn of PQ.

(iii) x_1, x_2 are solution to the set of simultaneous eqns

$$x_1 y_0 + x_0 y = 2 \quad \text{--- (1)}$$

$$\text{and } x_1 y = 1 \quad \text{--- (2)}$$

Sub $y = \frac{1}{x}$ into (1)

$$x_1 y_0 + \frac{x_0}{x} = 2$$

$$x^2 y_0 + x_0 = 2x$$

$$x^2 y_0 - 2x + x_0 = 0$$

(iv) $R = \left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2} \right)$

from (iii) $x_1 + x_2 = \frac{2}{y_0}$ (sum of roots)

$$\therefore \frac{x_1+x_2}{2} = \frac{1}{y_0}$$

Sub into eqn of PQ

$$\frac{1}{y_0} \cdot y_0 + x_0 y = 2$$

$$1 + x_0 y = 2$$

$$y = \frac{1}{x_0}$$

$$\therefore R = \left(\frac{1}{x_0}, \frac{1}{x_0} \right)$$


26(c)(v) If T moves on $xy = c^2$, then $x_0 y_0 = c^2$.
 For R $x = \frac{1}{y_0}$, $y = \frac{1}{x_0}$.
 $xy = \frac{1}{x_0 y_0} = \frac{1}{c^2}$ which is a hyperbola.

27 (a) $\tan(A + \frac{\pi}{2}) = -\tan(\frac{\pi}{2} - (A + \frac{\pi}{2})) = -\cot A$

(ii) $\tan\left[\frac{\pi}{2} + \frac{\pi}{4}\right] = \tan\left(\frac{3\pi}{4}\right) = -1$
 For $n=1$ LHS = $\tan\frac{3\pi}{4} = -1$
 RHS = $(-1)^1 = -1 =$ LHS.

is true for $n=1$.
 Assume true for $n=k$
 $\tan\left[\frac{\pi}{2} + \frac{\pi}{4}\right] = (-1)^k$
 For $n=k+1$, need to prove
 $\tan\left[\frac{\pi}{2} + \frac{\pi}{4}\right] = (-1)^{k+1}$

LHS = $\tan\left[\frac{\pi}{2} + \frac{\pi}{4}\right] = \tan\left[\frac{\pi}{2} + \frac{\pi}{4} + \frac{\pi}{4}\right]$
 $= \tan\left[\frac{\pi}{2} + \frac{\pi}{2}\right] = \tan\left[\frac{3\pi}{2}\right] = -1$
 $= -\cot\left[\frac{\pi}{4}\right] = -1$
 $= -\frac{1}{\tan\left[\frac{\pi}{4}\right]} = -\frac{1}{1} = -1 =$ RHS
 True for $n=k+1$ if true for $n=k$
 True for $n=1$, hence true for $n=2, 3, 4, \dots$
 \therefore True for $n \geq 1$.




For the slower stone
 $y_1 = -g$
 $y_1 = -gt + ut \sin \alpha$
 $y_1 = -\frac{1}{2}gt^2 + ut \sin \alpha$
 Similarly for the faster stone
 $y_2 = -\frac{1}{2}gt^2 + vt \sin \alpha$, $x_2 = vt \cos \alpha$
 At time t , let the stones be at A & B.

$M_{AB} = \frac{y_2 - y_1}{x_2 - x_1}$
 $= \frac{-\frac{1}{2}gt^2 + vt \sin \alpha - (-\frac{1}{2}gt^2 + ut \sin \alpha)}{vt \cos \alpha - ut \cos \alpha}$
 $= \frac{t \sin \alpha (v - u)}{t \cos \alpha (v - u)} = \tan \alpha$

(iii) When the faster stone is at Q, the top of the wall, the slower stone is at P.
 $M_{PQ} = \tan \alpha$, $\therefore \angle QPR = \alpha$

$\tan \alpha = \frac{QR}{PR} = \frac{h}{PR}$
 $\therefore PR = \frac{h}{\tan \alpha} = h \cot \alpha$
 (iii) When $y_1 = 0$,
 $-\frac{1}{2}gt^2 + ut \sin \alpha = 0$
 $gt^2 = 2ut \sin \alpha$
 $t = \frac{2u \sin \alpha}{g}$ ($t \neq 0$)

At this instant, the faster stone is at Q.



$\tan(\alpha + \beta) = \frac{-gt + V \sin \alpha}{V \cos \alpha}$
 $= \frac{\beta (2u \sin \alpha)}{V \cos \alpha} - V \sin \alpha$
 $= \frac{\sin \alpha (2u - V)}{V \cos \alpha} = \frac{\tan \alpha (2u - V)}{V}$
 $V \tan \beta = \tan \alpha (2u - V)$
 $\tan(\alpha + \tan \beta) = 2u \tan \alpha$
 (iv) If $\beta = \frac{1}{2}\alpha$,
 $\tan \alpha = \frac{2 \tan \beta}{1 - \tan^2 \beta}$
 $V \left(\frac{2 \tan \beta}{1 - \tan^2 \beta} + \tan \beta \right) = 2u \left(\frac{2 \tan \beta}{1 - \tan^2 \beta} \right)$
 $2V \tan \beta + V \tan \beta (1 - \tan^2 \beta) = 4u \tan \beta$
 $2V + V(1 - \tan^2 \beta) = 4u$
 $u = \frac{1}{4}V(3 - \tan^2 \beta)$
 $< \frac{3}{4}V$ ($\because \tan^2 \beta > 0$)

(b) $v_x = k(b-g)$
 $\uparrow m\ddot{x}$ $\downarrow \frac{mv}{k}$ (i) $m\ddot{x} = -mg - \frac{mv}{k}$
 $\ddot{x} = -\left(\frac{kg+V}{k}\right)$
 (ii) $\frac{dv}{dt} = -\left(\frac{kg+V}{k}\right)$
 $\frac{dt}{dv} = \frac{-k}{kg+V}$
 $t = -k \int_v^V \frac{1}{kg+V} dV$
 $= -k \left[\ln(kg+V) \right]_v^V$
 $= -k \left[\ln(kg+V) - \ln(kg+v) \right]$
 $= -k \ln \left(\frac{kg+V}{kg+v} \right)$
 $= k \ln \left(\frac{kg+v}{kg+V} \right)$
 $= k \ln \left(\frac{hk}{kg+V} \right)$
 (iii) Max. height when $v=0$.
 $t = \ln \left(\frac{kh}{kg} \right) = k \ln \left(\frac{h}{g} \right)$
 $v \frac{dv}{dx} = -\left(\frac{kg+V}{k}\right)$
 $\frac{dx}{dv} = \frac{-kv}{kg+V}$
 $x = -k \int \frac{V}{kg+V} dV$
 $= -k \int \left(1 - \frac{kg}{kg+V} \right) dV$
 $= -k \left[V - kg \ln(kg+V) \right] + C$
 When $x=0$, $v = kh - kg$
 $0 = -k \left[kh - kg - kg \ln(kg+kh-kg) \right] + C$
 $C = (kh - kg - kg \ln(kh))k$

Q8(b)(iii) Cont.
 $x = -k \left[V - kg \ln(kg+V) \right] + k \left[kh - kg - kg \ln(kh) \right]$
 $= k \left[kh - kg - V + kg \ln \left(\frac{kg+V}{kh} \right) \right]$
 At $x=H$, $v=0$.
 $H = k \left[kh - kg + kg \ln \left(\frac{kg}{kh} \right) \right]$
 (c) (i) $P(x) = (x^2 - a^2)Q(x) + px + q$
 $P(a) = pa + q$ (1)
 $P(-a) = -pa + q$ (2)
 (1) - (2)
 $P(a) - P(-a) = 2pa$
 $\therefore p = \frac{1}{2a} [P(a) - P(-a)]$
 (1) + (2)
 $P(a) + P(-a) = 2q$
 $\therefore q = \frac{1}{2} [P(a) + P(-a)]$
 (ii) $x^n - a^n = (x^2 - a^2)Q(x) + px + q$
 $P(x) = x^n - a^n$
 When n is even $P(a) = 0$
 $P(-a) = 0$
 $\therefore q = 0, p = 0$
 When n is odd
 $P(a) = 0$, $P(-a) = -2a^n$
 $px + q = \frac{1}{2a} [2a^n]x + \frac{1}{2}(0 - 2a^n)$
 $= a^{n-1}x - a^n$