

# Sydney Boys High School

## 4 unit mathematics

### Trial HSC Examination 1994


1. (a) Find  $\int \frac{2x}{1+x^4} dx$   
(b) Evaluate  
(i)  $\int_1^2 \frac{dx}{x^2-x+1}$   
(ii)  $\int_0^{\frac{\pi}{2}} \frac{dx}{2+\cos x}$   
(iii)  $\int_0^1 e^{\sqrt{x}} dx$   
(c) Prove that  $\int_a^b f(mx) dx = \frac{1}{m} \int_{ma}^{mb} f(x) dx$  ( $m \neq 0$ )
2. (a) (i) If  $z = 3 - 4i$ , express  $z^2$  and  $\frac{1}{z}$  in the form  $a + ib$  where  $a$  and  $b$  are real numbers. Represent them on an Argand diagram.  
(ii) If  $w^2 = z$ , express the two values of  $w$  in the form  $a + ib$ .  
(b) Illustrate on the Argand diagram the region  $\{z : 0 \leq \arg(z+4) \leq \frac{2\pi}{3} \wedge |z+4| \leq 4\}$   
(c) Let  $z_1, z_2$  and  $z_3$  be three complex numbers represented by  $Z_1, Z_2$  and  $Z_3$  respectively where  $z_1 \times z_3 = (z_2)^2$ . Show that  $OZ_2$  bisects  $\angle Z_1OZ_3$ .  
(d) A sequence  $z_1, z_2, z_3, \dots$  satisfy  $z_{m+1} = z_n^2 + z_1$  for all  $n \geq 1$ . If  $z_1 = i$ , find the distinct values that occur in the sequence.
3. (a) A hyperbola has equation  $9x^2 - 16y^2 = 144$   
(i) Prove that the eccentricity is  $\frac{5}{4}$ .  
(ii) Find the coordinates of the foci.  
(iii) Find the equations of the asymptotes.  
(iv) Find the length of the latus rectum. (A latus rectum is a focal chord parallel to the directrix.)  
(v) Find the equation of the normal to the hyperbola at the point  $(\frac{4\sqrt{10}}{3}, 1)$   
(b) The tangents at the two points  $P$  and  $Q$  with parameters  $\theta$  and  $\phi$  respectively on the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  intersect at the point  $T$ . Show that  
$$T = \left( \frac{a(\sin \theta - \sin \phi)}{\sin(\theta - \phi)}, \frac{b(\cos \phi - \cos \theta)}{\sin(\theta - \phi)} \right).$$
  
(c) Find the condition for the line  $px + qy + r = 0$  to be a tangent to the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ .
4. (a) Sketch, on separate diagrams, the curves  
(i)  $y = \frac{x-4}{x}$  (ii)  $y = \frac{x^2-16}{x^2}$ .




(b) (i) A particle  $P$  is projected, from a point  $O$  on the horizontal ground, with speed  $V$  at an angle  $\theta$  above the horizontal, where  $\tan \theta = \frac{1}{3}$ . The particle passes through the point with coordinates  $(3a, \frac{3a}{4})$  relative to the horizontal and vertical axes at  $O$  in the plane of motion. Show that  $v^2 = 20ga$ .

(ii) A particle  $Q$  is projected, from a point  $O$  at the instant when  $P$  is moving horizontally. It strikes the ground at the same place and at the same instant as  $P$ . Show that the speed of projected of  $Q$  is  $\sqrt{\frac{145ga}{2}}$  and find the tangent of the angle of projection.

8. (a)



Mr Keating's crystal ball rests on a solid stand which is in the shape of a square based frustum as shown.



The stand is constructed such that the crystal ball of radius  $R$  fits snugly inside and just touches the centre of the square base. The sides,  $EC$ , of the base slope so that, if extended, they would pass through the top-most point on the ball at  $A$  and make an angle  $\theta$  with the vertical  $AD$ .

(i) Show that  $OB = R \cos 2\theta$

(ii) Consider a slice  $PQR$  of thickness  $\Delta x$  as shown taken perpendicular to  $AD$  such that  $OQ = x$  units. Draw a neat sketch of the slice, determine its dimensions and show that it has a volume  $\Delta V$  given by  $\Delta V \approx [4 \tan^2 \theta (R + x)^2 - \pi(R^2 - x^2)] \Delta x$

(iii) Find the volume of such a solid where the angle  $\theta = \frac{\pi}{6}$ .